form frequently start with the usual symptoms, and thus a different variety becomes established imperceptibly.

What benefit is to be derived from describing epidemic catarrhal jaundice as a separate clinical entity? It is stated that outbreaks of the disease often accompany an increased incidence of diarrhoeal disorders and that the initial symptoms may be of the "influenzal" type. Only one form of the disease is described by this name, and therefore it is misleading. These facts surely support the adoption of a wider conception of influenza.

Summary

It is suggested that the application of the term "influenza" be generic; that the names "epidemic catarrhal jaundice" and "infective hepatitis" are unsatisfactory, and that "influenzal jaundice" or "hepatitis" would convey a clearer conception of the disease.

A case of acute lymphocytic meningitis occurring in influenzal jaundice is described.

I wish to thank Dr. P. F. Wilson for his permission to publish this case, and the pathologist, who desires to remain anonymous, for allowing me to include his reports on the cerebrospinal fluid.

PSEUDOSCLERODERMIA

BY

R. B. McMILLAN, B.Sc., M.D., M.R.C.P.Ed.

Physician-Superintendent, Eastern General Hospital, Edinburgh

The following case of pseudoscleroderma arising from multiple deficiency may be of interest owing to the rarity of the condition.

Case History

A man aged 63 was admitted to hospital on July 15, 1941, complaining of stiffness in his legs, which prevented him from walking. He stated that he had been perfectly well until approximately one month before admission, when his right knee-joint suddenly swelled and became painful, forcing him to rest in bed. This swelling was attributed to a slight injury received to the same knee two days earlier which caused only momentary discomfort. After a few days in bed he noted that the skin of both his legs below the knee was becoming darker in colour than usual, the right leg being more involved than the left. This discoloration extended, and within the next seven to ten days it had involved the skin all over his body, being most pronounced on the face and neck, the backs of the hands to the wrists, and the legs from mid-thigh down. During this time he had been rising to attend to his own wants, but about 14 days before admission he began to have difficulty in straightening his legs at the knees. This stiffness progressed, and for some days before admission he was confined to bed with his knees fixed in a position of semiflexion. He stated that he had been feeling tired for an indefinite period, and that for the past month his bowels had been loose. There was no history of sore tongue or of Raynaud's syndrome.

The patient was unemployed, and his total income was 16s. 6d. a week from public funds. His expenditure was 4s. 6d. a week for his bed in a lodging-house, 2s. for tobacco, and the rest for food, etc. He did not drink alcohol. He cooked for himself, and lived mainly on pies, sausages, white bread, and scones, never eating fruit or any vegetables except a very occasional potato, and using condensed milk and margarine in place of fresh milk and butter. There was no previous illness of any note.

Clinical Condition

Routine examination revealed a small, thin, wasted man with a light-sepia-coloured skin with uniformly darker areas on the face and neck, backs of hands, and legs from mid-thigh to mid-dorsum of the foot. There was no special pigmentation around the nipples, on the pubes, or on points where pressure from clothing normally occurs. He was edentulous, had pale mucous membranes, a dry glazed but not inflamed tongue, atrophic ribbed finger-nails, pure white luxuriant hair, and patches of brown pigmentation on his buccal mucosa. There was no circumcorneal injection or cheilosis. Both corneae appeared healthy, and there were no petechiae or evidences of deeper haemorrhages. He appeared comfortable, but was slightly inattentive and confused. The skin around both his legs from 6 in. to 9 in. above the knee-joints to about the middle of the dorsum of his feet, but not on the soles, was brown in colour, thick, hard, and glossy to touch, and could not be moved on the underlying tissues. The skin elsewhere, although discoloured, was of normal consistency and mobility. The thickened skin was raised slightly above the normal skin, and the junction between was sudden (within 1/4 in.) and uneven. The legs were

not swollen and were not tender on pressure. Sensation over the affected area was dulled but not entirely absent. Both legs were held flexed at 45 degrees at the knees, and movement at these joints, which were not swollen, was painless but limited to a range of 45 to 60 degrees flexion by the tightness of the surrounding skin. Movement at the ankle-joints was also slightly limited. Movement at all other joints was normal. Further examination revealed no physical defect, and the remaining points of particular interest which helped in making a diagnosis are:

Skin temperature steady at 97.4°; pulse 75 to 80 a minute, regular; blood pressure 145/80. Test-meal (including histamine): no free HCl. Faeces 4 to 6 soft pale motions daily; total fat 33.9 g.—split fat 27.2%, unsplit fat 72.8%. Urine: routine tests negative; ascorbic acid (estimated as reducing substance), nil; creatin, 14 mg. per 100 c.cm. Blood: W.R. negative; plasma proteins, normal; urea, etc., normal; B.S.R. (Westergren), 6 mm. per hour; plasma ascorbic acid, 0.10 mg. per 100 c.cm.; sugar tolerance, normal curve; R.B.C., 3,310,000 per c.mm.; Hb, 38%; C.I., 0.58; W.B.C., 8,200 per c.mm. Film showed poor filling of red cells, anisocytosis and poikilocytosis, and a normal differential white count. Radiographs of knees showed only slight atrophic arthritis. Cuff test (Hess), negative. Microscopical examination of biopsy from affected skin resulted in the following report:

"The tissue consists mostly of vascular scar tissue. Enveloped in it are small foci of adipose tissue and dense fibrous fascial bands; at one side is striped muscle, which, at margin of scar tissue, is partly atrophied, partly hyperplastic (sarcolemmal cells proliferating). There are recent haemorrhages in the scar and also scattered granules of old blood pigment (previous haemorrhages probably). The appearances suggest scar formation in and adjacent to muscle and perhaps accentuated by haemorrhage."

Diagnosis

On the history and findings recorded above a multiple deficiency state was diagnosed, obvious deficiencies being nicotinic acid, ascorbic acid, and iron. The general pigmentation was considered to be part of the pellagroid change, and the thickened skin on the legs to be the result of this plus very minute multiple haemorrhages, the result of the vitamin C lack. The probability that other deficiencies aided in producing the curious localized change in the legs was great, but it was impossible to estimate their part in doing so. The diagnosis was made, therefore, of pseudoscleroderma the result of vitamin deficiency, and support was lent to this by the earlier records of a precisely similar change occurring in the legs of old people with scurvy (Pfeiffer, 1918). Other causes of the changes that were considered and excluded on the facts which have been given were: (a) true symmetrical diffuse scleroderma (sclerodactylia); (b) chronic cellulitis of legs with, as superadded causes of the general pigmentation, Addison's disease, haemochromatosis, and chronic vermin infestation of the skin. The presence of creatine in the urine was considered normal at the patient's age.

Treatment

Treatment was designed to test this diagnosis, and was conducted in various stages. The first stage was the maintenance of the patient for a period on the ordinary hospital diet, in order that its action could be assessed; the successive stages consisted in the addition to this diet of the obviously absent vitamins, in order that their action in promoting cure could be assessed also. Ascorbic acid was replaced first, mainly because the condition had been described before in association with scurvy.

Table showing Results of Treatment

Clinical Feature	Stage I: 18/7/41 to 18/8/41 Ordinary Hospital Diet	Stage II: 18/8/41 to 8/10/41 Ord. Diet + 7,000 mg. Vit. C in 14 Days, then 200 mg. Daily by Mouth	Stage III: 8/10/41 to 24/1/42 Ord. Diet + 200 mg. Vit. C + 200 mg. Nicotinic Acid Daily by Mouth	Stage IV: 24/1/42 to 9/3/42 As Stage II + 1,000 I.U. Vit. B.I.M.I. Daily
Blood { R.B.C. Hb ..	3,970,000 50%	4,150,000 66%	5,000,000 75%	5,000,000 79%
Bowels	Stools still frequent and soft As on admis- sion	Improved but not normal	Normal	Normal
Mental state		Slight improve- ment	Marked im- provement to normal	Normal
Pigmentation	" "	Slight improve- ment	Marked im- provement	Normal
Skin thickening	" "	Very slight improvement; smaller area involved Knee move- ment 35°-60° flexion	Great improve- ment: small slightly thick- ened area on skin only re- mains Knee move- ments full and free	Completely healthy skin

The accompanying table shows the details of each stage of treatment, its duration, and the changes in the clinical features of the case as each stage was concluded.

On completion of treatment the initial tests were repeated, and it was found that all those previously abnormal were now normal. Creatine in small amounts was still being excreted in the urine. The patient was discharged well on May 1, 1942, and when last seen (Oct., 1942) was still free from recurrence of the skin lesion.

Comment

The little change shown in the skin, etc., of the legs when only ordinary diet was given, and the marked improvement ending in complete cure when the diet was supplemented and the existing deficiencies were corrected, suggest that the diagnosis of pseudosclerodermia arising from multiple vitamin deficiency was correct. According to the table the most marked improvement occurred when nicotinic acid was given, suggesting that it was lack of this factor which was mostly responsible for the condition. This may be so, but whether nicotinic acid would have had such a marked action in the absence of adequate vitamin C nutrition is doubtful, as in multiple deficiencies vitamins are notoriously interdependent for their action. It is only possible to suggest that the combination of deficiency of nicotinic acid, ascorbic acid, and thiamin, in that order of importance, precipitated the pseudosclerodermia.

The slow response to treatment was not considered abnormal, as this is often a feature in multiple-deficiency states, and particularly so in lesions affecting the skin (O'Leary, 1942). Quicker results could have been obtained had all the missing factors been supplied together. The failure of the lesion to respond to replacement of vitamin C only, plus the failure of the author to observe any condition really closely resembling sclerodermia—although skin thickening was often seen—in 58 old people with clinical scurvy observed over the past five years suggests that the lesion noted in association with scurvy by Pfeiffer, if really resembling sclerodermia as did the case described above, was the result also of multiple deficiency, vitamin C not being the most important lack.

Changes faintly resembling certain stages of true sclerodermia may result from vitamin deficiency and are not uncommonly found. The changes seen and the factors responsible are: (a) subcutaneous oedema, causing apparent skin thickening and fixation and muscle tenderness from lack of vitamin B (Eddy and Dalldorf, 1941); (b) atrophy of epidermis, papillary oedema, papillary capillary dilatation, slight oedema of the dermis, and pigmentation from lack of nicotinic acid (Eddy and Dalldorf, 1941); (c) subcutaneous haemorrhages causing thickened skin and pigmentation due to lack of vitamin C; and (d) atrophy of muscle fibres in experimental animals, due to lack of vitamin E (Pappenheimer, 1939; O'Leary, 1942).

Other records of pseudosclerodermia of nutritional origin are few. Reports of cases which might have had this origin are: (a) a lesion similar to that described in this paper in a patient in association with Sjogren's syndrome (Sheldon, 1939); and (b) a lesion in a poorly nourished woman in whom cure resulted when a good diet was given (Meakins, 1926). It is suggested that it might be more common than is thought, and, as support, attention is drawn to the reports of sclerodermia appearing in patients debilitated by cancer of various organs, in some of whom it disappears when the cancer has been treated and the general nutrition improved, good food being noted to be particularly beneficial in one instance (Behrman and Forman, 1939; Bezecky, 1935).

It is not suggested that vitamin deficiency is the cause of true sclerodermia, and a search has failed to reveal any example of vitamin deficiency or deficiencies causing a change which microscopically resembles the true condition. As the clinical resemblance may be very great the importance of a biopsy in making a diagnosis of sclerodermia is obvious, as without it mistakes could be made and the patient given an entirely wrong prognosis and treatment.

Lastly, although it is not intended to discuss vitamin C in general, the need of this substance by healing wounds (Crandon *et al.*, 1940) is well shown by the behaviour of the biopsy in this case. The wound made on Aug. 3, 1941, when the patient was subcorbutic, showed little healing by Aug. 18, after which date the rate of healing accelerated markedly, the wound being

firmly closed by Aug. 27. This acceleration corresponded to the addition to the patient's diet of 500 mg. of vitamin C daily by mouth, and was obviously governed by it, as all other factors were unchanged.

REFERENCES

- Behrman, S., and Forman, L. (1939). *Proc. roy. Soc. Med.*, **32**, 1575.
 Bezecky, R. (1935). *Arch. Derm. Syph.*, Berlin, **171**, 223.
 Crandon, J. H., Lund, C. C., and Dill, D. B. (1940). *New Engl. J. Med.*, **223**, 353.
 Eddy, W. H., and Dalldorf, G. (1941). *The Avitaminoses*, 2nd ed., Baltimore and London.
 Meakins, J. (1926). Quoted by McCallum, W. G., *Trans. Ass. Amer. Phys.*, **41**, 190.
 O'Leary, P. A. (1942). *Arch. Derm. Syph.*, Chicago, **46**, 628.
 Pappenheimer, A. M. (1939). *Amer. J. Path.*, **15**, 179.
 Pfeiffer, L. R. (1918). *Disch. med. Wschr.*, **44**, 625.
 Sheldon, J. H. (1939). *Proc. roy. Soc. Med.*, **32**, 255.

PNEUMONIA TREATED WITH SULPHAMETHAZINE

REPORT OF 77 CASES

BY

B. A. PETERS, M.D., D.P.H.

Medical Superintendent, Ham Green Hospital, Bristol
AND

M. L. EASBY, M.B., Ch.B.

Assistant Medical Officer, Ham Green Hospital, Bristol

During the latter part of the past winter we have treated 77 cases of pneumonia with sulphamethazine (2-sulpha-4:6-dimethyl pyrimidine), one of the latest additions to the sulphonamide group. It is built up on the same plan as sulphadiazine, with the addition of two methyl groups to the pyrimidine ring. This addition is claimed by the makers (I.C.I.) to increase the solubility of the drug and render its deposition in the kidneys unlikely. This effect was a serious disadvantage of sulphadiazine.

Of the 51 bronchopneumonias treated in the under-12 group 30 were secondary to measles, 2 to whooping-cough, and the rest primary. The ages varied from 4 weeks to 85 years. The normal dosage of sulphamethazine employed, as recommended by the makers, is 8 g. daily for an adult, given six-hourly in 2-g. doses. As children take these drugs better than adults we assumed the adult weight for the standard dose to be 8 st., and gave a fraction of this dose according to their weight—that is, a 4-st. child received half the adult dose and so on. The full dose was given for four days, and half the dose for a further two days. Plenty of fluids were administered, and oxygen tents were used as considered necessary.

We were unfortunately unable to estimate the blood concentration with these doses. All the cases responded more promptly to the drug the earlier in the disease it was given. We think the response in cases responding favourably is slower than with sulphapyridine, but the mortality was low, as only three died (4%). One 13-week-old baby with whooping-cough and bronchopneumonia was admitted on the sixth day and died four days later. Another fatal case had measles and bronchopneumonia, was admitted on the sixth day of disease, and died on the fifth day after admission. One patient with extensive bronchopneumonia—a female aged 26—survived only 12 hours after admission. Nine other cases showed no obvious response to the drug within six days, but eventually recovered.

Out of 77 patients, therefore, 65 appeared to respond and 12 did not; three of the latter died. Of those not responding, 6 cases in adults appeared to be aberrant cases of lobar pneumonia. From one of the children who did not respond a nearly pure culture of *Staph. aureus* was found in the sputum, and she developed a staphylococcal pyaemia. Not one case in the series developed an empyema.

Toxic Effects

Owing to the risk of renal blockage and urinary suppression with the related sulphadiazine, a careful watch was made on renal secretion. One child aged 2 showed almost complete sudden urinary suppression for one day. On stopping the drug the renal flow was restored within 24 hours. No crystals