sterilization. This can then be performed when the heart has not the added burden of the placental circulation with which to cope.

One complication of the method must be mentioned. A spinal headache seems to occur much more frequently in pregnant than in non-pregnant patients. The headache often lasts for as long as a week. Several methods have been tried both to prevent and to relieve this spinal headache; but none of them is uniformly successful, and these methods are beyond the aims of this paper.

I wish to thank Mr. C. Gordon Lewis, F.R.C.S.Ed., medical superintendent, and Dr. J. Lindsay Boyd, consulting physician, of the Stoke-on-Trent Hospital, for their help and consideration.

EFFECT OF DIET ON THE CONCENTRATION OF CHOLESTEROL IN BLOOD AND BILE

BY

NANCY GOUGH, B.Sc.

(From the Clinical Laboratory, Royal Infirmary, Edinburgh)

It is widely believed that the level of cholesterol in the blood and bile is influenced by the intake of cholesterol in the food, and for this reason the diet of patients with disorders of the biliary system has commonly been restricted to food low in cholesterol. The following investigations suggest that this irksome restriction may be unnecessary, since the concentrations of cholesterol in the blood and in the bile do not appear to be significantly affected by the ingestion of comparatively large amounts of cholesterol.

Effect of High- and Low-cholesterol Diets on the Blood Cholesterol

Hospital patients whose metabolism was supposedly normal were fed on low-cholesterol diets containing about 300 mg. of cholesterol a day for a fortnight, and blood specimens were analysed for cholesterol at intervals of three days, all blood samples being taken at the same time of day. Then the cholesterol content of the diet was increased by the daily addition of 900 mg. of pure cholesterol, three eggs (approximately 1,100 mg. of pure cholesterol a day), or 50 g. of sheep's brain (approximately 2,170 mg. cholesterol a day), the cholesterol content of the blood again being determined at intervals of three days.

FIG. 1.—Relation between cholesterol content of diet and concentration of cholesterol in plasma. The low-cholesterol diet contained approximately 300 mg. cholesterol and the high-cholesterol diet approximately 2,170 mg. a day, given as 50 g. of brain.

Of 10 such cases in which the estimations were made by the colorimetric method of Myers and Wardell, 8 showed the plasma cholesterol unaffected by diet, while in 2 cases the plasma cholesterol was definitely raised by the addition of

900 mg. of cholesterol a day to the basic diet, and also by the daily addition of three eggs. Of another 10 cases, in which the estimations were made by the digitonin-precipitation method of Okey, in no instance did a high-cholesterol diet result in elevation of the plasma cholesterol. The accompanying graph shows a typical result in a normal person (Fig. 1).

Cholesterol Absorption

Experiments on the effect of a single dose of cholesterol on the blood cholesterol were carried out on four normal subjects, who had been fasting for twelve hours before each experiment. The estimations were made by Okey's digitonin-precipitation method. A cholesterol-free, fat-free meal supplemented by 5 g. of pure cholesterol was given, and a blood cholesterol curve constructed over a period of six hours. The blood cholesterol curves did not differ significantly from those obtained from the same subjects when given the same meal without the added cholesterol. As it was thought possible that the absorption of crystalline cholesterol is dependent on the presence of adequate fat in the intestine, this experiment was repeated, 50 g. of fat as margarine being added to the test meal. Again no significant difference was observed. Fig. 2 shows the results in the case of two of these subjects.

FIG. 2.—Plasma cholesterol curves over a period of six hours. A=Test-meal (cholesterol-free, fat-free). B=Test-meal + 5 g. cholesterol. C=Test-meal + 50 g. fat. D=Test-meal + 50 g. fat and 5 g. cholesterol.

The conclusion from these experiments appears to be that 5 g. of crystalline cholesterol by mouth has no effect in raising the free or esterified cholesterol in blood. Whether these findings can be explained by a failure in absorption of crystalline cholesterol, or by some mechanism whereby it is removed from the blood or altered into some compound not detected by the analytical method employed, is uncertain.

Effect of High- and Low-cholesterol Diets on the Cholesterol Content of the Bile

These experiments were carried out on 10 patients with biliary fistulae. The clinical condition of 4 of the patients did not permit the giving of experimental diets, and they were therefore kept on a light ward diet, which has a fairly constant low-cholesterol content of approximately 300 mg. a day. The remaining 6 patients were given a special low-cholesterol diet containing less than 300 mg. of cholesterol. After a period of two or three days they were given a diet with a high content of cholesterol, as the pure substance or as brain, for a further period of three days. Bile was collected from the fistula on alternate days. The cholesterol content of the bile was invariably very low immediately after the operation, but rose gradually, presumably as the liver damage lessened, although there was considerable day-to-day variation. This variation was most pronounced (10 mg. to 45 mg. per 100 c.cm.) in a case with severe hepatic damage (gross jaundice). There was also a great daily variation in the volume of bile which drained externally.

In human beings with a biliary fistula it is impossible to determine what proportion of the bile is draining externally and what is draining into the intestine via the common duct; hence no calculation can be made of the total secretion of bile

or cholesterol. But it is possible to compare the concentration of cholesterol in a given sample of bile with the cholesterol ingested in the diet.

Of 10 patients with biliary fistulae, 1 showed a distinct rise in the bile cholesterol concentration on a diet containing a high content of cholesterol (about 2,000 mg., given as 50 g. of sheep's brain a day); 1 a rise when the high-cholesterol diet was first given, with a subsequent return to the original level (the patient still on the high-cholesterol diet); and 2 a fall in the cholesterol concentration of the bile on a low-cholesterol diet. In 6 no significant or sustained changes in the bile cholesterol concentration coincided with changes in the cholesterol content of the diet. The results in one of these six cases are shown in Fig. 3. The duration of these experiments was limited by the short period of time during which the fistulae remained open.

FIG. 3.—Concentration of cholesterol (free) in bile of a patient with biliary fistula. The low-cholesterol diet contained approximately 300 mg. a day, and the high-cholesterol diet 2 g. given as the pure substance.

To overcome the difficulty mentioned above with regard to determining the total output of cholesterol in the bile recourse must be had to operations on animals. Prof. Learmonth inserted a tube into the common duct of a dog so that all the bile drained externally. The animal remained in excellent condition for one month, the absence of bile in the intestine being compensated by giving 1 g. of sodium taurocholate by mouth every day. The total volume and the cholesterol concentration of the bile were measured daily; for the first three days insufficient bile was secreted for analysis, but thereafter it increased, although there was considerable fluctuation both in the volume and in the cholesterol content. After about ten days these became fairly constant (concentration of cholesterol in bile 13 to 14 mg. per 100 c.cm., and total bile cholesterol excreted about 20 mg. a day). Approximately 2 g. of cholesterol was then fed in the form of 50 g. of sheep's brain, and there was an immediate rise in the bile cholesterol (27 mg. per 100 c.cm. and total bile cholesterol 38 mg.), which was maintained for 48 hours; but subsequently both concentration and total output of cholesterol in the bile returned to the original level, although the high cholesterol content of the diet was maintained.

Conclusion

Further experimental work on animals is obviously required before any definite conclusion can be reached as to the effect of diet on the total output of bile and cholesterol, but the evidence of the foregoing experiments on human beings seems to indicate that there is no direct correlation between the intake of cholesterol and its level in the blood and bile. These observations give no support to the contention that foods rich in cholesterol should be prohibited in the diet of patients with cholecystitis and cholelithiasis. The exclusion of butter, fat, eggs, and liver from the diet of patients with diseases of the biliary tract is based essentially on the hypothesis that gall-stones are formed from cholesterol, and that a reduction in the intake of cholesterol accordingly reduces the liability to

gall-stone formation. Such a low-fat, low-cholesterol diet is unphysiological because fat is the natural stimulant to biliary contraction and drainage, and has the further disadvantage of being unpalatable, monotonous, and low in fat-soluble vitamins.

My thanks are due to Prof. L. S. P. Davidson and Dr. C. P. Stewart, under whose direction this work was done, and to Prof. J. R. Learmonth, who carried out the necessary animal surgery. Part of the expenses were defrayed by a grant from the Medical Research Council to Prof. Davidson.

Medical Memoranda

Contramine for Herpes

Herpes is very prevalent in this locality and is often of great severity; in some cases the acute stage has lasted for months and the after-effects for years. Hence my satisfaction when I stumbled on a specific which provided a rapid cure of this painful malady. By means of this treatment, if given early, the acute pain is often relieved in a matter of hours, the vesicles disappear in a week or so without pitting, and there are no after-effects.

Some time ago, while attending an elderly lady for rheumatoid arthritis, I gave a course of injections of contramine. When visiting her to administer the second injection she complained of acute pain in the right supraorbital region. The next day I had an urgent summons, and found a very extensive herpes rash present. To my surprise this completely disappeared in five days. On the assumption that the contramine cured rather than caused the herpes, I have in the last twelve months treated 15 cases by this method.

The results may be summarized as follows: Average duration of pain before case diagnosable, 4 days; average time between irruption and injection, 16 hours; average duration of irruption, 7½ days. In a few cases among elderly patients of the residential class there was some neuritic pain for a further week after the skin was normal.

The efficacy of the treatment was most pronounced in those cases in which the injection was given within a few hours of the first appearance of the spots, especially in the pre-vesicular stage. The following cases may serve as examples, taken in more detail:

REPORTS OF CASES

Case 1.—On Sept. 5, 1942, a single lady aged 70 consulted me about an acute pain which she referred to the eye. On the following evening I received an urgent message to visit her, as she had fainted with the pain, which was of a spasmodic character. I then arranged with a consultant ophthalmic surgeon to see her the following afternoon. The next morning I witnessed an acute attack severe enough to make her pulseless. By this time I suspected an acute herpes, which was confirmed by the consultant in the afternoon. Contramine was then given, with the result that there was no further pain, nor did any rash develop, and apart from a few days' weakness the patient was well.

Case 2.—On Dec. 24, 1942, a man aged 76 complained of violent occipital neuritis, and two days after this a slight rash appeared from occiput to orbit. Contramine was given, and the vesicles never fully developed but formed dry scabs. This would undoubtedly have been a very severe case, but he was well in a fortnight.

Case 3.—A working man aged 52 complained of an increasingly severe pain in the chest and loss of sleep for two nights. His work was hard, and as it was essential he wished to carry on, but doubted whether he could. On examination an early shingles rash was found over the whole of the left chest. An injection was given, and two days later he reported that the pain had ceased entirely in four hours. All symptoms had gone in four days.

In five cases drawn from the working-class similar results were obtained—much better than in elderly, and wealthier, patients.

ITS USE IN CHICKEN-POX

I have reason to believe that the treatment is equally effective for chicken-pox. Of the three cases in which I have used it details of one may be of interest—the only early case.

A boy aged 16, living at a hostel for evacuees, most of whom had not had this complaint, had an attack with a severe onset, and was injected with contramine 12 hours after the appearance of the rash. On the third day, being quite well, he was allowed up, did gardening, and mixed with the other children. None of these were infected.

COMMENT

The injection intramuscularly of 1 c.cm. (0.125 g.) of contramine (B.D.H.) gives a dramatic result in early cases of herpes, and is of great value in chicken-pox. In no case was pitting or