bear any relation to the frequency of fits. Blood pressure in these patients has shown little variation since the onset of their fits. Other evidences of cerebral arteriosclerosis—for example, strokes and mental deterioration—are notably absent. Only one case showed marked mental changes when he first came under observation; this condition improved with the response of the fits to treatment, and the man has been back at work for some years.

Jacksonian Epilepsy

The question arises whether Jacksonian epilepsy is especially suggestive of tumour. Among the total of 81 patients with onset over the age of 40, 7 are known to have had Jacksonian attacks (Table VI). Of these, 6 are dead, the cause being shown.

TABLE VI.—Significance of Jacksonian Epilepsy among 81 Patients

Number known to have Jacksonian attacks	7
Died	6
Cerebral arteriosclerosis	2
G.P.I.	1
Heart failure	2
Unknown cause	1

The mortality rate is probably exceptional, but the point which emerges is that Jacksonian attacks, while revealing the site of the disturbance, give no indication of the nature of the lesion.

Time of Epileptic Attack

A last point of clinical significance is the time of epileptic attacks. Among the 41 surviving patients of the last group the time of attack was: nocturnal only, 13; diurnal only, 16; irregular, 12. The purely nocturnal incidence of attacks has two important advantages: first, such attacks, occurring in bed, are unassociated with danger of injury, and the patient is able to lead an ordinary life by day; secondly, there is a better chance of controlling nocturnal attacks by maximum doses of an anti-epileptic remedy given at night.

Summary and Conclusions

The age of onset of epilepsy is considered in a series of 602 cases.

81 patients in whom epileptic attacks began after the age of 40 were followed up, and the results of the inquiry are recorded.

Making full allowance for errors in diagnosis, it is clear that the majority of these patients who began to have fits after the age of 40 could not be the subjects of cerebral tumour, severe cerebral arteriosclerosis, G.P.I., or other progressive disease. Most of them continue to have epileptic attacks from time to time without other evidences of disease.

These observations provide no contribution towards the aetiology of idiopathic epilepsy, but they suggest some conclusions which may be of practical value: (1) According to my experience idiopathic epilepsy is by no means uncommon in persons of mature years. (2) Though fits may be the first symptom of intracranial tumour it is rare for them to precede all other symptoms and signs by months or years. (3) In the later years of life epilepsy has remarkably little effect on intellectual powers and general health.

I am interested in this matter from the point of view of teaching. I would suggest that students should be advised always to regard epilepsy as a symptom and to search for the cause both in the nervous system and elsewhere; and, secondly, to try to discover the site of the disturbance by careful analysis of the aura or earliest stage of the fit; but that these rules should apply to epilepsy at any age, and we should not stress unduly the incidence of progressive disease as a cause of epilepsy in the later years of life.

REFERENCES

- Christian, H. A. (1942). *Osler's Principles and Practice of Medicine*, 14th ed., p. 1363, Appleton, New York.
Gowers, W. R. (1901). *Epilepsy*, London.
Penfield, W., and Erickson, T. C. (1941). *Epilepsy and Cerebral Localization*, Baillière, Tindall and Cox, London.
Turner, W. A. (1907). *Epilepsy*, London.

THE SCOPE OF PREVENTION IN OPHTHALMOLOGY

BY

IDA MANN, D.Sc., F.R.C.S.

(Nuffield Laboratory of Ophthalmology, Oxford)

The recent publication by Marshall and Seiler* of their statistical analysis of 3,219 persons certified blind during 1929–35 raises the question of the preventive function of ophthalmology. There is at the present time a total of 74,000 blind persons in England and Wales whose maintenance costs the State £4,500,000 annually, while practically nothing is spent on planned prevention of blindness by improved facilities for treatment, by education of the public, or by research.

Marshall and Seiler's figures bring out certain noteworthy points. First, the seven most important causes of blindness are, in order of occurrence:

Cataract	16.77%	of the total examined
Myopia	16.34%	" "
Vitreous disease	13.95%	" "
Uveitis	10.72%	" "
Glaucoma	8.69%	" "
Congenital anomalies and abiotrophies	7.45%	" "
Injuries	6.42%	" "

Secondly, although at first sight the first four items would suggest that the greatest amount of blindness occurs in the older age groups and does not therefore carry so great an economic importance, yet actually only half the cases became blind after 50, and in two-thirds of these the underlying cause had started before that age. In view of the longer expectation of life now than formerly, and the increasing average age of the population in general, as well as of workers in responsible positions, it is obvious that most of the causes of blindness begin to operate early enough to constitute a risk of economic importance. Thirdly, the relatively low percentage of blindness from injury appears surprising until we realize two points—namely, that the position of injury in the list of causes varies between 10.3 and 5.8%, according to whether the first eye, the second eye, or both, are involved; also, most severe industrial eye injuries lead to loss of one eye rather than both, only those of great severity such as explosions and head injuries appearing to any extent in statistics of blindness, yet the time lost from injuries short of this order is of great economic importance.

Although, because of the certification of blind persons and the payment of pensions to the blind, statistics are readily available, what we lack is a precise knowledge of the amount of working-time lost and of inefficiency and disability caused by eye injuries and diseases which do not end in blindness but which nevertheless should be, if possible, reduced in number. Even if the condition does end in blindness there is a latent period of illness and steadily increasing disability during which the patient is unable to work. This is brought out by Marshall and Seiler's tables, which show the average times between the onset of the cause of blindness and the actual certification. In some cases it is considerable; for example, in all accidents the difference between the mean age of onset and the mean age of blindness is 3.11 years; in glaucoma it is 2.94, in cataract 5.1, and in inflammatory conditions it is higher still (e.g., trachoma 25.58, and uveitis 12.4), while it is highest in myopia (36.24 years). Quite apart from these figures exists the mass of time lost from accidents which recover and from accidents leading to the loss of one eye only, which never come to certification. As pointed out before, these must be borne in mind as explaining the, at first, surprisingly low figures for blindness as a result of injury, and it is to be emphasized that in any discussion on prevention the non-blinding conditions must be taken into account.

When we consider the list of conditions involved from the point of view of possible reduction of eye disease it becomes obvious that advances could occur mainly in three directions—

* Marshall, John, and Seiler, H. E. (1942). *Brit. J. Ophthalm.*, 26, 337, 385, 434.

H. Weyler and C. C. Dustin (*New Engl. J. Med.*, 1942, 227, 785) record their observations on a woman aged 24 who presented paroxysmal auricular tachycardia throughout the latter part of four pregnancies over a period of five years.

namely, by improvement of the ophthalmic services of the country, by education of the public on ophthalmic matters, and by further research into aetiology and therapeutics.

Improvement of the Ophthalmic Services of the Country

From certain of Marshall and Seiler's figures it would appear that some relation exists between the number of blind, especially in the higher age groups, and the hospital facilities of the district. This is to be expected, and could only be countered by making modern ophthalmic hospital treatment equally available for everyone. This would entail geographical planning and co-ordination on a regional basis, together with the provision of a larger number of ophthalmic surgeons than are at present available. Some more standardized system of training for ophthalmologists at recognized centres and a reconsideration of the requisite examination standards would be advisable, as, at present, postgraduate teaching is somewhat haphazard and there is no uniformity of qualifications demanded by hospitals in various parts of the country. Thus the problem is linked with those of regional planning and of medical education. In addition, the vast questions of ophthalmic benefit and the relations between approved societies, ophthalmic surgeons, hospitals, and opticians would also appear to be involved. There is need, too, for a closer supervision and better facilities for diagnosis and treatment of children under school age and of adolescents leaving school but not yet eligible for ophthalmic benefit. At present no provision is made for these groups, and much preventive work might be done on them.

Education of the Public

This is a basic factor in maintaining the national health. One has only to consider the chaos which exists in the public mind on such subjects as the difference between a refractive error and a disease (and, through this, between an optician and an oculist), the value and nature of orthoptic exercises, and the relation between the condition of the eye and the general health, to realize that much could be done by clearing up misconceptions alone. It is well known that many people with glaucoma postpone medical treatment beyond the limit of safety by buying stronger glasses as a makeshift, and that practically all amblyopic eyes in children are due to the belief of parents that the child will outgrow the squint and that young children should not wear glasses. So little is known of the anatomy and physiology of the eye that it is usually difficult to explain to patients such simple problems as the nature of myopia and of cataract. This ignorance paves the way for the development of ocular neuroses and the accompanying vogue for magical or much-advertised systems of treatment. That these are successful in many cases points to the need for a higher standard of diagnosis among ophthalmologists and a higher level of educated co-operation from the patients. Education is needed at three levels. Some biological and physiological instruction in schools should be obligatory; it should include an understanding of the mechanism of sight and the nature of binocular vision and of errors of refraction. In adult life authoritative information should be available on such subjects as the care of the eyes (such as the futility of constant eye baths) and the nature of presbyopia. Most especially, however, instruction is needed on protection of the eyes in industry, since most minor injuries are due to the prevalent attitude towards safety-first measures and to ignorance of the possible effects of such injuries.

Research

The trend of medical research at the present time is towards greater collaboration between experts in different branches of science. Since medicine is largely an art there must be, for progress, close contact with the basic sciences underlying it. Advances in ophthalmology of recent years owe practically everything to basic discoveries in the realms of physics and chemistry—pharmacology and physiology also co-operating. We have only to mention the practical application of the diathermic current and of the cathode ray, the improvement in optical glass and plastics, the synthesis of the sulphonamides, the discovery of the nature of vitamins and their biochemical action, and the researches of psychologists into

the visual function to show how closely interlocked with advances in ophthalmology are discoveries in other fields. Too often, however, the ophthalmologist, overburdened with the mass of clinical work confronting him, lacks the requisite contact with the experimental sciences. The establishment of research teams in association with laboratories where physicists, physiologists, biochemists, and pathologists, among others, could unite with ophthalmologists for the solving of given problems would seem to be desirable. For example, to return to Marshall and Seiler's report and the seven most common causes of blindness, we can foresee that the problems of cataract and glaucoma will require the help of biochemistry for their solution and that uveitis must invoke experimentation in pathology, bacteriology, and immunology. Venereal disease is already yielding to chemotherapy, and myopia and the abiotrophies probably require genetical research for their unravelment. The question of injury includes experiments in the design of protective goggles and work on the best psychological approach to ensure their use.

If these desiderata be kept in mind it is possible that when the next survey of the statistical position can be undertaken a distinct improvement may be manifest.

RISE IN POTASSIUM CONCENTRATION IN THE BLOOD STREAM FOLLOWING ISCHAEMIA OF MUSCLE MASSES

BY

R. E. REWELL, M.D., M.R.C.P.

(From the Department of Pathology, Guy's Hospital Medical School)

It is well known that the concentration of potassium is much greater in cells than in tissue fluids. Thus Eichelberger (1941) found 25 times as much by weight in dog's muscle (corrected for blood and fat content) as in the serum. This great difference in potassium concentration on the two sides of the cell membrane is maintained only so long as this structure remains undamaged. Prolonged asphyxia is one method of causing such damage. Baetjer (1935) found, by measuring its concentration in the perfusate, that loss of potassium from cat's muscle increased as the rate of blood-flow was reduced, but that the actual excitability of the muscle remained unaltered. Fenn *et al.* (1939) found that there was a limit to the potassium reduction in cat's muscle that could be brought about by perfusion. Later, Fenn, Koenemann, and Sheridan (1940) observed that no increased loss of potassium or production of lactic acid followed periods of asphyxia in perfused frog's legs.

Method of Investigation

The present investigation was undertaken to determine whether or not any appreciable loss of potassium followed periods of asphyxia in the larger muscle masses of human beings. The subjects were patients to whose limbs tourniquets had been applied for periods ranging from 35 to 100 minutes in the course of orthopaedic operations. They were all muscular young adults, mostly men. Blood samples were taken from the antecubital vein just before and at seven minutes after the removal of the tourniquet. From the first few patients additional samples were taken—one at the start of the operation and another one hour after its conclusion; but since the former revealed only the usual slight fall in the concentration of potassium that occurs during the administration of an anaesthetic, and the latter a subsequent return to normal, they were soon discontinued. The clot and serum were separated as soon as possible, and any specimen with a trace of haemolysis was discarded. The potassium was estimated by the cobaltinitrite method of Tisdall and Kramer (1921), using duplicate samples and putting through a standard with each batch. In the later cases inorganic phosphate was determined