A CASE OF CONGENITAL DILATATION OF THE PULMONARY ARTERY

BY

H. L. HEIMANN, M.D., M.R.C.P.

Lieut.-Col., S.A.M.C.

AND

M. M. POSEL, M.D., M.R.C.P.

Major, S.A.M.C.

(From Oribi Military Hospital, Pietermaritzburg, Natal)

(WITH PHOTOGRAPHURE PLATE)

According to Bedford, Paff, and Parkinson (1941) congenital dilatation of the pulmonary artery is a rarity, while septal defect is the commonest congenital malformation of the heart. Those authors stress the great dilatation of the pulmonary artery and its branches that accompanies atrial septal defect, and, while attempting to differentiate between these two conditions, confess that differential diagnosis may be impossible. Other authors, notably Assmann (1928), describe this congenital dilatation as a primary condition due to asymmetrical divisions of the truncus arteriosus communis.

Jennes (1936) points out that dilatation of the pulmonary artery is as rare as dilatation of the aorta is common. In a total of 38,000 necropsies done at various English-speaking hospitals no case was discovered up to 1936. In all, 122 cases have been recognized, and, with the exception of 12, all at necropsy. From the descriptions of some of these cases they were probably of the type described by Bedford, Paff, and Parkinson. Master (1939) states that in atrial septal defect large auricles result, with enormous P waves. The opinion expressed by Bedford, Paff, and Parkinson (1941) is contrary to this statement: except in a certain proportion of cases, the right auricle was grossly enlarged, but not the left.

Assmann's classification of congenital dilatation of the pulmonary artery appears to be of value, and is as follows:

A. Acquired :

- (1) Aneurysm of the pulmonary artery
- (2) Perforation of aneurysm of aorta into pulmonary artery
- (3) Pulmonary insufficiency from endocarditis
- (4) Stasis in pulmonary circulation, as in mitral stenosis

B. Congenital :

- (1) Solitary single anomaly (asymmetrical division of truncus arteriosus communis)
- (2) Open ductus botalli
- (3) Pulmonary stenosis (very exceptional)
- (4) Defect of auricular septum
- (5) Defect of ventricular septum

Radiology plays an important part in the diagnosis of this condition, and the criteria of x-ray diagnosis seem to be as follows:

- (1) Increase of pulmonary bow
- (2) Small aortic knuckle
- (3) Broadening of hilar shadows
- (4) Enlargement of the pulmonary arterial tree. The oblique positions, together with the barium swallow, will show the characteristic picture of the enlarged pulmonary artery.

In view of the rarity of the condition, we are adding a description of a case which conforms, radiologically and otherwise, to those already described.

Clinical History

The patient, a man aged 23, was admitted to hospital because of "feverishness" and pains in the joints. Influenza was diagnosed, but routine examination revealed certain peculiar features. He was 6 ft. 5½ in. in height, and had clubbed fingers and a very blue tinge on his lips. At the age of 10 he had been told that he had "something wrong" with his heart, but this had not precluded him from having a fairly normal adolescence. He had played games at school, but could not run fast. He had had no illnesses and had lost no schooling. Three years previously he had coughed up a cupful of blood, but his doctor told him it was due to "lung strain." He had had no chest pains, cough, or expectoration. Neither had he had any sore throats, joint pains, or any kind of body rash.

The family history was not relevant, except that his father was a very tall man of 6 ft. 5 in., and his sister was also over 6 ft. The head and neck showed no abnormality. The apex beat was well out, five inches from the mid-sternal line. There was a palpable thrill, diastolic in time, over the praecordium—not like the pre-systolic thrill of mitral stenosis, but sharp and abrupt. On auscultation there was a mid-diastolic murmur, heard at the apex and conducted out. There were no moist sounds in the lungs. The pulmonary and aortic areas were clear. The liver was not palpable. The blood pressure was 140/70. The Wassermann reaction was negative, and the blood count showed 7,500,000 red cells, Hb over 140% (Sahli), and a white cell count of 13,000.

The electrocardiogram did not show a right axis deviation by measurement on the Einthoven triangle. There were inverted T waves of the coronary type, with depression of the S-T segment below the isoelectric line, in both arm and leg apex leads taken in the conventional manner. The three limb leads showed diaphasic QRS complexes suggestive of congenital heart disease as described by Katz.

X-ray examination demonstrated a huge pulmonary artery with a comma-shaped branch, also greatly enlarged in the antero-posterior position (Plate, Fig. 1). After a barium swallow a very large impression made by the right pulmonary artery in the right oblique position was a striking feature (Fig. 2).

We conclude that the above is a case of congenital dilatation of the pulmonary artery and its branches. Atrial septal defect might be present, but this is uncertain, as the right auricle is not grossly enlarged.

BIBLIOGRAPHY

- Assmann, H. (1928). *Die klinische Röntgendiagnostik der inneren Erkrankungen*, Leipzig.
Bedford, D. E., Paff, C., and Parkinson, J. (1941). *Brit. Heart J.*, 3, 36.
Jennes, S. W. (1936). *Johns Hopk. Hosp. Bull.*, 69, 133.
Master, A. M. (1939). *The Electrocardiographic and X-ray Configuration of the Heart*, Philadelphia.
Smith, Lester, et al. (1936). *Radiology*, 17, 141.

Medical Memoranda

A Case of Traumatic Ventricular Pneumocephalus

(WITH PHOTOGRAPHURE PLATE)

Cases of traumatic ventricular pneumocephalus reported in this country are few in number, and Davies (1943), who described one case of his own, could find only about twelve in the British literature. It may therefore be of value to put on record another instance of this rare condition.

HISTORY OF CASE

A man aged 33 was found lying alongside a road late at night, and was admitted to the Staffordshire General Infirmary on May 23, 1943, in a deeply comatose condition. The exact nature of the accident could not be ascertained, but from the time factors concerned it was judged that he had been lying out in the cold for at least one hour. On examination a small puncture wound, a quarter of an inch in diameter, was found over the left eye, with fairly large abrasions of the forehead and just below the outer canthus of the left eye. The eyelids on both sides were very swollen and could be opened only with difficulty, but enough to show subconjunctival haemorrhage in the left eye. There was no bleeding from the ears or nose, nor was any other injury found. His temperature was subnormal, but the pulse was fairly strong, with a rate of 78 a minute. Fracture of the skull was diagnosed; the patient was admitted and treated for shock. Because of his general condition, surgical toilet was not performed.

By the following morning he had passed into a state of cerebral irritation, and this irritation alternated with long periods of sleep. Dehydration treatment was begun, using a 50% solution of magnesium sulphate, 4 oz. six-hourly, per rectum. By the next day he was completely conscious, and complained of pain in the occipital region. This pain later shifted to the left temporal area, and during his stay in hospital he had repeated attacks of pain in this region.

On the fourth day after admission the patient complained of stiffness of his neck. Examination showed physical signs of meningitis, although temperature and pulse were normal. On lumbar puncture the cerebrospinal fluid was found to be of normal pressure and fairly heavily blood-stained. This fluid contained no coagulum, and the supernatant fluid was faintly yellow. Numerous red blood cells were present, but there was no increase of leucocytes, and films showed only Gram-positive bacilli, while, on culture, only aerial contaminants were found. The dehydration treatment was stopped and the patient put on a course of sulphapyridine, and the meningitic symptoms passed off in a few days.

Apart from the continued headaches, for which phenobarbitone was given, the patient made good progress until June 2, when he

H. L. HEIMANN AND M. M. POSEL: CONGENITAL DILATATION OF PULMONARY ARTERY

FIG. 1.—Large pulmonary artery with comma-shaped branch.

FIG. 2.—Oblique view after barium swallow, showing impression on oesophagus by right pulmonary artery.

M. GAINES: TRAUMATIC VENTRICULAR PNEUMOCEPHALUS

FIG. 1.—Lateral radiograph of skull, June 3, 1943.

FIG. 2.—Antero-posterior radiograph, June 3, 1943.