TRANSPLANTATION OF THE URETERS INTO THE LARGE BOWEL*

BY

G. GREY TURNER, LL.D., M.S., F.R.C.S.

Professor of Surgery, University of London; Director of the Surgical Unit, British Postgraduate Medical School

This subject has always interested me, for at the outset of my surgical career I saw cases of ectopia which made me realize the utter misery of the sufferers, so often to become unwanted outcasts who, like Job, survived to "curse the day wherein they were born." Treatment was most unsatisfactory, and though transplantation of the ureters into the bowel seemed the best solution I was brought up in the belief that the operation was extremely dangerous and sure to be followed by ascending renal infection and rapid death; the only justification for a return to the subject is because some members of the profession still appear to be obsessed by this fear.

For me the bright lamp of surgical progress in this field was lighted in 1911 when Stiles of Edinburgh read a paper in Denver City on two cases in which he had successfully transplanted the ureters by burying them obliquely in the wall of the bowel, in much the same way that the catheter is buried in the stomach wall in the Witzel method of gastrostomy.

After-results

In 1927 I delivered a Hunterian Lecture on this subject, and was able to give the after-history of 17 personal cases from a few months to 15 years after transplantation (Turner, 1929). The subsequent history of these same patients after the lapse of a

After-history of 9 Cases of Transplantation for Congenital Defects with Incontinence. All Previously Reported (Turner, 1929)

Case No.	Sex	Age at Operation (years)	No. in B.J.S., Vol. 17	Time since Operation (years)	Present Condition	Further Particulars
1	Male	8	1	30	In good health	Working long hours at munitions
2	Female	22	2	29	"	Three children born since operation. Does all own household duties. Twice had stone removed from right kidney and ureter
3	Male	19	3	25	"	Has worked as a farm labourer. Married
4	Female	6	5	25	"	Stone removed from kidney. Married; no children
5	Male	5	7	23	"	During one recent year several attacks of renal pain. Married
6	Male	10	12	19	"	Trouble with inguinal hernia. Otherwise well. Does heavy work
7	Male	8½	15	17	"	Nephrectomy for pyonephrosis—persistent urinary fistula. Does heavy work down a mine
8	Male	3	16	16	"	
9	Female	3½	17	16	"	Has always been splendidly well

further 16 years is now presented (see Table). The numbers of the cases recorded in the original paper are placed in the fourth column. The patients have been very closely watched, and those who have been admitted to hospital for various reasons have

been submitted to detailed examination. In no case has there been gross evidence of renal deterioration. In 1936 J. H. Saint examined several of these patients by the method of intravenous pyelography, and the results were published in a joint paper (Turner and Saint, 1936). In the original 17 cases there were four deaths directly due to the operation (peritonitis 3, septic dermatitis 1). Three patients subsequently died—from intestinal obstruction, sepsis following a plastic operation after removal of the ectopic bladder, and haemorrhage after plastic repair of the penis respectively. Thus none of the patients who recovered have subsequently died because their ureters were transplanted. Of the 9 surviving cases the essential details are set out in the accompanying table.

In Case 7 an enormously dilated ureter could not be buried and was simply tucked into the bowel end to the side. In Case 9 a submucous bed was prepared. In the other cases the original method of Stiles was used, one ureter being implanted at a time and with a varying interval between interventions. Full details of the operations are given in the original paper. "In good health" means that the patients can stand up to their environment and take their place among their fellows. No. 1 has never been a big strong person, but he has developed quite well, played games, including football, and has always been ready for work when it was available. No. 2 is a brave woman now 51 years of age. Her first child was born three years after the operation, and both pregnancy and delivery were without unusual incident. Ever since the operation she has had some pain in the back, but it has only been very troublesome at infrequent intervals during the last five years. In 1938 a phosphatic calculus was removed from the right kidney, which was considerably dilated, and early this year a large calculus was removed from the corresponding ureter. The patient is now very well. Except for these attacks her health has been good, and her greatest happiness has been to look after her family and household. She has carried out all the domestic duties which fall to the lot of a North Country woman of the artisan class. After her day's work she has often visited the pictures and has gone from home on holiday each year. For years No. 3 did the hard work of a general farm labourer. His marriage about three years ago was for companionship, but is not entirely Platonic. Case 4 has also had a stone removed from the right kidney 14 years after the transplantation; but she has always been well and active, carrying on the work of a ladies' hairdresser, which in her case entailed a considerable amount of travelling. Three years ago she married, and is anxiously awaiting maternity. No. 5 was a feeble child but has developed into a sturdy young fellow. Four years ago he was exceptionally well, was working long hours, and dancing in the evenings by way of relaxation. During the past twelve months he has had several attacks of renal pain; but these symptoms left him after passage of a considerable amount of grit, and he is now working regularly in charge of a factory. No. 6 has enjoyed good health except for troubles due to double inguinal herniae, one of which had to be operated upon in a state of strangulation; but despite this handicap he puts in a long day as a crane-man. No. 7 has had a tedious time with many vicissitudes, including nephrectomy for pyonephrosis. None the less he is well developed, strong and active, and capable of hard work. No. 8 has been difficult to keep in touch with, but he is reported to be well and strong. No. 9 was operated upon at an early age, but she made an easy recovery, has developed normally, has never shown any untoward symptoms, and has grown into an attractive girl who enjoys excellent health. Many other cases operated upon since the above series have been equally satisfactory, and I would just recall a charming girlie born with an ectopic bladder whose parents were thrown into the depths of despair as the result of the utterly pessimistic outlook of the

* Presidential address delivered at the Medical Society of London, Oct. 18, 1943, much abridged.

consulting surgeon who was called to see the case a few days after birth. The transplantations were carried out between 3 and 4, and now, eight years later, the child is perfectly well, highly intelligent, and especially fond of swimming. Also another girl, who wrote four years after operation: "I have to catch a bus in the mornings at a quarter to eight in the black-out, but it is fun!" An officer whose ureters were transplanted just previous to total cystectomy for neoplasm is now, two years later, back to his military duties, albeit on the home front, while in like case a street trader is actively employed in his traffic. I used to feel that every case of transplantation would sooner or later show some evidence of ascending infection, but I am now hopeful that with the great simplification of the technique there may be many in which it never occurs. The exact condition of these patients may be regarded as the results of an experiment in applied physiology, and can be detailed as follows.

Details of Results

Effect on the Economy of the Body.—Just after the transplantation the patients are invariably abnormally thirsty, and this may continue for about a year. During this period they often lose weight, and sometimes very considerably. After this time thirst diminishes and they not only regain lost weight but begin to put on flesh, so that they come to look plump and well nourished. Appetite is usually good and mental acuity more than average. They behave like normal individuals, are cheerful and happy, and able to work and play like others of their station. They can also stand up to the buffets of their environment, including physiological processes like pregnancy, operations, and illness.

Rectal Function and Control.—In nearly all my cases this has been satisfactory, though in two of them continence during the night has never been acquired, though these same patients have complete control by day. While there is some variation, most patients can hold the contents of the rectum for three to four hours at a stretch, and the call is not so urgent but what they can await a convenient opportunity with confidence. Some patients sleep throughout the night without being disturbed, while others may rise once or twice. All agree that the behaviour of the rectum is much influenced by the general health. With exhaustion or indisposition rectal calls are more frequent. The contents voided are usually an intimate mixture of faeces and urine, but sometimes the same patient may pass clear or opalescent urine at one time and a solid evacuation at another. The rectal mucous membrane is unaffected and, except for unusual moisture, looks normal when inspected through the sigmoidoscope.

Where the Urine is Stored.—At the time of my first communication many considerations led me to conclude that the urine was stored in the large bowel and not in the rectum. This conclusion is amply supported by urography.

Effect on the Kidneys and Ureters.—The changes in structure are sometimes very striking. In the first few weeks the ureters and pelves may be dilated, and this dilatation may persist and increase. In the series of examinations made in collaboration with J. H. Saint four out of six patients showed a moderate degree of unilateral hydronephrosis, and in another the condition was bilateral. The most striking feature was the general good health of the patients examined. Marked unilateral hydronephrosis was also found in a patient who died subsequent to operation for intestinal obstruction, but that boy had enjoyed good health until his fatal illness. In nearly all of the earlier cases there was clinical evidence of some ascending renal infection, but for the most part it was transient and did not materially interfere with recovery. From the clinical aspect only those cases in which there has been obstruction in the lower ureter have suffered seriously. This is in keeping with general pathology, and conforms to the rule that infection with good drainage is often neither very troublesome nor dangerous. The ureter may be dilated or may look normal, but the actual junction with the bowel and the opening into the lumen is not always demonstrable either by radiographs or by sigmoidoscopic examination. When visible it may take the form of a nipple with a terminal or lateral orifice, a rosette, or a tiny opening hidden behind one of the rectal folds.

Should Both Ureters be Transplanted at the Same Sitting?

Up till quite recently I have always taken the view that it is safer to transplant one ureter at a time, and the wisdom of this course has been borne out by statistics. With better understanding of the problems involved, a simpler technique, and more knowledge of after-care, I am tending more and more to carry out

a simultaneous transplantation, except in children, and even in them the hazards of a second anaesthetic may perhaps balance the only slight extra risk of the double transplantation.

What is the Best Technique?

The technique has now been very much simplified, and is carried out without any special apparatus. Clamps are not used either for the ureter or for the colon. Coffey's tubes are no longer employed, and Charles Mayo's catgut guide has been discarded. The main points are to make an oblique implantation into a submucous bed in the bowel without kinking or compression and with due regard for the blood supply of the ureter. Compression from too tight suturing or the result of haematoma or traumatic oedema is probably the most harmful thing that can occur. Fine 3/0 or 6/0 chromic catgut is used throughout. Drainage is not employed.

Preparation and Operation

Preparation will differ a little, depending on whether the surgeon is dealing with a congenital defect in an otherwise healthy child or the operation is for the relief of some condition, such as malignant disease of the bladder, likely to be associated with sepsis or general deterioration. No operation should be done until the renal function has been raised to as nearly the normal standard as possible. The bowel should be well cleared by a reliable purgative or by an enema not later than 12 hours before operation. If there is distension or constipation a full dose of castor oil is the best preparation. No attempt should be made to sterilize the bowel or even to cleanse it by repeated irrigation, which may result in the surgeon's encountering a wet puddle in the pelvic colon. The choice of anaesthetic will depend partly on the condition of the patient and partly on the practice of the surgeon, but whatever method is employed complete relaxation is necessary. The patient is placed in the Trendelenburg position and the abdomen opened by one of the vertical incisions, which must be long enough to give ready access to the pelvis. I have made it a practice to transplant the right ureter first, but I am not able to give any valid reason for the choice. If both are to be transplanted at the same sitting I am sure that it should be into the same side of the bowel, and the right or inner side is the more convenient. To bring this about it is usually best to pick up the left ureter on the outer side of the sigmoid and to bring it through the mesocolon to the inner side, where it is made to enter the bowel about 1½ in. higher than its fellow. It is usually easy to identify the ureter as it crosses the pelvic brim. Spontaneous vermication, or this movement elicited by mechanical stimulation, is characteristic. The posterior peritoneum is incised in the length of the ureter for about 3 in., and the ureter is gently lifted from its bed and traced towards the bladder. Not more than 2½ to 3 in. should be isolated: in this respect illustrations in surgical journals are misleading. At the lower end the ureter is clamped and divided obliquely; the end is securely tied, and its open end carbolized and allowed to retract. The upper end is held by a suture passed from its lumen and out through its upper surface a quarter of an inch from the cut end. This is tied, and is subsequently used for drawing the ureter into the lumen of the bowel and for fixing it to the bowel wall. The ureter is now turned backwards over a pad of gauze; this kinking is enough to prevent the escape of urine. A few interrupted sutures draw together the edges of the incision in the posterior parietal peritoneum.

The position of the incision to be made in the bowel is now determined by finding out where the ureter lies most comfortably against it without kinking. The aim is to carry the ureter obliquely into the bowel in the same alignment as it occupies in its retro-peritoneal bed. The site for the right ureter will probably be in the lowest part of the sigmoid or the uppermost end of the first part of the rectum. The actual incision in the bowel will usually be 1½ or 2 in. long, and this is marked out by guide sutures. The incision is made obliquely through the muscular wall down to the submucous layer. The edges will retract, but this process may require to be helped by gently spreading with dissecting forceps or a few touches of the knife, used with the blade on the flat. At the lower end of this oblique bed a small hole is made into the bowel lumen. The idea is to make such an aperture as will comfortably admit the ureter, but in practice it usually works out a good deal larger. Both ends of the stitch on the ureter are now threaded on a small round needle, which is passed through the hole into the bowel and out through the whole

thickness of its wall half an inch beyond the aperture. The ureter may have to be manipulated through the latter. Both are pulled upon to draw the ureter to this site and well up against the inner surface of the bowel. The ends are then tied firmly, but not so tightly as to cut through the tissues. The edges of the incision in the muscular layer are now sewn over the ureter with four or five interrupted stitches. If there has been any doubt about the fixation of the ureter the first muscular stitch takes a bite of its wall. The peritoneum is then drawn together over the first sutures, taking great care to bury the ureter-fixation stitch properly. One or two more than the number of deep stitches will be required. It is particularly important that the stitch nearest to where the ureter enters the bowel should be lax, as it is at this point that obstruction may be caused by subsequent oedema or kinking. If after tying this last stitch the ureter is seen to fill up, the suture is causing obstruction and must be removed. If excretion is going on, the urine should be escaping into the bowel while the anastomosis is being made, and the ureter should never distend. The ends of this last stitch are left long, and are used to anchor the bowel to the cut edge of the posterior parietal peritoneum. In this way at the conclusion of the operation the ureter passes directly from its retroperitoneal bed into the bowel, so that it has no intraperitoneal course. If the second ureter is to be transplanted at once the same steps are taken, but the incision in the bowel wall should be an inch or an inch and a half higher up than the first implantation. After completing the toilet of the peritoneum the omentum should be crowded into the pelvis and the patient brought to the horizontal position for closure of the abdomen. For some three years now I have omitted drainage from the site, but it is a perfectly rational safeguard, and the surgeon need not hesitate to use it. The abdominal wall must be very carefully closed, some through-and-through sutures being used. As a last step a tube about forefinger size is passed into the ampulla of the rectum and fixed to the anal margin by suture. This is to prevent the accumulation of a puddle in the rectum which might encourage infection.

The whole procedure takes from three-quarters of an hour to an hour and a quarter, depending upon whether or not both ureters are transplanted.

Immediate After-progress and Treatment

There should be no shock and very little upset of any sort. Urine may be discharged into the rectum from the outset, but it is more usual for uresis to be delayed for about six hours. At all events, only if there is no evidence of secretion by the end of that time should active steps be taken to encourage it. Some patients vomit a good deal and a moderate amount of distension is not uncommon. Such symptoms nearly always clear up as soon as uresis is freely established. I have always avoided the use of enemata, fearing to disturb the site of anastomosis. The patient reaches the ward with a glucose-saline drip *in situ*, and this is continued until there is a free discharge of urine from the rectal tube. When the patient recovers from the anaesthetic, the drinking of hot water is allowed and, in the absence of vomiting or distension, encouraged. It is only in cases with rather poor renal output or in which uresis is delayed that the intravenous sodium sulphate 4.3% is employed.* The rectal tube should be removed in four days, or earlier if it is much resented. Thereafter there may be incontinence for some days, especially in children, and adults may have very frequent calls—perhaps every hour. But even children acquire rectal control quite quickly, and nearly always by the time they are ready to leave hospital—in three or four weeks. Of course some education is usually necessary, and to this end it may be a help to keep the bed-pan in the bed. A capable and understanding nurse will usually manage to assist even small children to acquire control within this time.

Complications

In the early stage anuria, chest troubles, and peritonitis are the only likely troubles. At the end of about a week distension, local tenderness in either iliac fossa, quickening pulse, elevated temperature, and general malaise are very suggestive of intra- or extra-peritoneal infections about the site of anastomosis. Such a condition is grave but not necessarily fatal. In two cases (Nos. 4 and 5) a localized abscess formed and after evacuation

was followed by urinary fistula, yet spontaneous recovery ultimately ensued, and both patients are now, many years afterwards, alive and well. Some infection of the kidney used to be so frequent as to be looked upon as part of the normal convalescence, but in more recent years it has seldom occurred, possibly owing to the simplification of the technique. Most cases turn out to be mild, but, even so, the onset may be rather alarming and attended with considerable general disturbance. As a rule this soon settles down, leaving a swollen tender kidney with some fever as the only indication. In the more serious types things go from bad to worse until the patient presents the picture of acute ascending pyelonephritis, but I have not seen death from this cause for some years. In his anxiety and apprehension about possible renal infections the surgeon must not forget that these patients are liable to the occasional complications attending on laparotomy for any purpose. I have successfully dealt with a case (No. 3) in which a loop of small intestine escaped through the wound and became strangulated, and the burst wound is not unknown.

Mortality

The mortality rate depends to a considerable extent on the condition demanding the operation. In the non-malignant cases and especially the congenital deformities it should be very low—perhaps 5%—but in the malignant cases such a low rate is not to be expected if the possible advantages of the operation are to be offered to the greatest number of sufferers. In children some deaths have been due to chest complications and to the exanthemata, and I would stress the wisdom of choosing the summer months for the ordeal and of keeping children in hospital for a fortnight before operation.

When to Transplant the Second Ureter

There is no stated time for dealing with the second ureter, and the interval must depend on the progress made after the first intervention. If all goes perfectly well, three weeks has proved a proper interval; should it be otherwise, the surgeon must bide his time. The only error is to re-operate too soon.

Some Late Sequelae

If pyelitis is going to develop it usually comes on within a fortnight of the transplantation, but once having developed it tends to recur at varying intervals, perhaps over a lifetime. Hydro-nephrosis, on the other hand, probably develops over the course of months, and may not give rise to symptoms unless it becomes infected. Quite often it is discovered only because looked for. Calculus formation is not very rare, for it occurred three times among my first 13 patients who lived long enough to develop the condition. The symptoms became suggestive 15 years, 22 years, and 25 years after the original operation. Frequency of rectal micturition has seldom been a constant feature, but it may be an accompaniment of general ill-health. Rectal incontinence for urine is in my experience the most disturbing sequel. Fortunately it is unusual. In spite of painstaking investigation I have been unable to discover the cause, nor have I been successful in treatment. Wearing a rectal tube at night—a large catheter of the de Pezzer type—is a useful palliative measure.

Special Groups of Cases

Congenital Anomalies.—Of these the commonest is ectopia vesicae, which is said to occur once in every 50,000 births. The next are the lesser degrees of the same condition, such as complete epispadias in the male and the corresponding anomaly in the female known as subsymphysial epispadias. *The age at which to operate* is an important problem. I consider that the optimum time will occur somewhere between 4 and 6. But it is not so much a question of the number of years but of the condition, and whatever the age the operation must not be carried out until the child is in established good health. Many a child at 4 is better fitted for the operation than others at 6. But it is never too late to consider operation if congenital deformity is the indication, and my friend Mr. Norman Hodgson dealt most successfully with a case of ectopia in a man of 51 who, by a strange irony, had consulted me 30 years previously, only to be told that no operation was feasible. As a rule children do better than adults, and it is surprising how readily they accommodate themselves to the altered physiological condition.

* 42.85 g. of hydrated sodium sulphate (Glauber's salt) dissolved in one litre of distilled water gives an isotonic solution of 4.3%.

The exposed mucous membrane of the ectopic bladder is a distinct danger, and many examples of the development of epithelioma have been recorded. Quite apart from this risk it is offensive to the eye, may give rise to an unpleasant discharge, and is liable to mild injury and excoriation. In either sex some sort of plastic repair of the affected area is desirable.

With the lesser anomalies the question of marriage is often raised, and even in complete ectopia patients of either sex have sought the consolation of matrimony. The female with a normal partner may prove fruitful, and there is no reason why such marriages should not be happy. There appears to be no special risk in pregnancy, and, since the pubes is constantly separated, labour is often easy (Case 2). In the male sexual gratification is said to occur, but neither of my married male patients has as yet become a father. Now that the transplantation is less likely to be attended by sequelae, I welcome the opportunity of carrying out more careful and complete restoration of the genitalia.

Separation of the Pubic Bones.—In the case of complete ectopia the pubic symphysis is separated to the extent of from four to six inches, but the bones are connected by a very strong ligament. It is often assumed that such a degree of separation will result in marked disability, but though the thighs are more or less widely separated and the patient walks with a characteristic gait, they are usually quite strong and able to carry out ordinary activities without impediment.

Other pre-operative Problems that may Arise.—Sometimes there is weakness of the anus with incontinence, and on occasions prolapse. If this is associated with some degree of spina bifida and perhaps weakness of the lower limbs no improvement can be expected, and transplantation into the usual site is contra-indicated. Confronted with these circumstances, Hey Groves made the transplantation into the caecum, and combined this with a left inguinal colotomy. As a result not too frequent pul-taceous stools could be easily controlled by a colotomy belt, and for some years the patient was able to lead a sheltered but useful life.

Contraindications to Transplantation.—Marked pyelitis and pyonephrosis are usually regarded as contraindications to transplantation. It is always worth while establishing external loin drainage by nephrostomy and, if sufficient improvement follows, to proceed to transplantation while the drainage is still operating.

The Widening Scope for Transplantation

For many cases of *malignant disease of the bladder* total cystectomy holds out the best prospect of long palliation or even cure. In this operation it is necessary to divert the urine, and by far the best plan is to carry the ureters into the bowel. The technique is the same as that used in other cases. When the transplantation is a proved success—and this is usually assured in three or four weeks—the removal of the bladder may be carried out.

In some cases of *malignant disease of the cervix* extension to the adjoining bladder renders treatment both by radium and by hysterectomy impracticable. In such cases total hysterectomy, together with simultaneous removal of the bladder, holds out some prospect, and in the absence of glandular involvement or distant dissemination should be practised. The transplantation of the ureters should be a first stage, and the next part of the operation should not be undertaken until the patient has become stabilized under the new physiological conditions. An interval of 6 to 8 weeks will probably suffice, but there should be no unwarranted haste.

For some years now the operation has been used for certain intractable cases of *vesico-vaginal fistula*. In such cases transplantation should not be combined with a last desperate attempt at local repair, but should be a deliberate intervention only to be carried out after proper preparation. In the *systolic bladder*, which is sometimes a problem after the successful removal of a tuberculous kidney, diversion of the ureter into the bowel may bring great relief.

Transplantation has also been used in certain cases of severe *injury to the urethra* usually associated with fractured pelvis. Though this indication is among the rarest, it should be kept in mind.

REFERENCES

- Turner, G. Grey (1929). *Brit. J. Surg.*, 17, 114
— and Saint, J. H. (1936). *Ibid.*, 23, 580.

OCULAR MANIFESTATIONS OF NEUROSES COMMONLY FOUND AMONG SOLDIERS

BY

I. C. MICHAELSON, F.R.F.P.S.

Major, R.A.M.C.; Ophthalmic Specialist, British General Hospital;
Extra Surgeon, Glasgow Eye Infirmary

Fear and environmental stress are very prevalent in wartime and often give rise to chronic anxiety. Many soldiers react to such a chronic anxiety by developing a neurosis, of which there are two main varieties—hysteria and the chronic anxiety state. Whether one or the other will develop depends on the individual personality type.

The hysteric has an essentially childish personality with an undeveloped or perhaps only unsustained sense of social responsibility. He reacts to the anxiety by dissociation, which may affect nerve, muscle, bone, joint, an entire limb, memory, the conscious state, etc. He is very open to suggestion, and the type of somatic localization may be suggested by his observation of the infirmities of others or his memory of past disabilities within himself. In this way undue prominence is given to a condition already present, or else one is created entirely *ab initio*. The somatic incapacity is such as to still the anxiety or the conflict that may have arisen between it and duty. The more successfully this is done the happier is the patient, who may indeed wear a fixed look of quiet satisfaction throughout the day.

The patient who develops a chronic anxiety state is usually a more mature personality with some awareness of his obligation to society. His reaction to the anxiety provoked by fear or environmental stress is different from that of the hysteric. The early somatic upset appears to take place in the endocrinal and autonomic systems, leading to disturbances of the vasomotor, respiratory, sudomotor, or gastro-intestinal system. In this way are produced palpitations, vascular throbbing, headache, syncope, flushes, paraesthesia, breathlessness, sighing, undue sweatings, vomiting, diarrhoea, and dry mouth. The somatic incapacity is frequently not sufficient to assuage the anxiety of the patient, who may present a worried, unconfident attitude to the outside world.

As a result of hysteria and chronic anxiety state, various conditions arise in and around the eye. Indeed, it may be said without exaggeration that half of the soldiers presenting themselves at the out-patients' room of the eye department of this hospital in the Middle East have complaints which may fairly be described as functional. The most frequent, and those that will be dealt with in this report, are defective day vision, defective night vision, asthenopia, diplopia, headaches, photophobia, spots before eyes, and epiphora. All of these symptoms are found in organic conditions. Investigations into each of them should therefore take three main directions: (1) exclusion of organic disease, or, if present, an assessment of its role in the production of the symptoms (organ-symptom parallelism); (2) a detailed investigation into the nature of the symptoms; (3) some investigation into the patient's personality (while in difficult cases this is the sphere of the whole-time psychiatrist, the ophthalmologist should be capable of the simpler approaches). As investigation No. 3 applies to all the symptoms similarly, a brief generalization will be made now of the main points of inquiry.

Inquiry should be made into the presence of any worries, dislikes, or fears, and the degree of duty sense as evidenced by his former reactions to life. Search should be made for anything in the personal or family experiences of the patient, or in his general environment, that might have suggested the symptom to him. This may be a weakness of one eye which the patient had many years ago, or a history of blindness in his family, or the defective night vision of which his friend in the unit has complained. The relation of the symptom to change of environment should be noted, especially when a new climate provides factors, such as heat and glare, which can be blamed for it. It is noticeable how often among the troops in the Middle East a visual complaint began on the voyage out between Freetown and Capetown. It is as if there were an incubation period for the symptom.