"try and stick it out." He said he suffered from aches and pains in his back and legs, especially in the muscles of the calf and in both knee-joints. He also had a severe frontal headache. He said that he had had rheumatic fever eleven years previously, and since then had suffered almost every year from similar attacks. In civilian life he went to bed and was attended by his doctor; the last attack was two years previously. On examination his temperature was 104.8° and his pulse 120. The only physical sign detected was a soft presystolic mitral murmur, which was not heard four days after the onset of the pyrexia. He was confined to his hammock and given mist. sod. sal. gr. 20 and luminal gr. 2. The following day he was given mist. sod. sal. gr. 20 and aspirin gr. 10 t.d.s. On Aug. 14 all medicines were discontinued, and he returned to duty quite well.

FIG. 3.—Temperature Chart of Case 3.

Owing to the exigencies of the Service at the time it was only possible to obtain blood films from each of the patients on Aug. 12, 1943, and examine them microscopically in a large ship, in order to exclude malaria or leishmaniasis; blood counts were not done. No parasites were found in a careful examination of the slides. Paratyphoid, dengue fever, and influenza were all considered, and were rejected as improbable. Cases 1 and 2 were diagnosed as phlebotomus fever, and Case 3 as pyrexia of rheumatic origin. These patients suffered from intense supra-orbital headaches and generalized "rheumatic" pains, which varied in intensity. In all three cases there was absence of severe gastro-intestinal symptoms such as may be expected in the later stages of sandfly fever. The pyrexial period varied between 3 and 8 days. Bradycardia was not noticed in either of the cases of sandfly fever.

Conclusion

Case 3 resembled no particular syndrome with which I am familiar, but appeared to be related to the rheumatic series of diseases.

The cases resembled one another closely, and, by coincidence, occurred at the same time.

Cases 1 and 2 were diagnosed as phlebotomus fever because of the short duration of pyrexia, the presence of insect-bites, and the symptomatology. The fever, as is known, is caused by a virus introduced by the bite of *Phlebotomus papatasi*, which is present in large numbers in Sicily from May to October.

Almost constant "action station" routine caused the patients discomfort owing to the stuffy atmosphere in the ship, but it was possible to treat such cases of high pyrexia on board a destroyer provided that the patient was nursed efficiently.

According to Wenscher (*Dtsch. med. Wschr.*, 1942, 68, 984) neurological signs aid in the early diagnosis of typhus. The first indication is sudden high rise in temperature, accompanied by a typical and unusually stubborn headache which responds only slightly to analgesics. An additional sign is hardness of hearing, which occurs in almost all cases. If these signs are accompanied by speech disorders and the characteristic twitching of the perioral muscles the diagnosis of typhus exanthematicus is assured even without a distinct exanthema. Therapy consists in relieving the pressure. Apathy or restlessness responds quickly to lumbar puncture; circulation and respiration improve and the distressing headache disappears. Along with the essential support of the cardiovascular system, the main therapeutic concern is the prompt reduction of increasing pressure in the c.s.f. For this reason strophanthin should always be given in strongly concentrated glucose solutions, which exert a dehydrating action. After damage to the vasomotor and respiratory centres, the next most important danger is the occurrence of bulbar paralysis.

Medical Memoranda

An Unusual Form of Ectopia Testis

Sir John Thomson-Walker (1936) quotes Eccles (1903) as considering it extremely doubtful if the testis is ever spontaneously passed into the upper part of the thigh through the femoral ring. The following is the record of such a case.

Cpl. A., aged 43, was admitted to hospital with the diagnosis of right inguinal hernia. Seven years previously, while engaged in heavy work, he felt a sharp pain in the right groin followed by the appearance of a lump; the swelling had remained there since. Examination showed a swelling the size of a hen's egg in the right groin, over the junction of the middle and inner thirds of the inguinal ligament. The swelling was not reducible, felt cystic, and was attached to the deep tissues. The right testicle was present in the scrotum, the left half of which was empty.

Operation showed the swelling to be an ectopic testis which had descended through the femoral ring and canal, and had all the coverings of a femoral hernia. A hydrocele was present; this was aspirated, and the testis then made to retrace its path, the femoral ring being exposed from above as in Lotheissen's operation. Dissection revealed a hernia of the congenital type, the neck of which was very narrow. The fluid in the sac had been restricted to the distal extremity by the constriction to the sac as it passed through the femoral ring, forming a hydrocele of the hernial sac. The spermatic cord consisted only of a fine bunch of vessels, the testicle itself being extremely atrophied.

Two interesting features are that the testis had passed through the femoral ring to gain the upper part of the thigh, and that both normal and ectopic testes were on the same side, the surmise being justifiable that one testis found its way through the femoral ring because the normal passage through the inguinal canal was already occupied.

My thanks are due to Prof. R. C. Alexander, Surgical Director for the area, for permission to publish this case.

T. C. SKINNER, F.R.C.S.Ed.,
Surgeon, E.M.S.

REFERENCES

- Eccles, W. McAdam (1903). *The Imperfectly Descended Testis*, London.
Thomson-Walker, Sir J. (1936). *Surgical Diseases and Injuries of the Genito-Urinary Organs*, p. 844, London.

Recovery after 700-foot Fall

The following case would seem unusual enough to merit being placed on record.

CASE REPORT

A well-built paratroop pupil aged 25, weighing 11 st., was parachute-jumping from an aircraft travelling at 80 m.p.h., at a height of 700 feet, into a 10 m.p.h. wind. Unfortunately, although the parachute came out of its pack, the canopy did not develop properly, with the result that the pupil landed some 10 seconds later on a ploughed field, dropping at a speed which must have been in the region of 50 m.p.h. When examined a few seconds later he was found to be still alive and just conscious. He was lying on his left side in a crouched position, with his knees drawn up towards his chin. The pulse rate was 70 a minute and weak; breathing was shallow and difficult owing to much mucus being brought up, and there was a small amount of thin yellowish fluid exuding from his right ear. Physical examination showed no obvious signs of fracture or paralysis. The pupils were equal and of normal size.

A "tubunic" ampoule (morphine hydrochloride gr. 1/3, atropine sulphate gr. 1/120) was given, following which he was put on to a stretcher. A few minutes later he became very restless, and as he was now trying to move his arms and legs he had to be strapped securely to the stretcher during transit to hospital. Next day he looked very ill, but was cheerful and mentally clear, showing no signs of disorientation or amnesia. He could move his arms and legs without difficulty and his excretory functions were normal. Two days later the improvement was so marked that he was considered fit for x-raying, when it was discovered that he had a compression fracture of the 12th thoracic vertebra. After immobilization in a plaster cast progress was satisfactory and uninterrupted.

COMMENTARY

Partial failure of a parachute is extremely rare and is almost as unique as this man's survival of the accident, which was undoubtedly an extraordinary escape, especially in view of the fact that all previous occurrences of this nature had resulted in instantaneous death either from multiple injuries or from fractured cervical vertebrae.

Reliable eye-witnesses of this descent were of the opinion that the degree of inflation of this soldier's parachute was so slight and his fall so rapid that he could not possibly escape death. It is thought that he landed on one foot and fell sideways, thus preventing the full force of his landing being transmitted through the legs and spine to the base of the skull,

which would have resulted in a fracture in one or more of these places as in the other cases.

We desire to thank Squad. Ldr. J. D. Milne, R.A.F., and Major R. B. Davies, I.M.S., for permission to publish this case report, and would like to emphasize the co-operation between all the medical services concerned.

F. G. NEILD,
Capt., R.A.M.C.
C. M. MACKENZIE,
Flight Lieut., R.A.F.V.R.

Unusual Burn of Cornea from Molten Wax

Dr. H. Sewell Sims, of Palmer's Green, asked me to see a female patient whom he had attended the previous day. She complained of loss of sight after flicking out a lighted wax taper with which she had lit a gas stove, when she said some wax flew between the eyelids. There was no reaction to this foreign body, and the appearance of the cornea was quite normal till, on examination through a loupe, a perfectly transparent layer of wax was seen covering two-thirds of the cornea, which looked as if it had a contact lens lying on it. Until this was removed the cornea did not stain with fluorescein; afterwards approximately two-thirds stained, but the erosion was quite superficial. The wax was removed with a blunt instrument under cocaine, when the vision at once improved from one-tenth to one-third of full vision. Atropine and castor oil were instilled and the eye bandaged. The effects of the cocaine passing off, the eye wept profusely and was very painful. With the ophthalmoscope there was seen a fine line on the cornea, which marked the limit of where the wax had

rested. In spite of the extent of the lesion it had no depth, only the epithelium being affected. It is quite feasible that it acted as molten metals often do—forming a cast of the front of the eye, with a fine layer of protective vapour between. Usually, of course, the lids close so quickly that the cornea escapes injury.

Considering the risks that housewives run several times daily, it really is remarkable that injuries of a similar nature from splashes of boiling fat are not more common. Würdemann (1932) mentions boiling soap, fat, or lard as being possible causes of burns. Sir Stewart Duke-Elder (1938) makes no mention of this in his *Textbook of Ophthalmology*, though there is a diagram of a case of Mr. Doggart's which rather resembles this class of burn—a case of keratoconjunctivitis due to mustard gas. Haab (1909) has a plate in his *Hand-Atlas* showing a lime-burn of the cornea, which is more or less the same in appearance. Fuchs (1933) does not mention it.

Figs. 1a and 1b show the appearance after removal of the wax, Fig. 2a the line of demarcation, which, however, had entirely disappeared three days afterwards, when the eye was quiet and the vision normal again. Figs. 2b and 2c show the area stained by fluorescein. On the third day there was no staining, and the lesion had healed.

SYDNEY TIBBLES, L.R.C.P.&S.Ed.,
Late Refractionist, Royal Hospital, Richmond.

REFERENCES

- Duke-Elder, Sir Stewart (1938). *Text-Book of Ophthalmology*. 2, 1713, London.
Fuchs, H. E. (1933). *Diseases of the Eye*, 10th Eng. ed., Philadelphia.
Haab, O. (1909). *Atlas of External Diseases of the Eye*, Philadelphia.
Würdemann, H. V. (1932). *Injuries of the Eyes*, 2nd ed., p. 11, St. Louis.

Reviews

FRACTURES

Pictorial Handbook of Fracture Treatment. By Edward L. Compere, M.D., F.A.C.S., and Sam W. Banks, M.D. (Pp. 351; illustrated. 27s.) Chicago: The Year Book Publishers; London: H. K. Lewis and Co. 1943.

Fractures. By Paul B. Magnuson, M.D., F.A.C.S. Fourth edition, revised. (Pp. 511; illustrated. 30s.) London: J. B. Lippincott Company. 1943.

It is noticeable that the large number of textbooks on fractures which have appeared within recent years coincides with improvement in the results of treatment which has come about following segregation under specially trained staffs. Not all fractures, however, can be dealt with in special units, and it is necessary for both general surgeons and practitioners to be familiar with the most effective methods of reduction, fixation, and after-treatment. The scope for textbooks of fracture treatment is therefore wider than ever. The *Pictorial Handbook of Fracture Treatment* needs no apology such as we find in the preface. It is an excellent little manual and the pictorial part is particularly good—e.g., Fig. 130, p. 265, illustrates the method of production as well as the stages of treatment of "bumper" fractures; while Fig. 151, p. 295, depicts the way in which Colles's fracture and fractures of the os calcis and lumbo-dorsal part of the spine are produced. Fig. 163, p. 313, claims to depict Böhler's method of reducing a flexion compression fracture in the dorsal spine, but the credit for this method of treatment should, we believe, be given to Watson-Jones of Liverpool, and it is not usually desirable to anaesthetize the patient. Fractures of the patella are well done—it is pleasing to notice that the author does not support a recent radical and quite unjustifiable tendency to excise the simple transverse variety. Fractures of the skull may be thought to be rather briefly dealt with, but these fractures usually constitute a minor part of head injuries calling for treatment, and the surgical treatment—as opposed to first-aid treatment—of skull fractures may reasonably be considered to be beyond the ambit of this book. The book can confidently be recommended as a modern and well-illustrated outline of fracture treatment.

Like most new and the recent editions of older textbooks of surgery which have appeared in the last few years the fourth and latest edition of Magnuson's *Fractures* shows the influence of the war, and efforts have been made by the author to render the book more useful to surgeons in the Services. It is not surprising, therefore, that the principal additions deal with transportation and the early treatment of compound fractures. Written clearly in rather dogmatic style and well illustrated by line drawings and x-ray photographs this textbook is a straightforward guide which should be particularly helpful to medical students, practitioners, young surgeons generally, and Service medical officers. The line drawings, of which there is an abundance, appealed to us especially, as they indicate very clearly the steps in treatment of particular fractures—e.g., the method of reducing supracondylar fracture of the humerus (Fig. 5, p. 101)—a series of drawings showing the manipulations necessary and the reason for them. We are glad to find that the term "Pott's fracture" is still used despite a recent tendency noticed in some quarters to discard it. We disagree with the statement (p. 429) that "every patient with a skull injury accompanied by coma should be kept flat in bed at all times for a minimum period of two weeks." Nursing these patients with the head raised lowers intracranial venous congestion and has proved more effective; but in general the sections on skull fractures and their complications and those on fractures of the spine are as well done as the remainder and bulk of the book. Of the recently published books on fractures—of which there has been almost a spate—this is certainly one of the best.

FOOD POISONING

Food Poisoning: Its Nature, History and Causation: Measures for its Prevention and Control. By Elliot B. Dewberry. Foreword by Gerald R. Leighton, M.D. In three parts; with appendices. (Pp. 186; illustrated. 15s.) London: Leonard Hill, Ltd. 1943.

Food poisoning is a complicated and extensive subject, but within 180 pages the author has covered its main features. The book is severely handicapped by being a compilation from other sources and not by an author who has himself investigated