RESPIRATORY TUBERCULOSIS

EFFECT OF THE WAR ON THE LENGTH OF THE
INTERVAL BETWEEN NOTIFICATION AND DEATH

BY

E. LEWIS-FANING, B.Sc.Econ., Ph.D.

Of the Statistical Staff of the Medical Research Council

Among many factors that have been suggested as influencing the increase in tuberculosis mortality which accompanied the outbreak of the war is that of lowered resistance to the disease by those already infected. It is argued that if resistance were lowered to the point at which the average duration of the disease was shortened, this would tend to produce the higher mortality rates that were a feature of the 1940 and 1941 statistics.

Method of Investigation

The medical service of the Middlesex County Council, realizing the desirability of answering this question, offered for statistical analysis their records of fatal cases terminating in the years 1937 to 1941. The method of approach to the problem has been to compare the interval between notification

Analysis of the deaths in the pre-war period shows that the missing data were not distributed by age in the same proportions as the deaths officially recorded. According to official figures, 43% of the male deaths were at ages over 45, yet 72% of the male leakage occurred at these ages. Again, 20% of the recorded female deaths were at ages above 45, yet the female leakage was 41% in the same group.

Returns were submitted by each of the seven chest clinic or dispensary areas into which, during the period under review, the county was divided for the purpose of administering the tuberculosis service. The information supplied in each fatal case was: Sex, age at death, date first seen by the tuberculosis officer, whether diagnosis was of pulmonary tuberculosis or tuberculosis of other organs, and date of death. The analysis was restricted to deaths due to pulmonary tuberculosis, since there was not sufficient material from which to make satisfactory deductions in regard to the other forms. Two dispensaries supplied the date of notification instead of the date first seen, but exhaustive comparisons between the results given by these two areas as compared with the remainder showed no differences which could be ascribed to the distinction made.

The data were carded and the durations calculated to the nearest month. The frequency distributions in broad duration

TABLE I.—*Pulmonary Tuberculosis: Middlesex County Council's Data. Distribution of Fatal Cases according to Age and Duration.* Fatal Cases terminating in 1937-9 compared with those terminating in 1940-1*

Durations in Years	MALES									
	Age Groups (Age at Death)									
	Under 25		25-		35-		45-		55+	
	1937-9	1940-1	1937-9	1940-1	1937-9	1940-1	1937-9	1940-1	1937-9	1940-1
Under 1	109 (41)	78 (41)	111 (37)	77 (39)	117 (43)	78 (40)	111 (46)	87 (45)	93 (49)	78 (47)
1-	124 (47)	85 (45)	153 (51)	90 (45)	111 (41)	81 (42)	100 (42)	69 (35)	73 (38)	65 (39)
5+	32 (12)	26 (14)	39 (13)	33 (17)	46 (17)	36 (19)	28 (12)	39 (20)	24 (13)	22 (12.6)
All durations	265	189	303	200	274	195	239	195	190	165
Percentage age distribution	20.8	20.0	23.8	21.2	21.6	20.7	18.8	20.7	14.9	17.5
									99.9	100.1

Durations in Years	FEMALES									
	Age Groups (Age at Death)									
	Under 25		25-		35-		45+		All Ages	
	1937-9	1940-1	1937-9	1940-1	1937-9	1940-1	1937-9	1940-1	1937-9	1940-1
Under 1	161 (45)	128 (48)	119 (38)	90 (41)	47 (35)	44 (45)	57 (47)	41 (50)	384 (41)	303 (46)
1-	152 (43)	113 (43)	155 (49)	99 (45)	68 (50)	40 (41)	51 (42)	23 (28)	426 (46)	275 (41)
5+	44 (12)	25 (9)	42 (13)	32 (15)	21 (15)	13 (13)	13 (11)	18 (22)	120 (13)	88 (13)
All durations	357	266	316	221	136	97	121	82	930	666
Percentage age distribution	38.4	39.9	34.0	33.2	14.6	14.6	13.0	12.3	100.0	100.0

* Duration is defined throughout this paper as the period elapsing between notification (registration as a tuberculous person) and death.

and death for two periods in which death occurred: (1) the pre-war years 1937-9, and (2) the war years 1940 and 1941. Shorter average duration should appear in the second period if the postulate of the preceding paragraph is correct. The term "duration of disease" as used in this analysis is defined as the time elapsing between notification (or registration as an infected person) and death.

The data are not complete; only about 75% of the deaths recorded in the County of Middlesex by the Registrar-General during these five years appear to have come under the county council's scheme for the treatment of tuberculosis.* Of the remainder a number of persons escape notification as suffering from tuberculosis until after death; and a further proportion, who were notified, were treated privately and not under the county service. It can be shown that the proportion who escaped the net was constant in both periods, and hence the comparison is not invalidated even though the accuracy of the figures as measures of the absolute duration of the disease is somewhat reduced.

* A similar test could be applied by all tuberculosis centres to furnish a rough guide as to the extent to which tuberculosis cases in any administrative area were seen by the official medical service.

groupings, in both absolute and percentage form and distinguishing age and sex, are shown in Table I; also the percentage age distribution at death of these fatal cases.

Findings

Age Distribution.—Statistical tests showed no significant difference between the age distributions in the two periods.

Duration Distribution (All Ages).—Considering first all ages combined, Table I shows that in the three years 1937 to 1939 1,271 deaths of males occurred from pulmonary tuberculosis among patients under observation by the Middlesex Tuberculosis Service, as against 944 in the two-year period 1940-1. In the pre-war period the duration was of less than one year in 43% of the deaths, as compared with 42% in the war period. Those with durations of more than one year but less than five years accounted for 44% in the pre-war and 41% in the war years; while there were 13% in the pre-war and 17% in the war years with durations ranging from five to twenty-six years. Similarly for females, 41% of the 930 deaths in 1937-9 and 46% of the 666 deaths in 1940-1 were of less than one year's duration; 46% as against 41% were of more than one but less than five years; and 13% in both periods