SOME COMMON PAEDIATRIC PROCEDURES A REASSESSMENT FOR CHILD WELFARE PRACTICE*

BY

DOYNE BELL, D.M.

Physician, Belgrave Hospital for Children; Physician in Charge of Children's Department, Queen Mary's Hospital for the East End

It has seemed to me suitable, in addressing this audience, to confine my remarks to such aspects of the subject as are the day-to-day concern of the worker in the child welfare clinic. The choice of a few points of common paediatric procedure for your reconsideration has therefore been based on cases referred to me from welfare centre practice or presented as problems to midwives and students in maternity departments. If, then, my conclusions seem to many to be self-evident, I must plead that to others they have seemed less obvious.

The Crying Baby

Handbooks on the care of babies quite rightly tell their readers that the treatment of the crying baby is to find out the cause of its crying and correct it; yet in practice it is not at all uncommon for mothers to complain that their babies cry for hours on end, and an examination of such babies shows no abnormality to account for this behaviour. Endless and apparently motiveless crying is extremely disturbing to the mother, and may result in such anxiety that lactation fails. Even without such a serious consequence the symptom is distressing both to the baby and to all who hear it, and calls for treatment. I think in many cases, though the crying was started by some momentary discomfort, its persistence is due to the discomfort occasioned by the crying itself. The baby gets hot and its pillow wet, and so the crying continues. In these cases, after excluding by thorough examination an organic cause of pain, the giving of a sedative is amply justified. It is important, however, to give it in sufficient dose to ensure that the habit of crying is broken. Nothing discourages the mother more than to be given a medicine which she hopes will soothe the infant and then to have her hopes frustrated. For an infant 6 months old, for instance, phenobarbitone, gr. 1/2, would be an appropriate dose. The sedative need not be repeated for more than three or four nights as a rule.

Breast-feeding

The commonest errors in the handling of the breast-feeding of the newborn child seem to be due to anxiety and pessimism—anxiety lest the newborn infant should be underfed in the early days of life, and pessimism with regard to the mother's ability to suckle her child. Admittedly the factors influencing lactation are incompletely understood and complex—too complex for detailed discussion here; but it is clear that successful breast-feeding does not depend entirely on mechanical factors such as the stimulation of the nipple by sucking and the emptying of the breast. Endocrine influences of much complexity play an important part, and anxiety or pessimism or a display of even cheerful ignorance by nurse, doctor, or health visitor will often effectively disturb what should be a normal lactation. My plea, therefore, is for a much more intelligent and intensive field study of lactation. Such study would show, among many things, how great is the variation in the appetites of normal newborn babies and in the time taken for lactation to become established. I am depressed at the number of infants discharged from maternity hospitals at the age of 2 weeks or less, wholly artificially fed. Often in such cases I learn that on or about the tenth day after delivery the mother has been given tablets to dry up her milk and told that she has an insufficient supply. There is nothing pathological in the fact that on the tenth day the supply of breast milk is less than 2½ fl. oz. per lb. of the infant's body weight. I think it is a fair generalization to say that, in the case of a healthy mother and infant, it is never possible in the first three weeks to state that lactation will not be established and should be abandoned. I have made it a rule, in a midwifery department where the newborn babies are in my care, that no infant is ever to be given a complementary feed (I except water and saline) without my being notified. As a result such feeding is, in normal cases, practically unknown.

In contrast let me instance a case referred to me last year. A mother aged 20 was sent with her 9-weeks-old infant to see me because he was restless and crying, and vomited. The baby had been born in a private nursing home, and had never been put to the breast. The mother had been purged and had her breasts bound up on the third day, and a variety of proprietary foods had been given to the infant. On examining the mother's breasts I found that a few beads of milk could be expressed. She was obviously disappointed that she had been thought incapable of suckling her child. I gave her the ordinary advice for re-establishing lactation,

but warned her not to be unduly depressed if she failed. She wrote 4½ weeks later to tell me that her infant was fully breast-fed. The indiscriminate use of synthetic oestrogens has of course made the re-establishment of wrongly abandoned lactation more difficult.

The study of natural lactation can be carried out only by the paediatrician who sees the baby from birth onwards. The time available to the obstetrician or midwife—two or three weeks as a rule—is too short for a proper study to be made. This, incidentally, is a further argument, if one is still needed, for the paediatric supervision of the newborn babies in maternity units. The premature termination of lactation, like the termination of pregnancy, is a step which may have serious consequences, and should never be undertaken without a full knowledge of the factors involved and a careful consideration of the medical indications.

The formation of an abscess in the lactating breast is a complication of breast-feeding that needs much study. There is apparently a widespread failure to appreciate the aetiological factors and what should be the aim of treatment. Most cases of suppurative in the lactating breast are preceded by obstruction of the milk ducts, producing a tender, knotty breast. Treatment of this condition and prophylaxis of abscess formation should clearly aim at removing the obstruction and re-establishing the milk flow. Taking the baby off the affected breast and tight bandaging are therefore to be condemned.

When an abscess has definitely formed it is still too often forgotten by the surgeon that he has two patients—the mother and the infant. No treatment of breast abscess should be considered entirely satisfactory which does not achieve as its end-result normal lactation. A fissure in the nipple should be considered a danger signal, for it may be a portal of entry for infection to the deeper breast tissues. In any event, the fissure itself is extremely painful and may render breast-feeding from the affected side temporarily impossible. If, on account of pain, breast-feeding has to be suspended, particular care must be taken, while the fissure is under treatment, to prevent milky engorgement of the breast, either by manual expression or by the use of the breast pump. The objects of local treatment of the fissure are disinfection and rest, just as if it were an ulcer anywhere else in the body. There are many ways of achieving these ends, but a very practical method, and one that gives rapid relief of pain, is the application of a lead nipple-shield between feeds.

Artificial Feeding

The technique of artificial feeding should be elastic and be modified to suit the individual infant's requirements, and in this important branch of paediatrics rule-of-thumb methods have little place. One not uncommonly sees infants suffering from dietetic upsets who have been treated by many changes of the brand of dried milk offered but none in the composition or quantity of the food. The only change has been in the branded name on the tin. The slavish belief in the advertised merits of proprietary brands of dried milk has interfered with the wider use of national dried milk. In this respect some hospitals have been much to blame. An infant is admitted to hospital and fed on a proprietary brand of straight full-cream dried milk; if he does well the mother is naturally reluctant to change to national dried milk when the infant returns home. Thus she pays at the chemist's shop three or four times as much as she would pay for the identical product at the food office or clinic.

Respiratory Infections in Childhood

Chronic nasal discharge in small infants with resulting snuffles or nasal obstruction is a very common symptom. The origin of these snuffles is probably a mild catarrhal infection of the nose, but the persistence of the symptom is often due to mistaken attempts to clear the nose with pledgets of cotton-wool. These pledgets are useless for the purpose for which they are intended, and the delicate mucous membrane of the infant's nose is always irritated and often injured by their use, and a chronic infection results. Most cases of snuffles will clear up when the use of such pledgets is discontinued.

Whether tonsils should be removed or not is a question asked frequently in welfare centre practice. It clearly requires careful and individual consideration. No one who has seen some of the bad results of indiscriminate tonsillectomy will fail to realize the importance of a right decision. The tonsil undoubtedly plays a large part in the defence of the respiratory tract of the young child against infection, and its removal while still capable of healthy function will only weaken the defence. I do not think it is an over-simplification of the problem to state that there are but two indications for the removal of the tonsils in children under 5 years old. The first is hypertrophy of the tonsil to a degree which interferes with the child's swallowing; the second is chronic infection of the tonsil, resulting either in repeated auto-infection of the respiratory tract or in the child's becoming a carrier of pathogenic organisms which cannot be eliminated by other means. It is important to remember that a history of repeated acute infections in the child is not necessarily an indication for removal of the tonsils. The repeated infections may be due simply to repeated exposure to an outside source of infection. The same principles

* Lecture (abridged) given in London to the Maternity and Child Welfare Group of the Society of Medical Officers of Health.

should be applied to consideration of the removal of adenoids. The adenoids should be removed if they are the site of chronic infection, or if their hypertrophy obstructs the nasal air-way or interferes with the drainage of the paranasal sinuses or middle ear.

In the prophylaxis of respiratory infection in childhood it is still not uncommon to find that clinics are exposing children to ultra-violet light. The uselessness of this procedure was well shown by Dora Colebrook as long ago as 1929. She studied three groups of children, of which one was exposed to ultra-violet light, another to a screened lamp from which no ultra-violet rays came, and the third served as a control. She found no significant difference in the incidence or duration of respiratory infections in the three groups; the very small and insignificant difference being favourable to the control group. In the absence of any work to contradict these findings it seems to me that the exposure of children to ultra-violet light is quite unwarranted.

Strawberry Naevi

The strawberry naevus or capillary haemangioma is quite a common birthmark, and a great number of cases are still referred to hospital, often with the suggestion that they should be treated with CO₂ snow, with radium, or by excision. The natural history of these naevi was studied by W. A. Lister (1938), and he found, in a series of 77 children with 93 naevi who were watched for periods varying from one to seven years, that there was no exception to the rule that naevi which grew rapidly during the early months of life subsequently retrogressed and, of their own accord, disappeared, on the average at about the fifth year.

My experience is in accordance with Lister's findings. Any form of treatment is bound to leave a scar, whereas natural retrogression of the naevus leaves normal healthy skin. The chief difficulty in practice is to persuade the child's mother that the naevus will retrogress after reaching its maximum size at the age of about 12 months. My practice is to make an exact tracing of the naevus on a piece of "cellophane" or washed x-ray film and transfer this pattern to paper. This paper is dated and given to the mother to keep. She may then observe for herself that the naevus does not further increase in size, but gradually shrinks or fades. In many cases the retrogression starts in patches in the middle of the naevus, but after the age of about 12 months the edges, even if they are the last bits to disappear, do not spread.

Circumcision

This mutilation has a long history which is of great anthropological interest. But its popular justification on hygienic grounds seems to be of quite recent origin and of extremely doubtful validity. As it is not free from hazard and discomfort it is well, when considering it as a surgical procedure, to take stock of the indications for the operation. So far as I know, there is but one—phimosis; this being defined as a stenosis of the preputial orifice which prevents retraction of the prepuce behind the corona of the glans penis. In most cases in which circumcision is performed no attempt is made to assess whether phimosis is present or not; for normally the prepuce of the infant is lightly adherent to the glans, and it is these adhesions which prevent retraction. If the adhesions are divided—and this can easily be done either by digital retraction or with a probe, though I do not suggest that it is a desirable routine procedure—it will be found that phimosis is rare. If the preputial orifice is smaller than the urinary meatus—an extremely rare occurrence—interference with urination will occur and operation is certainly necessary. But I doubt whether operation is indicated in lesser degrees of phimosis in early infancy. Indeed, it has been suggested that some degree of phimosis, as above defined, is normal in the young infant.

I have heard it suggested that a long and inflamed prepuce or the so-called "cauliflower prepuce" indicates circumcision. This condition is in fact an absolute contraindication to circumcision. The "cauliflower prepuce"—the site of cellulitis, oedema, and even ulceration—is the result of an ammonia dermatitis. To remove such a prepuce is to expose the more delicate glans and the urinary meatus to the same trauma with a far more painful result—namely, meatal ulceration. One might as well urge a man to remove his steel helmet in an air raid because it was being dented by falling "flak."

REFERENCES

- Colebrook, Dora (1929). *Irradiation and Health*, Med. Res. Cncl. Sp. Rep. Ser. No. 131, London.
Lister, W. A. (1938). *Lancet*, 1, 1429.

P. A. O'Leary and W. E. Herrell (*Proc. Mayo Clin.*, 1944, 19, 20) record the treatment of a case of late cutaneous syphilis with penicillin. The patient was a 42-year-old married woman who for eight years had had a nodular syphiloderm of the nose. She was given 20,000 units of penicillin intravenously twice a day for 8 days. The result is described as striking, and three weeks after the course of treatment all that was left was some residual pigmentation. Serum tests for syphilis remained unchanged.

THE LISTER INSTITUTE

Long-term Investigations

The report of the Governing Body of the Lister Institute of Preventive Medicine, given at the recent annual meeting, contains brief accounts of many continuing investigations which cannot be expected to reach finality in any one year. Some of the members of the staff are on active service abroad or seconded to a Government Department. A number of the workers are enjoying research facilities at Oxford, and others at Cambridge. One event of the year was the establishment by the Medical Research Council, which is administering the London blood supply depots on behalf of the Ministry of Health, of a unit at the Institute for research into, and filtration of, blood plasma and serum for transfusion, and the Director of the Institute, Dr. Alan N. Drury, is chairman of the Blood Transfusion Research Committee.

Two of the workers, Dr. R. A. Kekwick and Dr. M. E. Mackay, have been studying problems arising in the large-scale processing of human serum and plasma. They have found that the adjustment of the reaction of plasma to pH 5.4 with sterile citric acid before ether extraction provides a method for the quantitative removal of fibrinogen, as well as for removing the unstable lipid complexes. After readjustment of the product to pH 7 followed by Setz filtration, a transfusion fluid is obtained which remains stable for at least twelve months. It is added that problems arising in the kaolin processing of plasma have been surmounted and the Serum Unit has been filtering the entire output of the London blood transfusion depots before dispatch to Cambridge for freeze-drying.

Serological and Bacteriological Studies

One of the serological studies undertaken concerns the treatment of louse-borne typhus by anti-proteus OX 19 serum. Dr. A. Felix has been working on this serum from the horse, and a few preliminary trials have been carried out in North Africa in collaboration with Capt. T. E. Woodward of the United States Typhus Commission. Six selected severe cases were treated; three patients received the serum intravenously and three intramuscularly. All did well, though one of them had been in a comatose state for several days. These trials have shown that large doses of the serum are well tolerated, and that as a result of the injections an initial low-titre OX 19 reaction in the patient's serum can be increased from twenty to thirty times.

Dr. Felix and Miss Callow have continued the typing of typhoid, paratyphoid, and food-poisoning bacilli with the Vi bacteriophage. A paper by them on the typing of paratyphoid B bacilli in this way was published in the *British Medical Journal* of July 31, 1943 (p. 127). One hitherto unknown Vi-phage type of the typhoid bacillus was identified during the past year as being responsible for a small outbreak of typhoid fever. This type does not seem to be indigenous to the British Isles, as it has been traced to a chronic carrier who acquired the infection during the war in South Africa more than forty years ago. On the other hand, a Vi-phage type of *Bact. paratyphosum* B has been found which must be considered indigenous. The number of known Vi-types and subtypes of the paratyphoid B bacillus is now five.

Nutritional Studies

The Ministry of Food has been anxious to ascertain more accurately the requirement for vitamin A as derived from pre-formed vitamin A and from the provitamin carotene. This has been done by a human experiment carried out on volunteers, who were conscientious objectors, at the Sorby Institute at Sheffield. Since July, 1942, these volunteers have received a diet made as deficient as possible in vitamin A and carotene; a small number have served as positive controls receiving supplements of these substances. Observation was specially directed to the measurement of capacity for dark-adaptation. At the end of a year none of the volunteers showed any definite signs of deterioration of the visual threshold, but after fourteen months one subject began to show this deterioration, with slight skin symptoms, and a very low value for plasma vitamin A, all of which signs progressed rapidly until he was given a small daily dose of the vitamin, when the symptoms slowly regressed. The experiment has indicated how large are the vitamin A reserves in the liver of the healthy human adult,