ment in favour of the rejection or early elimination of the vulnerable. There is no doubt whatever that those who do succumb show a high concentration of certain ominous points, such as a poor work record, compared with those who do not. According to certain investigations that have been made in the Navy, the proportion showing these ominous points amounted (with the technique used) to roughly 60% of psychiatric in-patients and 50% of psychiatric out-patients, as compared with only 15% among "normal" sailor controls. Yet naval psychiatrists keep 9 out of 10 of the men referred to them as out-patients at duty and send back to duty 2 out of 3 of their in-patients. Moreover, the relapse rate as judged by readmission to hospital does not seem unduly high. Thus, according to an investigation made last year, psychiatric cases were being returned to duty from hospital at about 5 times the rate that relapsed cases were being readmitted, and about half of these relapsed cases had remained outside hospital for more than 6 months. Finally, of those that are invalided for psychiatric disorders, two-thirds have survived in the Navy for over a year.

The significant thing about these findings seems to be that although a high proportion of those who break down not only give a past history which suggests their predictability as bad psychiatric bets, but also show that they were bad bets by having broken down, yet it is still possible to send the majority of them back to duty and to retain a relapse rate that seems reasonably small. This is, of course, not to doubt the desirability of selection procedures. The standard adopted for rejection or early elimination must naturally depend upon the available man-power; but a number do remain who should be rejected, no matter what the man-power position may be. It seems doubtful, however, whether in the Navy the proportion that should certainly have been rejected or eliminated amounted at most to more than 1 to 2%. This may seem a small number, yet the trouble these men cause is out of all proportion to their actual numbers, when considered in terms of hospital treatment, inefficiency, letting down others, and impairing morale. A great stand-by in the Navy has doubtless been the self-selection consequent upon men expressing a naval preference at the time of their call-up, for reluctant dragons do not make the best fighting material.

It seems reasonable to conclude that many "bad psychiatric bets" give very fair service before they break down and are also capable of giving further fair service after they have done so. These facts should counteract any undue pessimism that may exist as to the capacity of the neurotic to stand up to the stress of war. There can be little doubt that an additional factor inducing unwarranted pessimism is the abnormal or at least different situation that occurs in shore establishments, and particularly in hospital, compared with life afloat. The picture seen in hospital is especially misleading, and makes a correct assessment extremely difficult.

Treatment

Apart from the real, though limited, place that exists for pre-selection, probably the most that would be desirable prophylactically would be a warning to survivors and others going on leave from sea that anxiety would be experienced temporarily and should not be misinterpreted as illness. The vast majority of men, however, deal with their experiences quite adequately without outside help, and certainly any prophylactic "pep talks" of the above kind should be given by executive officers and not by medical men.

Once a pathological tension state has become established it must be dealt with either by reducing the tension or by raising the tempo. The latter can be accomplished by an early draft to sea before habituation, including doubts about future capacity, has occurred. Failure may result if the ship lies in harbour for too long a period.

Reduction of tension can be produced by sedation, and the barbiturates are eminently suitable for this purpose. Tremulous agitated patients should be put to bed, and 48 hours' sedation will make discharge to hospital unnecessary in the overwhelming majority. The average case seen as an out-patient in a naval depot shows a steady decrement of tension at a rate which is extraordinarily consistent. The administration of pheno-barbitone in three doses of half a grain daily for one week,

followed by the same dose twice daily for two weeks, usually brings about clinical recovery. With the other factors in treatment and management this paper is not primarily concerned. It may be worth noting that what medical men now effect with barbiturates humanity has accomplished for hundreds of years with the aid of alcohol. It is a sobering reflection that humanity was abreacting its pent-up emotions in parties with the aid of alcohol long before narco-analysis and group therapy were introduced for the same purpose.

"All theories are crutches, not sceptres or wizards' wands," and it is not suggested that the concept of post-operational strain that has been outlined is the whole story, but merely that it is perhaps a somewhat neglected part of it. Numerous other factors obviously help to make stress more tolerable and hence affect the incidence of neurosis, such as discipline, leave, mail, and the rapid solution of what are felt to be inequalities of treatment—to mention only a few. The Navy is, moreover, singularly fortunate in having a tangible object which is also a home around which group loyalty can centre—namely, the ship. An airman or member of the Tank Corps cannot feel about their contraptions in quite the same vividly personal way. Yet the life of a closed community, as in a ship, has its special dangers. Interesting browned-off (what the sailors call "chocker") paranoid reactions, rather similar to those seen among prisoners, may occur when the commission is too long or the ship is not a happy one. Dr. Johnson's dictum may be recalled: "No man will be a sailor who has contrivance enough to get himself into a jail; for being in a ship is being in a jail with the chance of being drowned." The rest of the quotation—"a man in jail has more room, better food and, commonly, better company"—is perhaps less acceptable.

There can also be little doubt that the group spirit and loyalty that exist on board ships are most potent factors in sustaining morale and that, conversely, the lack of this same group spirit and loyalty when a man leaves his ship and goes to a drafting depot is a most important factor in precipitating the breakdowns that occur there with so much greater frequency.

Summary

An endeavour has been made to show that psychiatric casualties attributable to operational stress occur most commonly after rather than during the stress.

Thus it has been pointed out that the psychiatric sickness rate is very much greater ashore than afloat.

The thesis is submitted that this is often due to a persistence of the physiological increase in tension developed at sea and rendered obtrusively uncomfortable by the suddenly reduced tempo on coming ashore; the reaction being analogous, in slow motion, to the normal experience of feeling worse after danger than during it.

While unworthy motives play an important part in perpetuation, their role in producing the state can easily be over-estimated.

While pre-selection has a definite value, the practical importance of selection procedures is much modified by the fact that a substantial majority of psychiatric casualties can be returned to full duty.

Symptomatic relief often follows reduction of tension by sedation and increasing the "tempo" by early draft to sea before habituation has occurred.

SCHICK REACTIONS IN RECENTLY CONFINED WOMEN AND THEIR INFANTS

BY

G. PAYLING WRIGHT, D.M., F.R.C.P.

AND

W. M. CLARK

(From the Department of Pathology, Guy's Hospital Medical School)

It is now generally recognized that the resistance of newly born infants to diphtheria depends chiefly upon the passage of maternal antitoxin through the placenta during the latter part of gestation. So readily can this antibody enter the foetal circulation that at birth its concentrations in the serums of mother and infant are almost equal. From early childhood onwards any insufficiency in the concentration of circulating antitoxin to ensure immunity to clinical diphtheria may be readily demonstrated by a positive reaction in the Schick test.

In very young infants, however, owing to certain physiological peculiarities in their skin, positive Schick reactions in susceptible individuals are much less readily evoked, and estimates of immunity obtained by the use of this test during the first few months of life are thus often fallacious. Consequently, the Schick reaction of the recently confined mother, which for practical purposes may be accepted as reflecting the antitoxin titres of both her own and her infant's blood, thus acquires a particular significance in the epidemiology of diphtheria in early infancy.

The following study was made during 1942 and 1943 in a maternity department of a hospital on the south-eastern outskirts of London. A large proportion of the 250 recently confined women examined came from districts in which housing conditions are good. Many of them, however, had spent the earlier part of their life as school-children in one of the Metropolitan boroughs of London, and, since it seemed possible that this might have had some effect upon their immunity as a group, the findings for these women have been recorded separately. The average age of the 250 women was 28.5 years. The Schick tests were carried out according to the usual technique 0 to 5 days after delivery. The sites of inoculation were examined daily for 7 days. The results of the tests are shown in Table I.

TABLE I.—Schick Reactions (Totals and Percentages)

Groups of Women Examined	+ve	ψ+ve	—ve	ψ—ve	Totals
London-bred ..	54 (45.7)	10 (8.5)	43 (36.4)	11 (9.4)	118
Others ..	75 (56.8)	5 (3.8)	45 (34.1)	7 (5.3)	132
Totals ..	129 (51.6)	15 (6.0)	88 (35.2)	18 (7.2)	250

Schick Reactions in Non-parturient Women

Since it seemed possible, from these rather high rates for positive reactions among recently confined women, that such persons might be more sensitive to diphtheria toxin in the puerperium than at other times, a further series of tests were made upon a group of 62 women who had been admitted to the same hospital for medical or surgical treatment. Their social surroundings were the same, and their average age—31.1 years—was nearly the same. Of this group 58% proved to be either positive or pseudo-positive reactors, so that there seems to be no reason to believe that the puerperal state gives rise to any heightened susceptibility to this toxin. The London-bred women in this group also showed a rather lower proportion of positive and pseudo-positive reactors (52.6%) than the remainder.

Positive Reactors in Recently Confined Women in Various Age Groups

In the group of 250 recently confined women there were statistically significant differences between the proportions of positive reactors in the following three subgroups: those 24 years and under; those between 25 and 34 years; and those 35 years and over. The numbers falling into these subgroups, the numbers showing positive or pseudo-positive reactions, and their respective percentages are set out in Table II.

TABLE II

Age Group	No. Tested	No. Positive	% Positive
24 years and under ..	88	60	68
25 to 34 years ..	116	67	58
35 years and over ..	46	17	37

Eighteen of these 250 recently confined women and 22 of the whole group of 312 stated that they had suffered from clinical diphtheria; 14 of them were among the 137 London-bred women—a proportion which might be expected from the ratio of notified cases to child population in the Metropolitan boroughs between 10 and 20 years ago (see Wright and Wright, 1942). Of these 22 women 12 reacted negatively to the Schick test, 7 positively, and 3 pseudo-positively. Only 5 of the 312 women had been immunized prophylactically against diphtheria; 4 of them had a negative and one had a pseudo-positive reaction.

Changes in Schick Reactions at Short Intervals after Injection of Toxoid

It appeared unlikely that so large a proportion of women living in a semi-urban environment could have escaped immunization through casual contact with diphtheria cases or carriers at some period of their lives. It seemed much more probable that most of them had undergone some degree of immunization in the past, but that the exposure was too slight or at too remote a date to maintain their circulating antitoxin titre at or above the critical level for the Schick test. To examine this possibility some of the positive reactors were injected intramuscularly with 1 c.cm. of T.A.F. on the third or fourth day after their Schick test. A second Schick test, in which toxin alone was used, was carried out four or six days after this immunization. The diameters of the hyperaemic areas present at the peak of the responses were measured, and in most instances the relative intensities of the first and second reactions were noted roughly for comparison. With the help of this combination—the comparatively large “recall dose” of toxoid and the two Schick tests—it was hoped to distinguish those women in the two subgroups: (a) “fully susceptible” and “sub-immune,” from (b) “positive immune” and “potentially immune” (see Glenny, 1925).

Of the two subgroups of 20 women each whose tests were repeated 4 and 6 days after receiving the “recall dose” of toxoid, 10 and 9 respectively showed second reactions two-thirds or less the size of those originally found. In the 6-day group, possibly owing to the rather longer interval, the intensity of the hyperaemia in the second test was in almost all instances notably less than in the first. Such changes in the size and intensity of the reaction almost certainly indicate that the second injection of toxin had been more fully neutralized than the first, and consequently that an appreciable amount of antitoxin had been formed actively by such women—a response to this antigenic stimulus that would be unlikely to take place so promptly except in the tissues of an already partially immunized person.

From these observations it seems likely that at least half, and perhaps more, of the positive reactors among the recently confined women fell into the categories termed by Glenny the “positive immune” and the “potentially immune.” At some time in the past they would probably have reacted negatively to this test, but, possibly owing to long residence in less densely populated districts, they have relapsed into the positive state.

Schick Reactions in the Infants

Schick tests were also carried out at some time during the first 5 days of life upon the infants of 145 of this group of women. The injections were made either in the scapular region or on the outer aspect of the forearm near the elbow. All the test sites were examined daily for 7 days, except in a few

TABLE III

	Infants All Under	Mother +ve Infant +ve	Mother —ve Infant —ve	Mother +ve Infant —ve	Mother —ve Infant +ve
Ruh and McClelland (1923)	7 days	20	75	4	1
Kuttner and Ratner (1923)	14 ..	4	39	7	0
Cooke and Sharma (1932)	10 ..	59	204	34	1
Greengard (1932)	14 ..	22	45	33	0
		Agreement		Disagreement	
Totals ..		468		80	
Percentages ..		85.4		14.6	
Present series ..	5 days	66	56	21	2
Totals ..		122		23	
Percentages ..		84.1		15.9	

instances in which the mother or infant had left the hospital on the fifth or sixth day. In the infants no reaction was regarded as positive unless it was seen to be present on at least two such examinations. The findings in the present series, together with those of certain earlier workers, are shown in Table III.

In 122 of the 145 mothers and infants tested their reactions agreed. In the remaining 23 there was disagreement—the mother being negative and the infant positive in 2 instances, and the reverse in 21 instances. The circumstances under which these disagreements occurred were as follows:

Mother Negative and Infant Positive (2 instances).—In the first of these the mother was pseudo-negative and the infant reacted only faintly on the second and third days. In the other the woman was definitely negative and the infant definitely positive. It may be observed that instances are recorded in which a relatively high antitoxin titre in the maternal blood is accompanied by a much lower, or even untitratable low, level in the cord blood. Von Gröer and Kassowitz (1915) found three such instances in their series.

Mother Positive and Infant Negative (21 instances).—In 6 of these instances the mothers were definitely positive. Three of their infants showed a positive reaction on one day only, and were accordingly classified as negative; in one other instance the mother had had diphtheria 24 years before. In 5 other instances the mother was classified as a faint reactor, and in none did the infant show any sign of a positive reaction for even so short a period as 24 hours. In the remaining 10 instances the mothers, three of whom had had diphtheria in childhood, showed pseudo-positive reactions. One infant in this group showed a positive reaction on one day only.

The most common form of disagreement was thus a failure of the infant to react positively when the mother was positive, and especially when she was either a faint or a pseudo-positive reactor. This state of anergy was present in about one-quarter of the infants in whom a positive reaction might have been expected from the result of the test upon the mother. The proportion is rather smaller than that found in the earlier studies, and is probably due to the frequent examination of the test sites in the present series. Moreover, the lighting conditions for examining the present group of infants were unusually favourable.

Discussion

The belief is still widely held that the great majority of young infants possess a high degree of congenitally acquired passive immunity to diphtheria for some months after birth. Two circumstances in particular have contributed to this belief. First, many of the observations upon specific immunity in newly born infants were made in Vienna and other European cities, in whose populations diphtheria was particularly prevalent, and the results of these early surveys have been applied rather freely to other places and times without allowing adequately for local differences in the endemicity of the disease. Secondly, insufficient attention has been paid to the phenomenon of skin anergy which is shown by young infants to diphtheria toxin and other noxious agents capable of producing delayed erythematous reactions, so that, for this age group, many negative Schick reactions have been interpreted erroneously in consequence. Together, these misconceptions have led to an exaggerated estimate of the prevalence of congenital immunity during the first few months of life, and the reliance which has been placed upon this supposed insusceptibility is unfounded in a large proportion of instances.

The present observations indicate that considerably less than half of the infants born to women living in a typical outer London suburb have any adequate congenitally acquired immunity to diphtheria. So infrequently do the mothers come into contact with toxigenic *C. diphtheriae* that their circulating antitoxin concentrations fall to levels considerably below the critical titre for the Schick test. While a large proportion of the positive reactors among these recently confined women are themselves immune in consequence of the promptitude with which their antitoxin-forming tissues respond to the secondary stimulus of the specific toxin, their low antitoxin titre during pregnancy confers little or no immunity upon their infants.

At the present time the trend of mortality of infants under 1 year from diphtheria in England and Wales is a declining one, and so long as this fall continues no additional prophylactic precautions seem to be required. In the event of an epidemic rise, however, either in some particular locality or in the country as a whole, two possible measures might be instituted. First, infants might be immunized considerably younger than is now the custom: this procedure has the disadvantage that very young infants seem to respond less well to prophylactic immunization than older infants or young children. Secondly, as Dudley (1934) suggested, it might be desirable to immunize all Schick-positive pregnant women during the latter part of the

gestation period. Such a procedure would at the same time immunize any woman who was susceptible and raise the maternal antitoxin titre to a high level about the time of the birth of the infant.

REFERENCES

- Cooke, J. V., and Sharma, B. M. (1932). *Amer. J. Dis. Child.*, **44**, 40.
Dudley, S. F. (1934). *Med. Res. Cncl. Sp. Rep. Ser. No. 195*, London.
Glenny, A. T. (1925). *J. Hyg., Camb.*, **24**, 301.
Greengard, J. (1932). *Arch. Pediat.*, **49**, 526.
Kuttner, A., and Ratner, B. (1923). *Amer. J. Dis. Child.*, **25**, 420.
Ruh, H. O., and McClelland, J. E. (1923). *Ibid.*, **25**, 59.
Von Gröer, F., and Kassowitz, K. (1915). *Z. Immunforsch.*, **23**, 108.
Wright, G. Payling, and Wright, H. P. (1942). *J. Hyg., Camb.*, **42**, 451.

RADIOTHERAPY OF ECTOPIC CALCIFICATION

BY

E. MILLINGTON, D.M.R.

Radiotherapist to the Royal West Sussex Hospital, Chichester

For some years radiotherapists have been treating with success deposits of calcium in the tissues, but unfortunately few outside the specialty seem to know of them. In two recent books on bone and joint conditions (McMurray, 1937; Watson-Jones, 1943) no mention is made of radiotherapy at all in this connexion, and few writers on general surgery have commented on this mode of approach, although in America the use of x rays in treatment is much more widely appreciated, and standard works, such as that by Comroe (1941), give full credit to radiotherapy. Ectopic calcium deposits may appear almost anywhere in the body, but of most interest here are those occurring in muscles, tendons, and bursae, as such deposits are in many cases painful and thus attract attention and demand treatment. The shoulder region is often affected, as in the calcified subacromial bursa or supraspinatus insertion, and the olecranon and Achilles bursae are also quite often involved in this process.

No convincing explanation of this deposition has as yet been given, and the formation of actual bone is even more obscure. Many theories have been put forward on the subject; these are summarized by McCurich and Millington (1943). It is only with calcification, and not with true ossification, that we are now concerned, as x-ray treatment has no effect on formed bone, but only on calcium deposits. Calcium is frequently deposited in dead or devitalized tissue, possibly because of the change in pH in such regions, and the lesions with which we are dealing are mostly traumatic in origin, with pain dating from some injury or strain and becoming more persistent with time and the usual methods of treatment. X-ray examination eventually shows the presence of shadows due to the deposition of calcium of an amorphous and flocculent nature. It seems questionable whether it is the calcium that causes the pain, but in most cases if the calcium can be removed the pain also goes to a large extent, and perseverance with physiotherapy will then complete the cure.

Many surgeons recommend and practise surgical removal of the calcium. It was sometimes found at operation, however, that what appeared radiographically to be a large mass of chalk could not be distinguished from surrounding tissues when exposed, and at other times only a small amount of chalky fluid was seen; so there is no guarantee of cure from operation. X-ray treatment offers a method which is devoid of trauma and risk, and in any case does not preclude any more drastic methods later should it fail. The mode of action of the rays is not known, any more than it is with other applications; it may be due to alteration of blood supply, or quite possibly to some change in the pH of the irradiated part. The dosage is small and not harmful, there are no known ill effects, and the patient is not laid up or incapacitated in any way; but it must be realized that the effects of x rays are not usually manifest for some time after the treatment has been given, and as a rule it is some four to six weeks before there is any change in the radiographic appearances even if the pain has been relieved earlier.