hereditary or acquired, temporary or permanent, and that it has been described in conjunction with various factors, local and general, thought to be of possible significance. Among other conditions we find thyrotoxicosis named more than once (Heller, 1927; Rossegger, 1936); curiously, thyroid deficiency has also been blamed (Schönstein, 1929). Reference may be made to the useful general accounts of Heller (1927), Cipollaro (1930), and Rossegger (1936).

The other chief point we desire to stress is that, in temporary koilonychia, attempts to associate the nail change with any particular factor should take into account the abruptness with which the nail growth may lose its abnormality, while the change in the nail, which has been the result of earlier abnormal growth, may continue to be shown for many weeks or perhaps months. Thus it is clear that the presence of a concave surface does not mean that the process responsible for it is still active; indeed, when the concavity is at the distal end of the nail it is probable that the cause has been for a long time in abeyance—a fact to which Toegel (1937) draws attention in his case which followed brief exposure of the nails to benzene.

Our thanks are due to Sir Thomas Lewis for his assistance with these cases, and to Prof. H. P. Himsworth and Dr. Kenneth Harris, of University College Hospital, for permission to use the case reports.

REFERENCES

- Cipollaro, A. C. (1930). *N.Y. St. J. Med.*, **30**, 380.
Heller, J. (1927). *Jadassohns Handb.*, p. 115, Berlin.
Rossegger (1936). *Med. Welt*, **10**, 1803.
Schönstein, A. (1929). *Zbl. Haut- u. GeschKr.*, **28**, 258.
Toegel (1937). *Derm. Z.*, **75**, 1.

HISTOLOGICAL EFFECT OF INJECTION OF MEPACRINE (ATEBRIN) DIHYDROCHLORIDE*

BY

FRANK HAWKING, D.M.

(From the National Institute for Medical Research, London)

Recently a note was published (Hawking, 1943) concerning the histological effects produced by the intramuscular or subcutaneous injection of mepacrine methanesulphonate (atebrin musonate). However, the preparation most commonly issued for injection in the Army is the dihydrochloride (atabrine dihydrochloride). An investigation has therefore been made to see whether the effect of the dihydrochloride is significantly different from that of the methanesulphonate, and the opportunity has been taken to compare both of these preparations with quinine dihydrochloride, which is the preparation of quinine most commonly recommended for injection.

A 4% solution of quinine dihydrochloride has a reaction of pH 3.08. A 10% solution of mepacrine methanesulphonate has a reaction of pH 4.0; and one of 5%, pH 4.24; a 3.3% solution of mepacrine dihydrochloride has a reaction of pH 4.40. All these reactions were measured at 17° C. There is consequently no substantial difference in the acidities of the two solutions of mepacrine, and both are somewhat less acid than that of quinine dihydrochloride. The chief physical disadvantage of mepacrine dihydrochloride lies in its lower solubility. The methanesulphonate is soluble in water up to 33%, but the dihydrochloride only up to about 3.3% at 17° C. To obtain a 10% solution the mixture must be cautiously warmed to about 50° C. (decomposition occurs on boiling), and then the solution is cooled to 37° C. and injected without delay.

Technique

Mepacrine dihydrochloride (atabrine dihydrochloride, Winthrop), 43 mg. in 0.5 c.cm. of warm water, or mepacrine methanesulphonate (May and Baker), in equivalent amount (50 mg.), was injected intramuscularly into the thigh or loin of rabbits; some injections were also made subcutaneously on the abdomen. Similar injections of quinine dihydrochloride (Howards), 20 mg. in 0.5 c.cm., were given to other rabbits: in the previous work it had been found that the toxic action of 20 mg. of quinine was approximately equal to that of 50 mg.

of mepacrine. The animals were killed at suitable intervals (1 to 10 days); the sites of injection were fixed in formalin and studied by the usual histological technique. Altogether mepacrine dihydrochloride was injected into 11 sites intramuscularly and 4 sites subcutaneously, and mepacrine methanesulphonate into 4 sites intramuscularly. Quinine dihydrochloride was injected into 7 sites intramuscularly and 3 sites subcutaneously.

A parallel series of observations were made with rats: 4.3 mg. of mepacrine dihydrochloride in 0.3 c.cm. of water was injected intramuscularly into the thigh and subcutaneously on the abdomen, respectively, in 8 sites of each type; 5 mg. of mepacrine methanesulphonate was injected intramuscularly and subcutaneously into 3 sites of each type; and quinine dihydrochloride 10 mg. in 0.3 c.cm. of water was similarly injected into 8 sites of each kind. The dose of quinine was twice that of the mepacrine, since the therapeutic dose of quinine (0.6 g.) is usually at least twice that of mepacrine (0.3 g.).

Experimental Results

Naked-eye Appearances.—When mepacrine dihydrochloride was injected into the lumbar muscles of rabbits a fusiform lesion was produced about 4 cm. in the long axis of the muscle, by 2 by 1 cm. The lesion, on section, was dark red surrounded by a narrow yellow or whitish zone (leucocytes). In the thigh the connective tissues along the sciatic nerve were usually swollen, and often there were haemorrhagic patches in the adjacent muscles, but the appearances were less regular than those in the loin. The thigh is much less suitable than the loin for observing the effects of intramuscular injection, since so many fascial planes are present that the injected material is deposited chiefly in the connective-tissue spaces between the muscles rather than inside the muscles. In the thigh of rats there was occasionally some yellowness and oedema of the connective tissue around the chief vessels and nerves, and in one animal killed after seven days there was contracture of the muscles; but on the whole the macroscopical changes were inconspicuous. The lesions produced by mepacrine methanesulphonate were indistinguishable from those described above, and so were those produced by quinine dihydrochloride (apart from the absence of any yellow staining).

When mepacrine dihydrochloride or methanesulphonate was injected subcutaneously it caused great thickening of the skin, extending 4 to 9 cm. and due to yellow gelatinous oedema of the deep connective tissue; later a hard necrotic scab was formed. With quinine dihydrochloride the swelling was less marked and there was no yellowness.

Since mepacrine is strongly fluorescent some of the animals were examined under ultra-violet light; diffuse yellow fluorescence was seen in the tissues around the site of injection, especially subcutaneously, but the distribution could not be definitely mapped out by this method.

Microscopical Appearances.—When mepacrine dihydrochloride was injected into the compact muscles of the loin an area of coagulative necrosis with haemorrhage was produced, as described above. A short distance inside the margin of this area there was a zone of disintegrated leucocytes (seen macroscopically as a yellow band), but the necrosis extended outside this zone, and the demarcation between necrotic and living muscle was initially indistinct. After some days there was a second cellular reaction at the demarcation line, and in 10 days a definite layer of granulation tissue was formed in which were embedded shrunken muscle fibres, a few giant cells, and new collagen. In the looser tissues of the thigh the main reaction occurred in the intermuscular connective tissue, which was initially greatly distended with fluid and infiltrated with polymorphs and mononuclears. Haemorrhages and collections of albuminous exudate were also present, and most of the tissue appeared necrotic. The superficial layers of the adjacent muscles were necrotic to a variable depth and the connective tissue inside the muscle often contained fluid and leucocytes; haemorrhage was sometimes present. After from 3 to 7 days some of the necrotic muscle fibres became calcified, and the necrotic areas were separated from the healthy tissue by a zone of young granulation tissue containing many mononuclear cells. In the thighs of rats the appearances were similar. The lesions caused by mepacrine dihydrochloride were indistinguishable

* Report to the Malaria Committee of the Medical Research Council.

able from those caused by the methanesulphonate, which were described in the previous note and which were confirmed in the present work. The lesions produced by quinine dihydrochloride could not be definitely distinguished from those due to mepacrine; at the margin between the surviving and the swollen coagulated muscle fibres, groups of shrunken basophilic fibres were found, but these also occurred to a less extent in some of the sections of tissue with mepacrine. The necrosis produced in the loin by 20 mg. of quinine was not quite so extensive as that produced by 43 or 50 mg. of mepacrine, but that produced by 10 mg. of quinine in the thigh of rats was not conspicuously greater than that due to 4.3 or 5 mg. of mepacrine.

When mepacrine dihydrochloride or methanesulphonate was injected subcutaneously into rabbits or rats it caused necrosis of the epidermis, subcutaneous layer of muscle, deep connective tissue, and part of the muscle of the abdominal wall. The deep connective tissue was greatly distended with fluid and contained variable numbers of cells; at first these were principally polymorphs, but after a few days mononuclears predominated. By seven days the necrotic area was surrounded by a zone of granulation tissue. With subcutaneous injections of quinine dihydrochloride (20 mg.) the oedema, necrosis, and cell infiltrations were somewhat less extensive than with mepacrine (43 mg.). In rats 10 mg. of quinine caused somewhat less swelling and cell infiltration than 4.3 mg. of mepacrine, but the necrosis was equally extensive.

Pieces of tissue injected with mepacrine were fixed in formalin, cut as frozen sections, and examined on a fluorescence microscope. The necrotic muscle fibres shone bright yellow; connective tissue and the outer coat of blood vessels from the areas of necrosis shone a little; other tissues were hardly visible.

Discussion

From the above experiments it is clear that there is no difference in the local toxicity of mepacrine dihydrochloride and that of mepacrine methanesulphonate, although methanesulphonate is more convenient to use owing to its greater solubility. The necrosis caused by mepacrine is indistinguishable in type from that caused by quinine dihydrochloride. The relative amounts of necrosis which will be produced in man by therapeutic injection of the two compounds are difficult to predict from animal experiments, since the size of the lesion is not proportional to the dose. As described above, when given by intramuscular injection 43 mg. of mepacrine caused somewhat more necrosis than 20 mg. of quinine; but the necrosis produced by 4.3 mg. of mepacrine injected into the thigh of rats was not conspicuously less than that produced by 10 mg. of quinine. When injected subcutaneously into rats, 4.3 mg. of mepacrine produced more swelling and cellular reaction than 10 mg. of quinine did, but the necrosis was no more extensive. Since the therapeutic dose of mepacrine by injection is only one-half to one-third that of quinine, it is probable that the necrosis produced by intramuscular injection of mepacrine will be less extensive than that produced by quinine; and this conclusion agrees with the clinical reports reviewed in the previous paper. Nevertheless it is clear that the local damage produced by intramuscular injections of mepacrine is far from negligible.

Mepacrine given by intramuscular injection is rapidly absorbed. As concluded in the previous note, the damage caused is not sufficient to contraindicate its use if the patient's life or health is in danger, but it is great enough to make this method of administration inadvisable if the patient can possibly take the compound by mouth. Intravenous injection of mepacrine has been found in American experience to be dangerous to life, and its use has been condemned. Subcutaneous injection of mepacrine is not recommended under any circumstances; it was included in the present experiments for technical reasons as being a sensitive method of displaying the effect of injected substances upon the tissues.

Summary

The histological effect of intramuscular and subcutaneous injection of mepacrine dihydrochloride (atabrine dihydrochloride) in rabbits and rats was compared with those of mepacrine methanesulphonate and of quinine dihydrochloride. Mepacrine dihydrochloride has the

same acidity in solution as the methanesulphonate, but it is less soluble (solubility 1 : 30 at room temperature) and therefore slightly less convenient to use; the local toxicities of the two compounds are indistinguishable one from another. The injection of therapeutic doses of mepacrine produces necrosis of the tissues around the site of injection; this necrosis is identical in type with that produced by the injection of therapeutic doses of quinine dihydrochloride, but it is probably somewhat less extensive.

It is considered that mepacrine dihydrochloride is suitable for administration by intramuscular injection, although it is less convenient than mepacrine methanesulphonate; but this route should not be used if it is practicable to give the compound by mouth. Mepacrine should not be given by subcutaneous or intravenous injection.

Grateful acknowledgments are due to Mr. F. J. Higginson for the histological preparations.

REFERENCE

Hawking, F. (1943). *British Medical Journal*, 2, 198.

STERILIZATION OF THE SKIN BY COLOURLESS FLAVINE (5-AMINO-ACRIDINE)

BY

VICTOR BONNEY, M.S., M.D., B.Sc., F.R.C.S., F.R.A.C.S.

AND

H. SANDEMAN ALLEN, O.B.E., M.B., B.Ch., F.R.C.S.

In the *British Medical Journal* of May 18, 1918 (p. 562), one of us (V.B.), in conjunction with Carl Browning, introduced a new antiseptic solution which we named violet-green (sometimes known as "Bonney's blue"), for sterilizing the skin; this, because it was entirely non-irritant, could be applied in strength for many hours without harming the skin. The bacteriological investigations then undertaken showed that, to sterilize the skin completely, the time over which the antiseptic can be applied is even more important than its absolute bactericidal power; or, in other words, more effect is obtained from a bactericide of moderate potency which can be applied for a long time than from one whose potency is of a very high order but which, on account of its effect on the skin, can be applied for only a very short time. Violet-green, however, combined high bactericidal power with non-irritancy and therefore amply filled the bill, and since its introduction it has been extensively used not only in surgery and obstetrics but in dermatology as well. It has, however, been objected against it that it intensely stains the bed-linen of the patient and the hands of the nurses. This is true, particularly when one is unaccustomed to its use. On the other hand, applied to the vagina, the staining is a great advantage when—as, for instance, in total hysterectomy—that canal has to be opened from above, for the deep coloration defines it very clearly.

The recent production of colourless flavine* (5-amino-acridine) suggested to one of us (V.B.) the making of a solution similar to violet-green, but containing, instead of 1% of equal parts of crystal violet and brilliant green, 1% of colourless flavine, in the same solvent—namely, equal parts of rectified spirit and water. The presence of water appears to be, in general, a necessity for the full development of antiseptic action; and, in connexion with that requirement and the pale yellow powder with which this paper is concerned, an analogy may be drawn between it and another and very homely yellow powder, called mustard, which, pungent and tear-provoking when mixed with water, can yet as a dry powder be placed on the tongue without any taste or discomfort. The spirit is added to counteract the grease of the skin, which checks a purely watery solution from spreading itself and delays absorption.

It should be noted that colourless flavine is not really colourless, but has a pale lemon tint, which just stains sufficiently to show the limits of its application—a very desirable property in any antiseptic used for the skin. To make the solution, the flavine is first dissolved in the water and then the spirit is added.

Method of Application.—The skin of the operation area, having undergone the usual cleansing, is first painted with the solution and then covered with a piece of double-thickness lint,

* Made by Glaxo Laboratories, Ltd.