Medical Memoranda

Cranial Nerve Paralysis following Cerebrospinal Fever

The chief interest in the case described lies in the picking out of two different cranial nerves on opposite sides and the escape of closely adjacent nerves.

CASE RECORD

A coolie aged 20 was admitted to hospital at 5.30 p.m. on Dec. 19, 1943. He was semiconscious, extremely restless, and resistant to examination. A history of headache and fever of one day's duration was obtained from his companions. On examination the temperature was 102.6 F., there was marked nuchal rigidity, but Kernig's sign was absent. The pupils were equal and circular and reacted sluggishly to light; there was no strabismus. Heart and lungs and abdomen, *nil* abnormal. Urine, *nil* abnormal. Lumbar puncture was performed and very turbid C.S.F. under low pressure obtained. Microscopically many pus cells and Gram-negative intracellular diplococci were seen. On culture a pure growth of *N. meningitidis* was obtained.

The patient was treated with sulphapyridine, 2 g. of "dagenan soluble" being given intravenously at once, and 1 g. sulphapyridine orally every 4 hours thereafter. A total of 31 g. was administered. On Dec. 20 ptosis of the right eyelid was noticed, and on closer examination a complete third-nerve paralysis on the right was discovered. The right eye was deviated laterally and the patient could not move it upwards or nasally. Slight downward movement accompanied by a clockwise rotation of the globe was possible, showing that the fourth nerve was unaffected. The pupil was dilated and fixed and there was slight oedema of the bulbar conjunctiva.

The general condition steadily improved but the third-nerve paralysis remained. It was not until Jan. 2, 1944, that it was noticed the patient was unable to close the left eye completely and that he had difficulty in keeping saliva in his mouth. On examination a bilateral facial paralysis was found. By Jan. 8 the seventh-nerve paralysis on the right side had cleared up, but it remained on the left. There was now inability to open the right eye fully or to close the left eye. The right side of the brow could be wrinkled and the right corner of the mouth elevated. An attempt was made to determine if there was any loss of taste sensation on the left, but owing to language difficulties and lack of co-operation by the patient no definite conclusion was arrived at. Similarly it was not found possible to determine if the eighth nerve on the left side was affected. There appeared to be no sensory loss in the distribution of the fifth nerve. The third, fourth, and sixth nerves on the left were unaffected. The Kahn test was negative.

After 3 months—on March 7—the condition had almost completely cleared up; there was slight external strabismus of the right eye when at rest, but full external ocular movements were possible. There was still slight weakness of the left side of the face.

J. F. K. HUTTON, M.B., Ch.B.,

India Command.

Capt., R.A.M.C.

Diphtheria in an Infant

It is appreciated that nasal diphtheria may occur even in the newborn, and the following report of such a case includes a record of the state of the mother in relation to diphtheria.

CASE HISTORY

A male child was born on March 24, 1944, the first child of its parents. At birth it weighed 7 lb. On March 31 it developed a purulent nasal discharge, and on April 9 was admitted to hospital, a nasal swab being positive. It was a normally nourished child but had a profuse watery nasal discharge. There was no clinical sign of faucial diphtheria. Routine swabs from nose and throat on admission gave growths of the intermedius type of Klebs-Loeffer bacillus. It was given 4,000 units of diphtheria antitoxin intramuscularly. The mother denied any symptoms to suggest diphtheria, and was found to be swab-negative and strongly Schick-positive. The nasal discharge continued and swabs were persistently positive until July 12. During the interval a virulence test was twice performed. On each occasion the result was positive. Throat swabs became negative after May 15. The child's Schick test at July 10 was negative. The only other item of interest during its stay in hospital was an intimate exposure to a case of measles developing in the next cot. Protection against this infection was apparently given by the intramuscular injection of 10 c.cm. of fresh convalescent measles serum.

The patient's general condition throughout was satisfactory, and on discharge from hospital on July 24, after three successive negative swabs from nose and throat, his weight was 11 lb. 8 oz.

F. F. KANE, M.D.,

Medical Superintendent, Purdysburn
Fever Hospital, Belfast.

Reviews

ALLERGY AND ANAPHYLAXIS

Allergy, Anaphylaxis and Immuno-therapy. Basic Principles and Practice. By Bret Ratner, M.D. (Pp. 834; illustrated. \$8.50 or 47s.). Baltimore: The Williams and Wilkins Company; London: Baillière, Tindall and Cox.

Allergy is a medical topic much less talked and written about in Britain than in America, where, indeed, it seems to be a more prevalent phenomenon than it is here. It is not a popular subject with the medical student, and most practitioners have rather hazy notions about what it is or is not. This volume of over 750 closely written pages (and 56 pages of references) will lighten his darkness, for all aspects of allergy and anaphylaxis, from the experimental and theoretical to the clinical and practical, are fully described and often graphically explained. It tells about the nature of the proteins, antitoxins, and the like, which are mostly responsible for allergic reactions, the clinical manifestations and laboratory findings in serum sickness, and the methods for recognizing the hypersensitive individual and for preventing serum accidents. Transfusion risks are discussed, and the chapter on sulphonamide sensitivity is particularly valuable and topical; numerous examples are given of the accelerated type of allergic reaction to the drug, especially in patients who had shown sulphonamide sensitivity during an earlier course. The practitioner will find the section on immuno- and chemo-therapy of bacterial infections, listed alphabetically, very useful, though the expert may question some of the views and recommendations expressed there. Unfortunately there are no chapters devoted specifically to such allergic diseases as asthma, hay-fever, eczema, and food allergy. For the immunologist there are good descriptions of the mechanism and rationale of allergic and anaphylactic phenomena, including physiological pathology, blood chemistry, and cytology.

The book, which is well produced and freely illustrated, should prove useful alike to the fevers expert, the general practitioner, and the laboratory worker.

GYNAECOLOGICAL DISORDERS

The Symptomatic Diagnosis and Treatment of Gynaecological Disorders. By Margaret Moore White, M.D., F.R.C.S., M.R.C.O.G. With foreword by F. J. Browne, M.D., F.R.C.S., F.R.C.O.G. The General Practice Series. (Pp. 229; illustrated. 16s.) London: H. K. Lewis and Co. 1944.

This little book of just over 200 pages has been written for general practitioners. As the title indicates, the subject is discussed from the point of view of symptoms and symptomatic treatment. Each of the first 17 chapters opens with a tabulated summary of the contents which recalls memories of the "synopsis" type of textbook. A brief description of each condition with notes on appropriate treatment follows. The last two chapters, on contraception and radiation therapy, are by Dr. Mary Redding and Mr. I. G. Williams respectively.

In spite of the excellent detail it contains there is something unsatisfactory about this book. Criticism may be epitomized by a consideration of Chapter 17, which is devoted to pre- and post-operative treatment and complications. This is surely the province of the specialist and not the practitioner. If the latter is sharing the control of a case with the former he should be given his instructions according to the specific needs of the case and the practice of the surgeon. In other words, there must be a close relation between the practitioner and the specialist. There are things the general practitioner should take responsibility for and things he should not, but there is little in this book to guide him through the difficult no-man's-land of borderline cases. Accurate diagnosis must obviously precede intelligent treatment. The practitioner may well expect to investigate and treat pruritus vulvae or vaginal discharge, but if the tragedies of gynaecological practice are to be avoided he must know the dangers and be able to realize when the time has come to seek help. He must be warned of the dangers of leucoplakia and kraurosis, and not wait until carcinoma has developed before he refers his patient elsewhere. As for accurate diagnosis, he cannot find much help in an academic summary of possible causes. For example, in dealing with