hepatoma (Dardinski, 1932), or recent thrombosis of a splenic vein (Rhame, 1928).

It is probable that in some cases there is an arteriolar rupture. The splenic artery degenerates early. Black states that at 36 years 50% of spleens show degenerative changes. The splenic artery may show gross degenerative changes with aneurysm formation when the rest of the arteries in the body are normal (Parsons, 1937).

In the case reported above the spleen was not abnormally mobile—there was no evidence of venous obstruction, and in view of the arteriolar degeneration present it is possible that the haemorrhage was arteriolar in origin.

Summary

A further case of spontaneous rupture of an apparently normal spleen is reported and a brief review of the literature is given. It is suggested that rupture is secondary to an intrasplenic haemorrhage. It is unlikely that any one common cause of the haemorrhage will be found to explain all cases.

REFERENCES

- Bailey, Hamilton (1930). *Brit. J. Surg.*, **17**, 417.
Black, J. M. (1933). *Ibid.*, **20**, 526.
Burnett, E. C., and McMenemy, W. H. (1935). *British Medical Journal*, **1**, 1122.
Coleman, A. H. (1939). *Brit. J. Surg.*, **27**, 173.
Dardinski, V. J. (1932). *J. Amer. med. Ass.*, **99**, 831.
Dubash, J., and Langley, G. F. (1944). *British Medical Journal*, **1**, 183.
Halliwell, A. C. (1933). *Ibid.*, **1**, 919.
Parsons, C. G. (1937). *Brit. J. Surg.*, **24**, 708.
Rhame, J. S. (1928). *Ann. Surg.*, **88**, 212.
Susman, M. P. (1927). *Brit. J. Surg.*, **15**, 47.
Zuckerman, I. C., and Jacobi, M. (1937). *Arch. Surg.*, **34**, 917.

PNEUMOPERITONEUM

BY

DUNCAN LEYS, M.D., F.R.C.P.

Physician, Royal Northern Infirmary, Inverness; Consulting Physician, Highlands and Islands Medical Service

The term "pneumoperitoneum" is used to describe the presence of free gas in the peritoneal cavity without evident peritonitis. The condition is regarded as rather a curiosity, but d'Allaines (1932) was able to analyse 34 published cases, and to add one from his own experience. The fact that three cases can be reported here, all of which were observed within the space of a few months, suggests that spontaneous pneumoperitoneum is not really very uncommon, and that to be aware of the existence of the syndrome is to make it likely that one will observe it.

Case I

An unmarried woman aged 30 was admitted to the Royal Northern Infirmary under Dr. T. N. MacGregor for removal of fibroids of the uterus. Her previous health had been good except for dyspepsia for several years. She developed a post-operative collapse of the right lung with secondary infection. This was treated with sulphanilamide, but resulted in a small right-sided pleural transudate (sterile). After this had subsided she complained of a sharp pain in the right side, had a rise of temperature to 101° F., and x-ray screening (Dr. G. Gottlieb) showed gas beneath the right diaphragm (which had not been present in previous radiographs). She had some tenderness and some muscle spasm in the upper abdomen, and there was complete disappearance of liver dullness. Her general condition, however, was not that of generalized peritonitis. Further treatment with sulphadiazine was given, and after four days her temperature was normal and symptoms absent, but there was still some abdominal tenderness. This gradually disappeared, and she was discharged from hospital. She was x-rayed again three weeks later (Dr. Gottlieb), and gas was still to be seen below both domes of the diaphragm. There was an ill-defined pyloric antrum with some deformity and stiffness, apparently due to adhesions. The duodenal cap was deformed, probably from adhesions to the gall-bladder. No ulcer crater was seen. The emptying time was delayed, with a moderate two-hour residue. The appendix was partially filled, fixed to the midline, and tender on palpation.

This patient reports, 18 months later, that she has since remained in good health.

Case II

A married woman aged 60, with six children, was admitted to the Royal Northern Infirmary for attacks of abdominal pain with nausea and occasional vomiting, constipation, anorexia, loss of weight, and attacks of diarrhoea preceded by severe abdominal pain, the stools being fluid and dark, without excessive mucus. Her

general condition was good except for recent loss of weight. Her complexion was sallow. Red cells numbered 4.8 millions, with haemoglobin 89%. The heart action was normal except for extrasystoles. Stomach splash was present three hours after food. The benzidine test was consistently positive in stools. Fractional test meal: Maximum free acid, 0.15%; traces of blood throughout; starch present to the third hour. Radiograph (Dr. Gottlieb): ptosis of the stomach with no clear definition of pylorus or duodenum; a large residue of barium present at four hours; gas seen below both domes of the diaphragm.

At this time there was no tenderness, distension, or rigidity, and little discomfort. She was kept in bed, given a diet largely of milk and regular gastric lavage, and was x-rayed again three weeks later, when the appearances were much the same, though an ulcer crater could be seen at the base of the deformed duodenum. A large barium residue was seen at five hours (Dr. Gottlieb). After further gastric lavage the residue steadily diminished and the patient was symptomless. She left hospital to continue under observation as an out-patient.

Over a period of about four months she was seen at intervals of a few weeks, her only symptom being a feeling of abdominal distension "towards the evening, when there was a sensation as of something crackling." She was readmitted, however, four months after her discharge in a coma from alkalosis following several days of repeated vomiting. Tetany was present. Blood urea was 220 mg. and blood calcium 6.8 mg. per 100 c.cm. Intravenous glucose-saline rapidly improved her condition, and she later submitted to gastro-enterostomy (Mr. A. J. Hamilton), from which she made an uninterrupted recovery. Operation showed a narrow scarred pylorus, but no signs of perforation.

Case III

An unmarried woman aged 35 was referred to me in 1941 for abdominal pain of irregular occurrence, but sometimes of the hunger type and relieved by food; the pain had prevented her from working as a cook for 10 months. She had been previously studied, without a positive diagnosis being made, at the Glasgow Western Infirmary. Her general condition was good. No physical signs were found. A fractional test meal showed a maximum free acid of 0.11%, with starch present to the third hour. Radiographs (Dr. Campbell Tainsh) showed a normal oesophagus, stomach, and duodenum, with no gastric residue at six hours; the appendix was not outlined. No blood was found in test meal or stools.

A year later she was admitted as an emergency to the Lawson Memorial Hospital, Golspie, under Mr. B. S. Simpson, with abdominal pain and distension of three days' duration. She was in a state of surgical shock. General restorative measures and the passage of a flatus tube improved her condition, but the next day there was increased distension and pain. At operation, as soon as the peritoneum was opened, there was an outrush of non-odorous gas and the abdomen became deflated. No evidence of peritonitis was found, nor could any other lesion be discovered. A caecostomy was carried out. After operation she was in good condition for a few hours, but there was then a recurrence of pain and distension, with deterioration of her general condition, and the abdomen was reopened 48 hours after the first operation. Once again there was an outrush of odourless gas and no evidence of obstruction or peritonitis. The abdomen was closed, and on the following day there was once more an acute recurrence of pain and distension. A needle was inserted in the posterior axillary line beneath the ninth and tenth ribs, and there was a prompt outrush of gas, with complete relief of symptoms.

There was no further recurrence of symptoms. The patient made an uneventful recovery and the caecostomy wound was closed. I saw her at the Lawson Memorial Hospital at this time and could detect no physical signs. She was admitted to the Royal Northern Infirmary two weeks later for further study, and immediately after admission developed a pain beneath the ribs on the left side which was worse on taking a deep breath and prevented her from sleeping. The abdomen was slightly distended and tender on the left side of the epigastrium. These symptoms disappeared after a bowel movement, and subsequently there was no further pain or distension. The patient was afebrile. Radiographs (six weeks after the first symptoms) showed no abnormality in oesophagus, stomach, and duodenum, and no evidence of gas beneath the diaphragm (Dr. Campbell Tainsh).

Comment

It will be noted that these three patients were women, and I have only been able to discover a report of one male case—that of d'Allaines (1932)—and no case in childhood. All the patients described here and a great majority of those reported elsewhere have had previous symptoms referable to the gastro-intestinal tract, usually suggestive of ulceration. The gas in two analyses quoted by d'Allaines (made by Montmollin and Urban) was atmospheric except for a rather higher content

of nitrogen. The symptoms in published cases have been sometimes trivial and sometimes, as in the third case reported here, very severe. None of my patients complained of the pain beneath the left shoulder which is reported by Neumann (1943) as characteristic of artificial pneumoperitoneum (an observation confirmed by personal experience). The normal intra-abdominal pressure as measured by a manometer connected with the peritoneal cavity was -9 cm. of water in the knee-elbow position, but slightly positive in the erect position (Eisenmenger, 1935; quoted by Moberg, 1937). Moberg himself found the intra-abdominal pressure in a rabbit normally negative. My own observations in two cases of artificial pneumoperitoneum gave a normal slightly negative pressure with a very quick rise to positive pressures after introduction of quite small quantities of air at the site of the manometer needle; one could, however, continue to introduce air, without much further rise in pressure and with very little discomfort to the patient, to a total of anything up to one or more litres. No pain except slight discomfort at the shoulder was complained of by these two patients. Coombs (1922) found no immediate alteration of pressure when air or fluid was introduced into the peritoneal cavity.

Almost all the reported cases of spontaneous pneumoperitoneum have been thought to be due to escape of gas from the stomach or the first part of the intestine by a perforation so situated as to allow the escape of gas only, and healing so rapidly as not to be followed by the escape of other duodenal or gastric contents. There are at least two other possible sources of free gas in the peritoneum: (a) vaginal, via the Fallopian tubes; and (b) thoracic, via the diaphragmatic hiatus. The forcing of air through the Fallopian tubes, presumed due to vaginal douching with careless use of a syringe, has been reported by Moberg (1937), and, by the same author in another case, from supposed alteration in intra-abdominal tension secondary to gastric dilatation. This latter patient, an elderly woman with carcinoma of the pylorus, was in the habit of adopting a stooping position to provoke vomiting, and Moberg considered that she produced an unusually low pressure in the peritoneum with subsequent suction of air through the Fallopian tubes. Dr. T. N. MacGregor informs me that it is unusual to require a pressure exceeding 4 cm. Hg for artificial inflation of the Fallopian tubes, and that he considers spontaneous suction of air through the vagina very unlikely. Only one of my patients (Case III) had been using vaginal douching for leucorrhoea before the occurrence of her symptoms, and the recurrence of distension in her case makes it quite obvious that this could not have been the source of her pneumoperitoneum. It is nevertheless odd that almost all the reported cases should have been in women. An exceedingly rare cause of intraperitoneal gas is so-called "pneumatosis cystoides peritonei," which is due to infection by parasites.

Lastly, at least one case has been recorded (Hinkel, 1940) in which pneumoperitoneum was thought to be the result of an extension of gas from the thorax to the peritoneum by way of the diaphragmatic hiatus. His patient, a woman aged 70, had sudden symptoms of abdominal distension and tenderness, without severe constitutional symptoms and without fever. She was discovered to have an old left basal pleurisy with a high left dome of the diaphragm, and a bronchogram showed, just above the diaphragm and close to the heart shadow, a large emphysematous bulla which was considered to be the probable source of the gas.

The case reported by d'Allaines is the only one I can find in the literature which bears a close resemblance to Case III, reported here as having had severe symptoms without any definite evidence of perforation at operation. d'Allaines suggested that the explanation in his case lay in a perforation of the stomach so minute and so close to the oesophageal opening as to be inaccessible to observation. He recalled one patient of his, with a perforation on the anterior wall of the stomach just beneath the diaphragm, who gave a history suggesting an escape of gas preceding the appearance of the actual peritonitis by several days. He recalls, in support of his explanation, the fact that many surgeons have observed perforations to heal satisfactorily after laparotomy in which no procedure was undertaken except that of drainage of the peritoneal cavity.

Summary

Three cases of free gas in the peritoneum without evident peritonitis (pneumoperitoneum) are reported, all presumed due to minute perforations of stomach or duodenum.

REFERENCES

- Coombs, H. C. (1922). *Amer. J. Physiol.*, **61**, 159.
Hinkel, C. L. (1940). *Amer. J. Roentgen.*, **43**, 377.
Moberg, G. (1937). *Acta radiol., Stockh.*, **18**, 798.
d'Allaines, F. (1932). *Bull. mem. Soc. nat. Chir.*, **58**, 940.
Neumann, H. (1943). *British Medical Journal*, **1**, 9.

Medical Memoranda

An Unusual Case of Meningococcal Meningitis

The occurrence of two attacks of meningococcal meningitis in the same patient within a short period seems worthy of record. The case is one of a series of 30 severe cases of this disease in which there were four deaths (two moribund on admission).

FIRST ATTACK

The patient, a pagan private aged 25, was of good muscular physique. On admission he was unconscious, with marked signs of cerebral irritation. The general condition showed toxæmia, with temperature 104° . C.N.S.: neck very rigid, incontinence, Kernig's sign positive, knee- and ankle-jerks increased, with bilateral plantar extensor reflex, abdominal reflexes absent. Lumbar puncture yielded turbid fluid under pressure; globulin was increased; centrifuged deposit contained pus cells and scanty meningococci. Two grammes of soluble sulphapyridine were given intravenously, and the same amount four hours later; then 1 g. of sulphapyridine intramuscularly four-hourly. (Total course of 35 g.)

Two days after admission the breathing became distressed, with moist rales in both lungs and patchy consolidation. Recovery took place from this hypostatic pneumonia. Five days after admission the patient suddenly became unconscious and had repeated convulsions. Spasmodic twitching began in right facial muscles, with conjugate deviation of eyes to left. The spasm spread to the muscles of the upper and lower limbs on the right side, finishing with conjugate deviation of eyes to right and spasms of the left side of the body. Convulsions ceased after one hour, and further recovery was uneventful. Lumbar puncture gave clear fluid; no increase in globulin, cells 7 c.mm. in centrifuged deposit. No micro-organisms seen. The patient was discharged fit on March 1 (21 days after admission). He had no complaints and was ambulant during the last 12 days. He returned to full duty.

SECOND ATTACK

The patient was readmitted on March 27 complaining of headache and pain in the back of the neck for 24 hours. Condition on admission: temperature 103° ; neck rigidity marked; Kernig's sign positive; knee, ankle, and abdominal reflexes absent; plantar reflexes not obtained. Lumbar puncture yielded cloudy fluid under pressure; marked increase of globulin; plus cells and meningococci present in centrifuged deposit. Soluble sulphapyridine, 2g., was given intravenously, and sulphapyridine, 1 g. intramuscularly, every four hours (total course of 53 g.).

Sixteen days after admission patient was afebrile and ambulant. On the 17th day he complained of weakness on the right side of body. The temperature was 100.8° . There was right facial palsy with general loss of tone and power of the muscles of right upper and lower limb. Pupils reacted equally to light and accommodation and were equal in size. Movements of the tongue were unaffected. Hearing and vision were normal. Reflexes were generally sluggish except for increased abdominal reflexes on the left side.

His general condition deteriorated, with increasing right-sided weakness and increasing neck rigidity and a positive Kernig's sign. White cell count showed a leucocytosis of 17,000 with 70% polymorphonuclears. Lumbar puncture: turbid fluid with increase of globulin, pus cells, and scanty meningococci in deposit. He was transferred to another hospital so that further investigations could be carried out. Death took place shortly after transfer. Post-mortem examination disclosed a superficial cerebral abscess over the left temporal lobe. Direct smear showed scanty meningococci.

I am indebted to Major P. E. Rees-Davies, officer commanding, for permission to publish this case.

L. R. L. EDWARDS, M.R.C.S., L.R.C.P.

C.C.S., West African Force.

Lieut., R.A.M.C.