angle. It is divided between ligatures, and the slit in the deep fascia is closed. The skin in these conditions heals poorly; it is often eczematous, or there may be adjoining ulceration. One stitch and a firm pressure dressing only are required, and patients stop in bed until healed, getting up morning and evening for lengthening periods. The results are excellent.

The communicating veins in the lower third of the thigh seldom call for direct attack, and are settled either by incision, ligature, and injection of the groin or by a similar procedure to the internal saphenous vein, 3 to 4 in. above the knee on its antero-internal surface.

Subsequent Injections.—In the majority of cases no further treatment is required, but occasionally small areas of veins remain patent, and these respond to immediate injection by ethamolin or quinine and urethane, 2 to 5 c.cm. of each. Sodium chloride 30% is not used for ordinary injection because gross sloughing follows if a few drops leak outside the vein.

Conclusion

The radical operation for varicose veins depends on the pathology and diagnosis of the varicosities. There are two types of varicosities: those due to loss of elasticity of the internal and external saphenous venous system with incompetence of the valve guarding their terminations, putting the superficial veins in direct communication with the right side of the heart, and those due to enlargement of the communicating veins between the deep and superficial venous systems—these are mainly at the lower third of the thigh and leg.

There are seven main diagnoses made up of combinations of these.

Up to five small but essential operations are required to cure the varicose veins in a limb.

REFERENCES

- Dodd, H., and Oldham, J. B. (1940). *Lancet*, 2, 8.
Russell, H. S. (1941). *British Medical Journal*, 1, 626.
Wright, A. Dickson (1940). *Ibid.*, 1, 665, 699.

SULPHONAMIDE ALLERGY THE PERSISTENCE OF DESENSITIZATION

BY

R. G. PARK, M.B., M.R.C.P.

Major, N.Z. Medical Corps

(From a Military Hospital in the Middle East)

There are two types of allergic sensitivity which can arise during sulphonamide therapy—according to whether the drug is taken internally or is applied to the skin. The internal type may take many different clinical forms, of which fever, morbilliform erythema, and leucocytosis are the commonest features. In the cutaneous type the reaction is an eczematous dermatitis. In either case violent reactions are apt to follow administration by mouth, and one question which comes up is, Can these people be desensitized as a safeguard for the future?

The answer is that they can. Schlesinger and Mitchell (1938) reported that cases of sulphanilamide fever and rash cleared up in three or four days even though the drug was continued, and that one such case was followed up two months later and found to be insensitive. In 1942 Erskine stated that allergic cases could be desensitized either by keeping on with full doses at the time of the reaction or by giving ascending doses at a later date. In a previous report (Park, 1944) on a patient who developed sulphapyridine neutropenia and was still sensitive after three months, details of a successful desensitization were given.

In the eczematous type desensitization was introduced by Tate in 1943 (personal communication), either at the expense of a severe reaction by full doses over about four days or by smaller doses for about three weeks, causing milder reactions.

Methods Used

My experience over the last 18 months has been that desensitization has been consistently successful in these allergic

states. The drug has been given in all cases by mouth, in four or five divided doses per day—enough to keep up a fairly constant level in the blood stream. Beginning with an amount too small to cause a reaction, the dose is doubled each day until a mild reaction appears. It is then kept the same until all symptoms have subsided, when it is again increased.

The two forms of allergy differ sharply in their response to desensitization. In the internal type, in which the first sign of a reaction is a rise in temperature, it is usually possible to start with 0.1 g. at each dose, doubling up daily to reach 1 g. in four or five days. More sensitive cases, such as the one reported below, take longer; the most sensitive one yet encountered had to start with 0.005 g., and took nine weeks to work up to 1 g. In the eczemas this high degree of sensitivity is the rule. The average case takes about six weeks to desensitize if the dosage is kept below reaction level.

The next question which springs to the mind is, How long does this benefit last? Does the allergic state return in the way that it does each season after hay-fever desensitization, or is it permanent? This question has not been answered up till now, and it is only from its answer that we can learn whether this desensitization to sulphonamides is really worth while. The following case is reported as a single pointer to the possible solution of this problem.

Case Report

Sapper X., aged 40, began treatment for chancroid with sulphanilamide on July 21, 1943, taking 1 g. four times a day. On the sixth day his temperature rose to 103° F. and he developed a general morbilliform erythema, showing no tendency to photosensitivity. It was slightly irritable. He complained of pains in the knees and other joints, but there were no objective findings. The drug was stopped at once. He gave a history of having received sulphapyridine for pneumonia six months previously without any reaction. On successive days his temperature reached peaks of 103°, 101.6°, and 100° F. A leucocyte count showed a total of 10,000. From July 30 onwards he was afebrile. On Aug. 8 he was put on divided doses of 1 g. of sulphapyridine. A few hours after the first dose he developed a temperature of 101° F. and further joint pains, but no rash. The fever lasted 48 hours.

Desensitization with sulphanilamide was started on Aug. 8 with four doses per day of 0.125 g. After two doses, however, he again reacted with a fever up to 102° F., a recurrence of the rash, and further joint pains. The drug was stopped and his symptoms subsided. Sulphanilamide was continued as follows:

Date	Doses per Day	Amount (g.)	Result
Aug. 11	4	0.02	No reaction
12	4	0.04	" "
13	4	0.08	" "
14	4	0.125	Fever of 99.6°
15	3	0.125	" 101.6°
16	Nil		Still febrile
17	"		Almost afebrile
18	"		Afebrile
19	"		" "
20-22	4	0.02	No reaction
23-24	4	0.04	" "
25-26	4	0.08	Mild reaction
27-28	4	0.04	No reaction
29-30	4	0.06	" "
Aug. 31-Sept. 2	4	0.08	" "
Sept. 3-5	4	0.125	" "
6-11	4	0.15	" "
12-14	4	0.19	" "
15-16	4	0.4	" "
17	4	0.5	" "
18	4	0.6	" "
19	4	0.75	" "
20	4	0.9	" "
21	4	1.0	" "

He was discharged from hospital and nothing more was seen of him until Sept. 1, 1944, when he reported at the V.D. Dept. to have a routine injection for syphilis, which he had contracted in the meantime. He gave the story that in June, 1944, he developed a sore on his penis. At this time he was given sulphanilamide in seven doses of 1 g. over two days. This caused a febrile reaction accompanied by a recurrence of his previous symptoms, subsiding in two or three days. A diagnosis of syphilis was then made, and from that time he had been receiving weekly N.A.B. and bismuth.

He was brought in for further investigation and given the following drugs:

Sept. 3,	sulphathiazole,	1 dose of 1 g.
" 4,	"	4 doses of 1 g.
" 6,	sulphapyridine,	1 dose of 1 g.
" 7,	"	4 doses of 1 g.

No reactions to the above were seen. On Sept. 9 a dose of 1 g. of sulphanilamide was given, and caused a pyrexia up to 100° F., with the same pains in the joints. This was confirmed by a similar reaction the next day to a dose of 2 g. On Sept. 12, 13, and 14 single doses of 3 g. each of sulphapyridine, sulphathiazole, and sulphanilic acid respectively were given, with no reactions. At this stage he had to leave hospital urgently for other reasons, and it was not possible to give him any further desensitization.

Discussion

To sum up, this patient developed a highly allergic state during sulphanilamide therapy, so that he reacted to as little as 0.25 g. with a sharp pyrexia. He was also allergic to sulphapyridine in a lesser degree, with a milder reaction to a dose of 1 g. Desensitization took six weeks, but succeeded in working him up to full doses without reaction. Nine months later he apparently had a reaction to further full doses.

Reinvestigation one year after desensitization showed mild reactions to doses of 1 g. and 2 g. of sulphanilamide, but none at all to doses up to 3 g. of other sulphonamides. His allergy had therefore partially returned, but only to a fraction of its original level.

If we may be allowed to theorize on the possible immunological mechanisms underlying this and other drug allergies, the following working hypothesis would fit the facts as they are at present known. In some people sulphonamides stimulate the formation of antibodies. These will then unite with future doses of the drug to cause allergic reactions. Their formation goes on after the drug has been stopped, but gradually tapers off and finally stops. Small desensitizing doses will use up these antibodies and abolish the allergic state. If desensitization is carried out before they cease to be formed, the allergy may partly recur; but if an interval is allowed to elapse before starting desensitization, it may be expected to be permanent. It now remains for this hypothesis to be tested in practice.

Summary

Desensitization to sulphonamides has been a consistent success by the method described.

An opportunity of following up a desensitized case showed that allergy had returned after one year to only a fraction of its former level.

There is theoretical reason to believe that if such cases are left for a time before being desensitized the benefit may be permanent.

I wish to thank the Director of Medical Services, 2 N.Z.E.F., for permission to publish this report.

REFERENCES

- Erskine, D. (1942). *Lancet*, 2, 568.
Park, R. G. (1944). *Ibid.*, 1, 401.
Schlesinger, E. R., and Mitchell, W. L. (1938). *Amer. J. Dis. Child.*, 56, 1256.

PENICILLIN IN CIVILIAN PRACTICE

BY

H. B. MAY, M.R.C.P.

Assistant Director, Hale Clinical Laboratories, London Hospital

Supplies of British-made penicillin have now been available for civilian use for three months, being distributed by the Ministry of Health to the Deans of Medical Schools and, through them, to their own and associated hospitals. An indication of the types of case which have received penicillin from one centre might be considered to be of general interest.

Forty million units of penicillin a month were allotted to this centre, and the administrative heads of 43 hospitals were circularized, but, largely owing to the publicity given in the Press to the fact that supplies were limited, the amount asked for in the first month was only 3 million units; in the second month it rose to 13.5 million units, and in the third month to 22 million units. No request for penicillin was refused in cases in which the infection was caused or was probably caused by a sensitive organism, and systemic treatment when needed was as a rule started before full laboratory investigations had been carried out, although in cases in which local treatment was indicated penicillin was usually withheld until a sensitive organism had been isolated. It was suggested that for systemic treatment the penicillin should be given as a continuous intramuscular drip, using 100,000 units a day dissolved in a pint of

saline and administered into the outer side of the thigh. In each case 5 days' supply was provided.

Conditions Treated

Cases receiving penicillin comprised 15 of osteomyelitis, 5 of staphylococcal pyaemia, 2 of brain abscess, 2 of sulphonamide-resistant gonorrhoea, and 1 case each of staphylococcal orbital cellulitis, staphylococcal jugular phlebitis after mastoidectomy, streptococcal puerperal septicaemia, septic abortion, actinomycosis, pneumococcal meningitis, staphylococcal meningitis, salpingitis of undetermined aetiology not reacting to sulphonamides, generalized staphylococcal pyoderma, post-operative pneumonia of doubtful aetiology not reacting to sulphonamides, staphylococcal pericarditis, suppurative pneumonitis with multiple small abscesses demonstrated in radiographs of the chest (this patient subsequently had a pneumonectomy performed with success), and a case of severe human bite which showed a mixed flora, including *Staph. pyogenes*, *F. fusiformis*, and an anaerobic streptococcus.

Local treatment was employed alone in 50 cases of staphylococcal skin infection, mainly sycosis barbae, 20 cases of staphylococcal infections of the hand, 5 cases of staphylococcal or pneumococcal conjunctivitis, 4 cases of pneumonectomy or lobectomy to prevent post-operative empyema, 4 cases of empyema—one being a case of tuberculous empyema with an associated staphylococcal infection—2 cases of staphylococcal carbuncle, and 1 case each of infected gangrene of the foot, streptococcal arthritis of the shoulder-joint, and generalized peritonitis secondary to a perforated gangrenous appendicitis.

Results

A number of these cases will no doubt be reported in detail by individual clinicians, but certain general conclusions emerge from the series as a whole. The results of penicillin treatment have confirmed the findings of earlier clinical trials: very serious staphylococcal infections, such as meningitis and pyaemia with a high colony count in the blood culture, have been cured, but two patients with multiple lung abscesses died. In very early cases of osteomyelitis it was possible to bring about apparent cure without operative intervention, but in those seen late in the disease penicillin seemed to have disappointingly little effect. It is therefore more than ever necessary for practitioners to get cases of osteomyelitis into hospital at the earliest possible moment. The cases in which the bacteriology was unknown—viz., those of salpingitis, post-operative pneumonia, and septic abortion—all did badly; as would be expected, the result of the administration of penicillin without bacteriological control is likely to prove unsatisfactory. The case of human bite improved rapidly under general and local penicillin treatment. The patient was a negro who had been bitten on the thumb by a negro, and, as Boland (1941) showed, these bites carry a high morbidity rate due to spreading cellulitis. In this case the patient was treated with injections of 20,000 units every three hours and with local penicillin cream; the inflammation remained localized, and she was discharged from hospital in 27 days. One patient complained of considerable local pain when being treated with a continuous intramuscular drip; it was noted that blood concentrations of penicillin were lower than usual, and at necropsy there was evidence of considerable muscle damage.

Systemic Treatment.—The number of patients for whom penicillin was requested for systemic use was the same in the second and the third months in which supplies were available, although in the latter, owing to increasing supplies, greater individual doses were prescribed. It is probable, therefore, that, although spectacularly successful in suitable cases, the number of patients in hospitals needing systemic penicillin is by no means large, and the total quantity which will be used even when supplies are freely purchasable will be smaller than many of us expected.

Local Application.—The general practitioner will find local penicillin treatment of great value in suitable cases of conjunctivitis and staphylococcal skin diseases. For conjunctival instillation the solution of 1,000 units per c.cm. is less painful than the application of penicillin cream, but for sycosis the lotion was less effective than the cream; this was prepared according to the formula of Clark *et al.* (1943), and results from