Published in final edited form as:

J Am Stat Assoc. 2007 ; 102: 952–967.

Longitudinal Studies With Outcome-Dependent Follow-up: Models and Bayesian Regression

Duchwan Ryu, Debajyoti Sinha, Bani Mallick, S. L. Lipsitz, and S. Lipshultz

Duchman Ryu is ???. *Debajyoti Sihna is Professor, Department of Biostatistics, Bioinformatics, and EPI, Medical University of South Carolina, Charleston, SC 29425 (E-mail: sinhad@musc.edu).* *Bani Mallick is Professor, Department of Statistics, Texas A&M University, College Station, TX 77843 (E-mail: bmallick@stat.tamu.edu).* *S. L. Lipsitz is ???.* *S. Lipshultz is ???*

Abstract

We propose Bayesian parametric and semiparametric partially linear regression methods to analyze the outcome-dependent follow-up data when the random time of a follow-up measurement of an individual depends on the history of both observed longitudinal outcomes and previous measurement times. We begin with the investigation of the simplifying assumptions of Lipsitz, Fitzmaurice, Ibrahim, Gelber, and Lipshultz, and present a new model for analyzing such data by allowing subject-specific correlations for the longitudinal response and by introducing a subject-specific latent variable to accommodate the association between the longitudinal measurements and the follow-up times. An extensive simulation study shows that our Bayesian partially linear regression method facilitates accurate estimation of the true regression line and the regression parameters. We illustrate our new methodology using data from a longitudinal observational study.

Keywords

Bayesian cubic smoothing spline; Latent variable; Partially linear model

1. INTRODUCTION

In many observational longitudinal studies, individuals are not measured at prespecified regular intervals. We consider studies where the time points of measurements are unequally spaced and the risk of having a follow-up measurement at any time can depend on the history of past clinic visits and of previous outcomes of that individual. This situation, often called *outcome-dependent follow-up*, can arise when an individual with a poor disease history and physical condition requires more care and hence more frequent visits to a doctor/hospital. For example, Lipsitz, Fitzmaurice, Ibrahim, Gelber, and Lipshultz (2002) investigated the cardiotoxic effects of doxorubicin chemotherapy for the treatment of acute lymphoblastic leukemia in childhood. Although doxorubicin has proven to be successful in curing leukemia, it can cause progressive abnormalities of the heart in long-term survivors. One longitudinal outcome variable of the study was the patient's heart-wall thickness, which was measured via echocardiogram during hospital visits for thorough medical evaluation. The time of the next evaluation was based on the physician's judgment about required urgency of the hospital visit, taking into account the child's history of cardiac disease, and the cardiac condition at the current evaluation, as well as other unrecorded factors such as overall physical condition of the child (as perceived by the attending physician). Consequently, the unequally spaced random times of medical evaluations (visits) are considered as important indicators of a patient's health progression. Covariates of interest include doxorubicin dose, gender, and age at the end of doxorubicin chemotherapy. Our interest lies in estimating the regression models for the longitudinal outcome as well as the model of hospital visits, treating both processes as important responses of the progression of the cardiac condition and the overall health status of the patient. Our new models and

associated Bayesian methods can deal with the situation when the interval between two visits might also depend on other unobserved health factors beyond what was captured directly via the recorded history of echocardiogram results. For such a study to monitor the progression of cardiotoxicity, the goals also include prediction of future outcome and understanding the relationship between the outcome process and follow-up process.

There has been a recent surge of interest in methods to analyze longitudinal data with outcome-dependent follow-up. Classical parametric regression methods for longitudinal data have been proposed by Liang and Zeger (1986), Lipsitz et al. (2002), and Chen (2003). Several authors, notably Severini and Staniswalis (1994), Hu, Wang, and Carroll (2004), Wang, Carroll, and Lin (2005), and Lin and Carroll (2006), proposed the regression methods of longitudinal data using the *partially linear model*, which consists of a parametric linear regression for the exposure variables and a nonparametric function of the time. All these existing approaches assume, without any critical assessment of the justification and consequence of the assumption, that the process of the follow-up times is ignorable, that is, modeling of the follow-up process is not required for inference about the longitudinal outcome model. In Section 2 we explore the required conditions for the validity, the practical implications, and the limitations of this assumption.

In Section 3.1 we develop a Bayesian fully parametric regression model of longitudinal response with an ignorable assumption; however, we allow a more general longitudinal correlation structure that accommodates subject-specific correlation parameters. It is well known that in nonparametric longitudinal data regression problems with correlated errors, the performances of most of the popular data-driven smoothing parameter selection methods are unsatisfactory (Opsomer, Wang, and Yang 2001). Furthermore, misspecification of the correlation structure can cause the data-driven bandwidth selection methods to perform badly. To avoid this problem for the partial linear regression analysis of longitudinal outcome under the ignorable follow-up process assumption, in Section 3.2, we develop a model based Bayesian alternative to the profile-kernel estimating equation approach of Wand et al. (2005) that uses a natural cubic smoothing spline with a partially improper Gaussian prior and unknown smoothing parameters (Berry, Carroll, and Ruppert 2002) for the nonparametric function of time. Using our Bayesian partially linear model approach, we obtain the exact uncertainty intervals for the unknown regression parameters of the exposure variable and the nonparametric function of time. Later, we also explore the advantages of this Bayesian partially linear regression via a simulation study.

One of the limitations of the previously proposed models and methods is that they do not specifically model the association between the longitudinal process and the follow-up time process. As a consequence, the inference based on these models may be highly affected by the assumed correlation structure of the longitudinal process. The estimates of the models may have unknown bias when the simplifying ignorable assumption is not true. These existing methods cannot address the inference about the important response variable of hospital visit times. In Section 2 we develop novel models that address the joint modeling of the follow-up times and longitudinal response using latent frailty (Vaupel, Manton, and Stallard 1979) variables. In Section 3.2 we then extend our Bayesian methods for partially linear longitudinal regression models to accommodate the outcome-dependent follow-up process. In spite of extensive recent literature on the partially linear models for longitudinal and repeated measure data, there has been a complete lack of any development of a method to deal with such models in the presence of informative follow-up times.

Our next contribution is to propose a model choice methodology in Section 4 to compare these vast sets of models. This is a daunting task because the collection of possible models is not nested and has different structures, as well as having different dimensions of observed data

variables. We present a novel extension of the conditional predictive ordinate based model diagnostics (Gelfand, Dey, and Chang 1992) to compare the performance of these heterogeneous sets of models.

To explore the effect of misspecification of the correlation structure, Section 5 provides an extensive simulation study to compare the performances of several model based procedures and the likelihood based procedure of Lipsitz et al. (2002) (under ignorable follow-up process assumption) in terms of the coverage probability and the width of the interval estimates of the regression parameter when the true model is the model with informative follow-up process. Finally, in Section 6 we illustrate the proposed methods by analyzing the data from the cardiotoxicity study of doxorubicin chemotherapy (Lipshultz et al. 1995).

2. MODEL

For individual $i = 1, \dots, n$, the unequally spaced random follow-up times are denoted by $\{T_{i1} < \dots < T_{im_i}\}$. Let $\{Y_i(t)\}$ be the longitudinal outcome process. We use the simplified notation $Y_{ij} = Y_i(T_{ij})$ for the response from individual i at the random follow-up time T_{ij} . The corresponding j th *observed* follow-up time and the *observed longitudinal response* are denoted, respectively, by t_{ij} and y_{ij} . The first observed follow-up time t_{i1} (the time of first hospital visit since the end of chemotherapy) is considered by cardiologists to be a key measure in predicting the future course of heart function. The subsequent follow-up times (after entry into the study) can be modeled by a counting process of follow-up times $\{N_i(t)\}$, where $N_i(t)$ equals the number of follow-up visits by time $t > 0$. We note that modeling $N_i(t)$ is equivalent to modeling the time elapsed between follow-up times $U_{ik} = T_{i,k} - T_{i,k-1}$ for $k = 2, \dots, m_i$. Let $\mathbf{T}_i = [T_{i1}, T_{i2}, \dots, T_{im_i}]'$ be an $m_i \times 1$ vector of all random follow-up times, $\mathbf{Y}_i = [Y_{i1}, \dots, Y_{im_i}]'$ be an $m_i \times 1$ vector of all outcome measurements for subject i , and let the corresponding vectors of observed values be denoted by $\mathbf{t}$ and $\mathbf{y}$. The interval U_{ik} is expected to be dependent on the history of previous follow-up times and responses at those time points ($\mathcal{U}_{ik}; \mathcal{Y}_{ik}$) = $(T_{i1}, \dots, T_{i,k-1}; Y_{i1}, \dots, Y_{i,k-1})$.

To understand the implications and consequences of the ignorable follow-up process model, it will be advantageous to think of the process $N_i(t)$ over a discrete time axis with K time points. Our use of terminology such as “ignorable follow-up,” which is reminiscent of the typical missing data terminology of Little and Rubin (2002), is intentional as well as appropriate. The following paragraph will show the deep connection between this concept of ignorable follow-up process and the idea of ignorable missing pattern mechanism espoused by Little and Rubin (2002). For convenience of presentation, we temporarily suppress the patient index i and covariates x_i . We use the notation $[W = w|Z = z]$ to denote the conditional density of W given $Z = z$ evaluated at w . For a given subject, let $\mathbf{N} = [N_1, \dots, N_K]'$ be the $K \times 1$ random vector that indicates time points where observations of the process $Y(t)$ have been recorded, where N_t is an indicator random variable that equals 1 if $Y(t)$ is observed at time t and equals 0 otherwise.

Note that $m_i = \sum_{j=1}^K N_{ij}$. Let the values of the longitudinal response $Y(t)$ at observed follow-up times and at other time points be represented as $\mathbf{Y}_o$ and $\mathbf{Y}_c$, respectively. We denote the values of $Y(t)$ at all K time points as $\mathbf{Y}_H$, which includes the combined information from $\mathbf{Y}_o$ and $\mathbf{Y}_c$. For any subject, the joint density of the observed data is given by

$$[\mathbf{N}, \mathbf{Y}_o | \Theta, \Psi] = \int_{\mathbf{Y}_c} [\mathbf{Y}_H | \Theta] \times [\mathbf{N} | \mathbf{Y}_H; \Psi] d\mathbf{Y}_c, \quad (2.1)$$

where (Θ, Ψ) are model parameters. Existing methodologies are based on the distribution of $[\mathbf{Y}_o | \Theta]$, where Θ is the vector of parameters associated with the longitudinal response $\{Y(t)\}$. For such a method to be valid, we need modeling assumptions that ensure that $[\mathbf{N}, \mathbf{Y}_o | \Theta, \Psi]$ in (2.1) is proportional to $[\mathbf{Y}_o | \Theta]$ as a function of Θ . From (2.1), we can see that the assumptions

will be that (1) $[\mathbf{Y}_H|\Theta]$ and the conditional distribution $[\mathbf{N}|\mathbf{Y}_H; \Psi]$ have no common parameter, and (2) $[\mathbf{N}|\mathbf{Y}_H; \Psi]$ is free of $\mathbf{Y}_c$. However, it is difficult to evaluate whether the validity of these two assumptions is reasonable in a clinical application without understanding their practical implications and physical meanings. In particular, Lipsitz et al. (2002) noted that

$[\mathbf{N}|\mathbf{Y}_H; \Psi] = \prod_{j=1}^K [N_j|N_1, \dots, N_{j-1}, \mathbf{Y}_H; \Psi]$ and they assumed (in effect) that

$$\begin{aligned} & \prod_{j=1}^K [N_j|N_1, \dots, N_{j-1}, \mathbf{Y}_H; \Psi] \\ &= \prod_{j=1}^K [N_j|N_1, \dots, N_{j-1}, \mathbf{Y}_{oj}; \Psi], \end{aligned} \quad (2.2)$$

where $\mathbf{Y}_{oj}$ is the observed history of $\mathbf{Y}_o$ up to the $j-1$ th time point. When there is no shared parameter between Θ and Ψ , under the assumption of (2.2), we get $[\mathbf{N}, \mathbf{Y}_o|\Theta, \Psi] \propto [\mathbf{Y}_o|\Theta]$ as a function of Θ . It is clear from (2.2) that under such an assumption, the processes $\{Y(t)\}$ and $\mathbf{T}$ are not necessarily independent. However, to obtain either the maximum likelihood estimator or the posterior distribution of the parameter vector Θ associated with the process $\{Y_i(t)\}$ using (2.1), one needs only to consider the piece of the likelihood

$$\begin{aligned} L(\Theta|\mathbf{Y}_o, \mathbf{X}) &\propto \prod_{i=1}^n \prod_{k=1}^{m_i} f_{Y_i(t_{ik})}(y_{ik}|Y_i(t_{i1}) = y_{i1}, \dots, \\ &Y_i(t_{i,k-1}) = y_{i,k-1}). \end{aligned} \quad (2.3)$$

Thus, under this class of models, the stochastic model of $\{N_i(t)\}$ can remain unspecified (ignored), because the joint density of $(T_{i1}, \dots, T_{im_i})$ does not make any direct contribution to the estimation of the parameters Θ associated with the longitudinal process $\{Y_i(t)\}$. The assumption of (2.2) is hard to verify and somewhat difficult to interpret for many practical data examples and studies. It is a sufficient condition to ensure the likelihood of (2.3) for inference about the model of $\{Y(t)\}$, but it is not a necessary condition. Other estimating equation based approaches use similar assumptions [e.g., assumption (1) of Wang et al. 2005] to ensure that the model of $\{N_i(t)\}$ can be ignored for the inference about $\{Y_i(t)\}$. Only a fully specified joint stochastic model of the two processes (presented later in this section) can demonstrate the practical implications and physical meanings of these assumptions to ensure ignorable follow-up.

In this article we first consider a typical member (called Model 0) of the ignorable follow-up class of models. Under Model 0, the stochastic model of $\{N(t)\}$ is assumed to be ignorable and we define only the longitudinal model of $\{Y_i(t)\}$, which is given by

$$Y_i(t) = \mu_i(t|\mathbf{x}_i(t)) + \varepsilon_i(t), \quad (2.4)$$

where $\mu_i(t|\mathbf{x}_i(t))$ is a regression function of the $1 \times (q+1)$ row vector $\mathbf{x}_i(t)$ that contains a constant 1 (for the intercept) and the q covariates measured at time t . The error process $\varepsilon_i(t)$ is a mean zero Gaussian process that is assumed to be independent across different individuals, but dependent within each individual, and to have common variance $\text{var}(\varepsilon_i(t)) = \sigma^2$ and covariance $\text{cov}(\varepsilon_i(r), \varepsilon_i(s)) = \sigma^2 \rho_{irs}$. For the likelihood based analysis given by Lipsitz et al. (2002), we need to assume that $\rho_{irs} \equiv \rho_{rs}$ for $i = 1, \dots, n$, where ρ_{rs} depends only on the time elapsed $|r-s|$. For example, the (autoregressive) AR(1) structure with fixed ρ (same for all subjects) assumes that $\rho_{irs} = \rho^{|r-s|}$ (FAR1 structure), whereas the distance-invariant correlation structure of the FEXH assumes that $\rho_{irs} = \rho$ for $r \neq s$. However, for a Bayesian analysis, we can also allow the subject-specific ρ_{irs} . For example, the AR(1) structure with subject-specific ρ_i (RAR1 structure) is given by $\rho_{irs} = \rho_i^{|r-s|}$, and the REXCH structure assumes the distance-

invariant correlation structure $\rho_{irs} = \rho_i$ for $r \neq s$ with subject-specific ρ_i . For the regression function $\mu_i(t|\mathbf{x}_i(t))$, we consider an additive model given by

$$\mu_i(t|\mathbf{x}_i(t)) = f(t) + \mathbf{x}_i(t)\boldsymbol{\beta}, \quad i = 1, \dots, n, \quad (2.5)$$

where $f(t)$ is a function of the follow-up time t , and $\boldsymbol{\beta}$ is a $q + 1$ dimensional column vector of regression coefficients. Note that because the intercept term is contained in $\mathbf{x}_i(t)$ and the corresponding $\boldsymbol{\beta}$, the $f(t)$ does not contain any intercept term. For the parametric regression model, $f(t)$ can be assumed to be, say, a known order polynomial function of t with unknown coefficients, for example, a quadratic $f(t) = \theta_1 t + \theta_2 t^2$. As an alternative, we can use a natural cubic smoothing spline for the nonparametric function $f(t)$ of the partially linear model for $\mu_i(t|\mathbf{x}_i(t))$. To avoid identifiability problems, we set $\sum_{ij} f(t_{ij}) = 0$ for the partially linear model.

As an alternative to Model 0, we now consider Model 1 to address the joint modeling of the follow-up process $\{N_i(\cdot)\}$ and the longitudinal process $\{Y_i(\cdot)\}$ via a shared latent variable W_i , reflecting the subject-specific random effect. Let $\mu_i(t|\mathbf{x}_i(t))$ be the regression function and let $\varepsilon_i(t)$ be the mean zero Gaussian error process similar to Model 0. Given the subject-specific W_i , the model for $\{Y_i(t)\}$ is given by

$$\begin{aligned} Y_i(t|\mathbf{x}_i(t)) &= \mu_i(t|\mathbf{x}_i(t)) + \eta \log W_i + \varepsilon_i(t), \\ \log(W_i) &\sim N(0, \kappa), \quad i = 1, \dots, n, \end{aligned} \quad (2.6)$$

where κ is a variance of the random subject-specific W_i and η is the regression coefficient of W_i to $Y_i(t)$. We assume W_i to be independent of the process $\varepsilon_i(t)$. Furthermore, we assume T_{ik} , $k = 2, \dots, m_i$, given the frailty W_i and the observed history $(\mathcal{U}_{ik}, \mathcal{Y}_{ik})$ follows a survival distribution with the conditional hazard rate

$$\begin{aligned} h_{ik}(t|\mathcal{U}_{ik}, \mathcal{Y}_{ik}, W_i) &= h_0(t)W_i \exp(\gamma Y_{i,k-1}), \\ k &= 2, \dots, m_i, \end{aligned} \quad (2.7)$$

for $t > t_{i,k-1}$, where $h_0(u)$ is a baseline hazard rate function and γ is a regression coefficient of the most recent observed outcome on the risk of next visit. In Model 1 the stochastic model of $\{N_i(t)\}$ is not ignorable and it contributes to the estimation of the model of $\{Y_i(t)\}$ through the shared frailty (Vaupel et al. 1979) W_i . The model in (2.7) for $\{N_i(t)\}$ reduces to a nonhomogeneous mixed Poisson process (NHMPP) model (Oakes 1992; Sinha 1993) when $\gamma = 0$. However, the actual dependence of the model of (2.7) on the observed history is certainly more complex when γ is not zero. Please note that $\eta = 0$ implies that the conditional density $[T_{ik} = t_{ik} | \mathcal{U}_{ik}, Y_i(t_{i1}) = y_{i1}, \dots, Y_i(t_{im_i}) = y_{im_i}]$ does not share any parameter with the joint density $[Y_i(t_{i1}) = y_{i1}, \dots, Y_i(t_{im_i}) = y_{im_i}]$. This is the sufficient condition to ensure that the distribution of $\mathbf{T}_i$ does not contribute to the estimation of parameters of $\{Y_i(t)\}$. However, $\eta = 0$ does not imply the stochastic independence of the two processes $\{N_i(t)\}$ and $\{Y_i(t)\}$ as long as $\gamma \neq 0$. For the data examples considered here, we assume $\mathbf{x}_i(t)$ influences $h_{ik}(t)$ in (2.7) only via its influence on $Y_{i,k-1}$. The model for the $\{N_i(\cdot)\}$ considered in (2.7) is similar to the lognormal frailty model for recurrent events data considered by Hougaard (2000) and Ibrahim, Chen, and Sinha (2001b), but other choices (such as the gamma) of the frailty density with unit mean and unknown variability can be considered. We recommend using a time-constant frailty instead of a multivariate frailty model (e.g., the correlated gamma frailty model of Henderson and Shimakura 2003), because of the identifiability issues related to the unspecified multivariate frailty density in this type of data structure. For details about semiparametric models, identifiability issues, and analysis of frailty models for recurrent events data, we refer to Klein and Moeschberger (1997) and Oakes (1992) for classical approaches and Ibrahim, Chen, and Sinha (2001b) for Bayesian approaches.

3. PARAMETRIC AND PARTIALLY LINEAR REGRESSIONS

We consider Bayesian analysis using Model 0 and Model 1. For unknown parameters of these models, we provide the priors and calculate posterior quantities of interest utilizing the Gibbs sampling (Geman and Geman 1984). We start with the parametric $f(t)$ in (2.5) and then proceed to the nonparametric $f(t)$ for the partially linear model of $\mu_i(t|\mathbf{x}_i(t))$. In each model of $f(t)$, we first consider inference under Model 0, and then proceed to Model 1 and comparison of two competing models.

3.1 Parametric Regression

For the function $f(t)$ in (2.5), we consider a polynomial function of known order p ,

$$f(t) = \theta_1 t + \cdots + \theta_p t^p, \quad (3.1)$$

where $\theta = (\theta_1, \dots, \theta_p)'$ is the unknown parameter vector. For subject i , let $\mathbf{X}_i$ denote the $m_i \times (q+1)$ matrix of covariates where the j th row $\mathbf{x}_{ij}$ is the $1 \times (q+1)$ row vector of covariates at j th observed follow-up time t_{ij} with a constant 1 for the intercept. Let $\mathbf{Z}_i = [g(\mathbf{t}_i), \mathbf{X}_i]$, where $g(\mathbf{t}_i)$ is a $m_i \times p$ matrix $(\mathbf{t}_i, \dots, \mathbf{t}_i^p)$. Finally, we define $\mathbf{R}_{yi} = (\mathbf{y}_i - \mathbf{Z}_i \boldsymbol{\Theta})$, where $\boldsymbol{\Theta} = (\boldsymbol{\theta}', \boldsymbol{\beta}')'$ is a $(p+q+1) \times 1$ vector of regression coefficients $\boldsymbol{\theta}$ from follow-up times and $\boldsymbol{\beta}$ from constant terms and other covariates, $i = 1, \dots, n$. Then, based on the joint density of the $\mathbf{Y}_i$'s given observed follow-up times $\mathbf{t}_i$, the likelihood in (2.3) for Model 0 is given by

$$L(\boldsymbol{\Xi}) \propto (\sigma^2)^{-N/2} \prod_{i=1}^n |\boldsymbol{\Sigma}_i|^{-1/2} \exp \left\{ -\frac{1}{2\sigma^2} \mathbf{R}_i' \boldsymbol{\Sigma}_i^{-1} \mathbf{R}_i \right\}, \quad (3.2)$$

where the $(m_i \times m_i)$ matrix $\boldsymbol{\Sigma}_i$ is the i th diagonal element of the $(\sum_i m_i \times \sum_i m_i)$ dimensional block diagonal correlation matrix $\boldsymbol{\Sigma}_e$ of $(\boldsymbol{\varepsilon}_1, \dots, \boldsymbol{\varepsilon}_n)$ with $\boldsymbol{\varepsilon}_{ij} = \boldsymbol{\varepsilon}_i(t_{ij})$. Because, under Model 0, the model of the $\{N_i(\cdot)\}$ does not contribute to the likelihood given in (2.3), we can derive the joint posterior and full conditionals of the parameters [to be used for the Markov Chain Monte Carlo (MCMC) computation] using likelihood in (3.2) and the joint prior of $(\boldsymbol{\Xi})$, where $\boldsymbol{\Xi}$ contains the parameter vector $\boldsymbol{\Theta}$, the correlation parameters (there can be either one or n of them, depending on the correlation structure assumed), and the variance parameter σ .

We assume uniform priors for $\boldsymbol{\theta}$ and the regression parameters $\boldsymbol{\beta}$ so that their posterior means are comparable to the maximum likelihood estimates. For the model with common ρ , we use the uniform prior $\rho \sim U(-1, 1)$. For the model with subject-specific ρ_i , we consider independent uniform priors $\rho_i \sim U(-1, 1)$ for $i = 1, \dots, n$. We use a $\sigma^2 \sim \text{IG}(A_s, B_s)$ prior with known A_s and B_s for the variance σ^2 of $\boldsymbol{\varepsilon}_i(t)$. We choose $A_s = 1$ and $B_s = 10$ to have a near diffuse prior.

To illustrate our methodology for Model 1, we assume a special case for (2.7) using $h_0(u) = \lambda$ for this data example. We use the same notation of $\mathbf{y}_i$, $\mathbf{t}_i$, $\mathbf{X}_i$, $\mathbf{Z}_i$, and $\boldsymbol{\Theta}$ as before. Let $\mathbf{R}_{yi}^* = (\mathbf{y}_i - \mathbf{Z}_i \boldsymbol{\Theta} - \eta \log \mathbf{W}_i \mathbf{1})$, $i = 1, \dots, n$, and stack them into $\mathbf{R}_y^*$ over all individuals. Using the same priors for regression parameters and nuisance parameters as in Model 0, we can modify the previous likelihood in (3.2) to

$$\begin{aligned} L_1(\boldsymbol{\Xi}, \xi) &\propto (\sigma^2)^{-N/2} \prod_{i=1}^n |\boldsymbol{\Sigma}_i|^{-1/2} \exp \left\{ -\frac{1}{2\sigma^2} \mathbf{R}_{yi}^{*'} \boldsymbol{\Sigma}_i^{-1} \mathbf{R}_{yi}^* \right\} \\ &\times \prod_{i=1}^n (\kappa^{1/2} W_i)^{-1} \exp \left\{ -\frac{1}{\kappa} (\log W_i)^2 \right\} \\ &\times \prod_{i=1}^n \prod_{j=2}^{m_i} \lambda W_i \exp(\gamma Y_{i,j-1} - \lambda U_{ij} W_i e^{\gamma Y_{i,j-1}}), \end{aligned} \quad (3.3)$$

where $\xi = (\lambda, \kappa, \gamma, \eta)$. We use the $\text{Ga}(A_l, B_l)$ prior for λ , the uniform prior for η , and the IG (A_0, B_0) prior for the variance κ of the subject-specific frailty W_i . We choose A_0 and B_0 in such a way that the prior distribution centers around 1 but has large enough variance. We choose $A_0 = 10^{-2}$ and $B_0 = 10^{-2}$. We assign the uniform prior for γ , and choose A_0 and B_0 similar to A_s and B_s to obtain a near diffuse prior. By utilizing the Gibbs sampling, we can sample from the joint posterior $p(\Theta, \sigma, \xi)$.

3.2 Nonparametric $f(t)$

For a nonparametric $f(t)$, we use a Bayesian natural cubic smoothing spline (NCS) to model the $f(t)$. Let $\mathbf{f}_i = [f(t_{i1}), \dots, f(t_{im_i})]'$, $i = 1, \dots, n$, be the regression function for the i th individual so that $\mathbf{f} = (\mathbf{f}_1', \dots, \mathbf{f}_n')'$. We consider a partially improper prior (Berry et al. 2002) such that

$$\mathbf{f} \sim \text{singular normal} [0, \alpha^{-1} \sigma^2 \mathbf{K}^-], \quad (3.4)$$

where α is a smoothing parameter for the NCS, $\mathbf{K}$ is $N \times N$ matrix with rank $N - 2$ satisfying $\mathbf{f}'\mathbf{K}\mathbf{f} = \int \{f''(t)\}^2 dt$, and $\mathbf{K}^-$ is a generalized inverse of $\mathbf{K}$. The matrix $\mathbf{K}$ based on the sorted $\mathbf{t}$ can be achieved by the method of Eubank (1999, p. 244). Instead of specifying a prior for α , without loss of generality, we put a conjugate $\tau \sim G(A_t, B_t)$ prior for $\tau = \alpha/\sigma^2$, and for σ^2 and ρ_i , $i = 1, \dots, n$, we assume the same priors as in the parametric regression. We followed Berry et al. (2002) to specify $A_t = 3$ and $B_t = 1,000$.

For the implementation of Gibbs sampling, the only significant changes from the parametric regression case of Model 0 are the conditional posteriors of $\mathbf{f}$ and τ . Similarly, under Model 1 with nonparametric $\mathbf{f}$, the joint posterior distribution is very similar to the joint posterior of the parametric $f(t)$ model under Model 1, except the priors of $\mathbf{f}$ and t .

4. MODEL DIAGNOSTICS

The conditional predictive ordinate (CPO) statistics introduced by Gelfand et al. (1992) is a popular and useful model assessment tool based on the marginal posterior predictive density of the response for subject i given the observed data from the rest of the subjects. Let ξ be all parameters in the model, let $\mathbf{D}$ be the data from all subjects, and let $\mathbf{D}_{(i)}$ be all data, excluding subject i . Further let $\mathcal{Y}_i$ be the observed response vector from subject i . Then the CPO statistic for the subject i is defined as

$$\begin{aligned} \text{CPO}_i &= f(\mathcal{Y}_i | \mathbf{D}_{(i)}) = \int_{\Xi} f(\mathcal{Y}_i | \Xi) \pi(\Xi | \mathbf{D}_{(i)}) d\xi \\ &= \left\{ E_{\Xi | \mathbf{D}} \left(\frac{1}{f(\mathcal{Y}_i | \Xi)} \right) \right\}^{-1}, \end{aligned} \quad (4.1)$$

where $\pi(\Xi | \mathbf{D}_{(i)})$ is the posterior distribution of Ξ given $\mathbf{D}_{(i)}$. The last expression of (4.1) (Gelfand et al. 1992) enables us to use MCMC samples from the posterior based on the full data to compute the CPO_i . For Model 0, the distribution of $\mathbf{T}_i$ is ignorable for the predictive distribution of $\mathbf{Y}_i$; hence, for given $\mathbf{t}_i$ and the observed data $\mathcal{Y}_i = \mathbf{y}_i$ from subject i , the CPO can be computed as

$$\text{CPO}_{i0} = (f(\mathbf{y}_i | \mathbf{t}_i, \mathbf{D}_{(i)})) = \left[E_{\xi | \mathbf{D}} \left\{ \frac{1}{f(\mathbf{y}_i | \mathbf{t}_i, \Xi)} \right\} \right]^{-1}, \quad (4.2)$$

where $\Xi = (\theta, \beta, \Sigma)$.

For Model 1, the CPO_i based on the predictive distribution of $(\mathbf{T}_i, \mathbf{Y}_i)$ is not comparable to the CPO_{i0} based only on the predictive distribution of $\mathbf{Y}_i$. To remedy this problem of comparing two cross-validated densities of different dimensions, we introduce the novel idea of conditional CPO (CCPO). For Model 1, the CCPO of the subject i based on $\mathbf{Y}_i | \mathbf{T}_i = \mathbf{t}_i, \mathbf{D}_{(i)}$, is given by

$$\begin{aligned} \text{CCPO}_i &= \frac{f(\mathbf{y}_i | \mathbf{t}_i, \mathbf{D}_{(i)})}{\int f(\mathbf{y}_i | \mathbf{t}_i, \mathbf{W}, \boldsymbol{\Xi}^*) f(\mathbf{W}, \boldsymbol{\Xi}^* | \mathbf{D}_{(i)}) d\mathbf{W} d\boldsymbol{\Xi}^*} \\ &= \frac{E_{\mathbf{W}_i, \boldsymbol{\Xi}^* | \mathbf{D}} (f(\mathbf{t}_i | \mathbf{W}_i, \boldsymbol{\Xi}^*))^{-1}}{E_{\mathbf{W}_i, \boldsymbol{\Xi}^* | \mathbf{D}} (f(\mathbf{y}_i, \mathbf{t}_i | \mathbf{W}_i, \boldsymbol{\Xi}^*))^{-1}}, \end{aligned} \quad (4.3)$$

where $\boldsymbol{\Xi}^* = (\boldsymbol{\Xi}, \boldsymbol{\Theta})$. Equation (4.3) shows that the CCPO can be also computed using samples from the full posterior $\pi(\mathbf{W}, \boldsymbol{\Xi}^* | \mathbf{D})$ (similar to CPO).

Let the superscripts (q) , $q = 1, \dots, Q$, be the q th MCMC samples for corresponding parameters or subject-specific random effect. Then, under Model 0, the Monte Carlo approximation of the CPO is given by

$$\widehat{\text{CPO}}_{i0} = \left[\frac{1}{Q} \sum_{q=1}^Q \left\{ \Psi \left(\mu^{(q)}, \sigma^{2(q)} \sum_i \right) \right\}^{-1} \right]^{-1}, \quad (4.4)$$

where $\Psi(\mathbf{a}, \mathbf{B})$ is a multivariate normal density with mean vector $\mathbf{a}$ and covariance matrix $\mathbf{B}$. To compute the CCPO for Model 1 using (4.3), we need to evaluate the density of $\mathbf{T}_i$ and the joint density of $(\mathbf{Y}_i, \mathbf{T}_i)$ given $(\mathbf{W}_i, \boldsymbol{\Xi}^*)$. Using results from Section 3, we obtain the density functions

$$\begin{aligned} &f(\mathbf{y}_i, \mathbf{t}_i | \mathbf{W}_i, \boldsymbol{\Xi}^*) \\ &\propto \Psi[\mu(\mathbf{t}_i) + \eta \log \mathbf{W}_i, \sigma^2 \Sigma_i] \\ &\times \prod_{j=2}^{m_i} \lambda W_i \exp(\gamma y_{i,j-1} - \lambda u_{ij} W_i \exp(\gamma y_{i,j-1})), \\ &f(\mathbf{t}_i | \mathbf{W}_i, \boldsymbol{\Xi}^*) \\ &= \frac{f(\mathbf{y}_i, \mathbf{t}_i | \mathbf{W}_i, \boldsymbol{\Xi}^*)}{f(\mathbf{y}_i | \mathbf{t}_i, \mathbf{W}_i, \boldsymbol{\Xi}^*)} \\ &\propto \prod_{j=2}^{m_i} \left\{ \lambda W_i \exp(\gamma y_{i,j-1} - \lambda u_{ij} W_i \exp(\gamma y_{i,j-1})) \right\}, \end{aligned} \quad (4.5)$$

where $u_{ij} = t_{ij} - t_{i,j-1}$. Hence, using Q MCMC samples from the joint posterior of $(\boldsymbol{\Xi}^*, \mathbf{W})$, the CCPO can be approximated by

$$\widehat{\text{CCPO}}_{i1} = \frac{1/Q \sum_{q=1}^Q (f(\mathbf{t}_i | \mathbf{W}_i^{(q)}, \boldsymbol{\Xi}^{*(q)}))^{-1}}{1/Q \sum_{q=1}^Q (f(\mathbf{y}_i, \mathbf{t}_i | \mathbf{W}_i^{(q)}, \boldsymbol{\Xi}^{*(q)}))^{-1}}. \quad (4.6)$$

For each subject, larger values of CPO or CCPO imply a better fits of the model. As a summary statistic of CPO and CCPO over all subjects, we utilize the logarithm of the pseudomarginal likelihood (LPML; Ibrahim et al. 2001b), which is defined by

$$\text{LPML} = \begin{cases} \frac{1}{n} \sum_{i=1}^n \log(\text{CPO}_i) & \text{for Model 0} \\ \frac{1}{n} \sum_{i=1}^n \log(\text{CCPO}_i) & \text{for Model 1.} \end{cases}$$

Note that the LPML based on the CCPO proposed by us is a weighted version of the LPML based on CPO.

5. SIMULATION STUDY

We have performed an extended simulation study to compare the performances of the maximum likelihood estimators (MLE) using SAS PROC MIXED [SAS Institute Inc., Cary, NC (version 9.1), as demonstrated by Lipsitz et al. 2002] and the partial linear regression methods estimators (Wang et al. 2005) with the Bayesian estimates in terms of the width and coverage probability of the interval estimates. The MLE can be obtained only under Model 0 and with the assumption of a fixed correlation structure $\rho_{irs} = \rho_{rs}$ for all subjects. We are interested in evaluating the effects of misspecifying the correlation structure ρ_{irs} and the assumption of the ignorable visits model on the interval estimates. The Bayesian methods using parametric regression $f(t)$ (BP) and nonparametric regression $f(t)$ (BNP) under both Model 0 and Model 1 are considered here. We also consider simulations from models using the four correlation structures of $Y(t)$ that are presented in Table 1.

The REXCH model can be a useful approximation when ρ_{irs} is not a simple parametric function of $|r - s|$. The independent (IND) model is used merely to demonstrate the effect of the extreme naive assumption of within-subject independent regression errors on the estimation of the regression parameter. The RAR1 and REXCH models allow subject-specific correlation structures. The interpretations of all these correlation structures are in Section 2. This extensive simulation study simulates 500 datasets of $n = 50$ sample size from Model 1 using four correlation structures: FAR1, RAR1, IND, and REXCH. Each dataset was then analyzed using 16 methods: MLE method using FAR1 and Model 0; BP [Bayesian with parametric $f(t) = \theta_1 t + \theta_2 t^2$] using RAR1 and Model 0, FAR1 and Model 0, REXCH and Model 0, RAR1 and Model 1, FAR1 and Model 1, REXCH and Model 1; Wang's methods [based on Model 0 and nonparametric $f(t)$] using RAR1, FAR1, and REXCH; BNP [Bayesian with nonparametric $f(t)$] method using RAR1 and Model 0, FAR1 and Model 0, REXCH and Model 0, RAR1 and Model 1, FAR1 and Model 1, REXCH and Model 1.

The first visit time t_{i1} was generated from $\text{Exp}(.5)$, where $\text{Exp}(a)$ is the exponential distribution with mean a^{-1} . The subsequent follow-up times were generated via the intervals between consecutive visits $\mathbf{U}_i = (U_{i2}, \dots, U_{im_i})$ with $\text{Exp}(\lambda W_i \exp\{\gamma Y_{i,k-1}\})$, where we take $\lambda = .1$, $\gamma = 1$, and the subject-specific random effects $\log(W_i)$ as iid $N(0, \kappa = 2)$. We have allowed 20 maximum visits and the visit times are all censored at time 20 (i.e., $m_{ij} \leq 20$ and $t_{ij} < 20$).

For any given follow-up time t_{ij} , responses from subject i were generated from

$$Y_i(t_{ij}) = \mu(t_{ij}|x_i) + \eta \log W_i + \varepsilon_i(t_{ij}), \quad (5.1)$$

where $(\varepsilon_{11}, \dots, \varepsilon_{nm_n})' \sim N(0, \sigma^2 \Sigma_e)$, $\sigma^2 = .1$, $\eta = -1$, and $\log(W_i) \sim N(0, \kappa = 2)$. For simulation, we used a piecewise linear $f(t)$ with a discontinuity at $t = 10$ so that we can demonstrate the effect of wrongly specifying the parametric form $f(t)$ [we consider quadratic $f(t) = \theta_1 t + \theta_2 t^2$ to be the wrongly specified parametric $f(t)$]. The subject-specific ρ_i were generated from $U(.2, .8)$ and the fixed ρ were taken as equal to .8.

In all the examples (simulation or real), we used the same prior specifications as described in Sections 3.1 and 3.2. We found that the results are insensitive to moderate modifications of these priors. The flexibility of these priors is demonstrated by their success capturing the different types of regression functions used in the simulation. We ran a MCMC chain for 50,000 times after convergence. The convergence and mixing of the chain were checked using two independent chains and the methods described by Gelman (1996). Our final results are based on the pooled samples from these chains.

Instead of presenting all the detailed results of our vast simulation study using ten methods of estimation for eight models, we present only a summary of our findings here. The detailed simulation results can be obtained from the first author. We omit the table that shows the width of the interval estimates. Widths of the interval estimates increase significantly when we use a Bayesian partially linear model with a nonparametric $f(t)$ and a wrongly specified correlation structure. Widths of the MLE based estimates are slightly lower than the widths of the corresponding Bayes estimators under Model 0, but this may come at the price of lower coverage probability of the MLE based interval. Table 2 presents the coverage probabilities of different interval estimates when the data are generated from Model 1 using three types of correlation structure. This table shows the effect of wrongly specifying the correlation structure when we use the MLE along with Model 0 (the true model is Model 1). Even under Model 0 (when the ignorable visits assumption is correct), we have found that it is extremely important to specify the correct correlation structure while making inference using the MLE. The true coverage probability of the 95% maximum likelihood interval estimates can be as low as 78% when the true correlation structure is RAR1 (subject-specific ρ_i) while the assumed correlation structure is either IND or FAR1. Use of a Bayesian method with either a parametric [e.g., quadratic $f(t) = \theta_1 t + \theta_2 t^2$ in Table 2] or nonparametric $f(t)$ can improve this situation, mainly because it allows for a subject-specific ρ_i when it is needed. However, the coverage probability of the interval estimate for the BP method can be still around 83% for assuming the FAR1 correlation structure when the true structure is RAR1. The nonparametric method has better coverage probability even when the structure of ρ_i is misspecified, but it comes at an additional cost of an almost 20% larger interval width compared to the interval estimates using the BP method.

As shown in Table 2, correct specification of the correlation structure is necessary for good coverage probability. Among regression methods, the Bayesian partially linear model based method (BNP) has the best results in terms of coverage. Also Model 1 performed better than Model 0 under both parametric and nonparametric $f(t)$ assumptions.

The simulation studies do not show any significant gain in coverage probability and width of the interval estimate of the regression parameter from using the Bayesian nonparametric $f(t)$ instead of Bayesian parametric $f(t)$ (the gain for using Model 1 when it is correct is obvious). This is not an unexpected surprise, because it is well accepted, for the case of frequentist partially linear models, that the gain in regression parameter estimate is remarkable over the parametric linear model only when a covariate is time dependent. In our simulation study in Table 2, there is no time-dependent covariate. It is important to note that even when the underlying model uses outcome-dependent informative follow-up, the gain in coverage probability for using the nonparametric $f(t)$ (in terms of regression parameter estimate) is moderate also in the Bayesian analysis. However, Figure 1 shows very explicitly that the parametric Bayes estimate of $f(t)$ may miss the true underlying form of $f(t)$ [particularly when the unknown $f(t)$ has a sharp change point]. On the contrary, the nonparametric Bayes model of $f(t)$ can approximate the underlying shape of $f(t)$ reasonably well [the 95% credible band covers the true $f(t)$]. This suggests that for the purpose of predicting the future course of the progression of cardiotoxicity in a patient, a nonparametric $f(t)$ (partially linear model) will be

helpful and a prudent choice. For a time-stationary covariate effect, nonparameteric modeling of $f(t)$ may not be crucial.

As pointed out by a reviewer and the editor, a more equitable comparison of the Bayesian nonparametric $f(t)$ based methods would be with the partially linear models of Wang et al. (2005). In Table 2 we also report the coverage probabilities obtained from Wang's method under different error structures. The results are very similar to those obtained by the Bayesian nonparametric method under Model 0. This is obvious because Wang's method is also based on the ignorable follow-up assumption of Model 0. However, under Model 1, the coverage probability of the BNP method using Model 1 is better than the corresponding coverage probability of Wang's estimate. This gain in coverage probability for using Model 1 and BNP $f(t)$ is not as remarkable for a fixed covariate. Later, we will show the significant gain in coverage probability of Model 1 with BNP $f(t)$ for the time-dependent random covariates.

Utilizing the Bayesian model assessment techniques discussed in Section 4, larger values of the summary statistic LPML indicate a better fit of the model. For every value of γ , the parametric $f(t)$ model yields a smaller CPO statistic relative to the BNP model $f(t)$, indicating a better fit for the BNP model. For comparison of Model 0 with Model 1, the latter gives higher LPML value than Model 0 for all assumed $f(t)$ and correlation structures. In addition, Model 1 brings less variation in CPO statistics than Model 0 when we vary the assumed correlation structure. This indicates the robust behavior of the LPML under Model 1 relative to Model 0 irrespective of any assumed correlation structure.

In another follow-up simulation procedure, we showed the utility of our Model 1 under time-dependent covariates. To simulate realistic datasets for comparing different models, we have used the cardiotoxicity data as a prototype. The posterior means of the parameters from the Bayes analysis of the cardiotoxicity data (detailed analysis given in Section 6; summary of results is in Table 4) have been taken as the parameter values of the simulation using Model 1 with REXCH correlation structure. To make it more complex, we have taken a piecewise linear $f(t)$ for the simulation model and a random time-dependent binary covariate $\{x_i(t)\}$ (rather than the fixed one) generated using an independent binary process with increasing probability of $x_i(t) = 1$ as t increases. This simulation was repeated 500 times to generate 500 different datasets, which were later analyzed using Bayesian parametric and nonparametric models under Model 1 and REXCH correlation structure. We used Wang's method and the BP method as competing methods (versus the BNP method) for this simulation study. The results of the coverage probability and the width ratio of the interval estimates of the regression coefficient of the time-varying $\{x_i(t)\}$ are reported in Table 3. For the BNP method, we see significant improvement in terms of coverage probability as well as relative width of the interval estimates (all widths are compared to the width of the interval estimate using the BNP method). The performance of Wang's method is remarkably inferior to the BNP method because Wang's method is based on the wrong assumption of ignorable follow-up process. We also found that the BNP model under Model 0 performs miserably for time-varying covariates (results omitted). Figure 1 shows that even with the Bayesian parametric method using cubic $f(t) = \theta_1 + \theta_2 t^2 + \theta_3 t^3$, the shape of the true $f(t)$ can be missed. However, the posterior estimate under the BNP method can approximate the true shape of $f(t)$ to a satisfactory level.

6. ANALYSIS OF THE CARDIOTOXICITY STUDY

Doxorubicin chemotherapy has been very efficacious in treating childhood leukemia: the 5-year survival rates for patients who receive doxorubicin is at least 80%. However, doxorubicin can also be toxic to the heart and heart function becomes progressively worse over time. Patients may require frequent hospital visits for thorough and invasive medical evaluations that seriously compromise their quality of life. Measurements (via echocardiograms) of the

thickness of the heart wall were obtained for 111 patients during their hospital visits over time (Lipshultz et al. 1995). The covariates of interest are cumulative doses of doxorubicin per square meter of body surface (dichotomized as 1 = 350 mg or more; 0 = less than 350 mg), age at the end of chemotherapy (range from 1.4 to 20.1 years), and gender (1 = female; 0 = male). The times for echocardiograms (recurrent event times T) for patients were not prespecified, but depended on the physician's assessment of the child's overall health status. The hospital visit times (times of echocardiogram) ranged from .1 to 13.86 years, and there were 329 total visits from the 111 patients.

Following Lipizt et al. (2002), the MLE based analysis of the data used the regression function

$$\begin{aligned}\mu(t) = & \theta_0 + \theta_1 t + \theta_2 t^2 + \beta_1 \text{age} + \beta_2 \text{gender} \\ & + \beta_3 \text{dose} + \beta_4 (\text{gender} \times \text{dose})\end{aligned}\quad (6.1)$$

for the mean wall thickness of the heart at time t with the correlation structures FAR1 ($\rho_{irs} = \rho^{|r-s|}$) and FEXCH ($\rho_{irs} = \rho$ for $r \neq s$). We also considered additional subject-specific correlation structures RAR1 and REXCH, and both quadratic and nonparametric $f(t)$ under Model 0 and Model 1 for the Bayesian analyses. For Model 1, we used a special case of $h_0(t) = \lambda$ (constant) in (2.7).

Table 4 shows the 95% interval estimates of the regression coefficients. Although, in most cases, the interval estimate of the dose–effect on heart-wall thickness does not vary much across different models, we found that there is an approximate 10–20% reduction in the width of the interval estimate using Model 1 under the REXCH correlation structure (a 10% reduction in the width amounts to an approximately 20% increase in the sample size). It is, however, clear that under any model, there is significant evidence of the effect of dose on heart-wall thickness. For the interaction of dose and gender, we found that the MLE interval estimates do cover 0; however, one of them is almost below 0. The corresponding Bayes estimates under Model 0 may or may not contain 0, depending on the correlation structure used. However, under Model 1, this confusion is finally settled by showing that all Bayes interval estimates contain 0, suggesting that there is not enough evidence for a dose and gender interaction effect on the heart-wall thickness. Wang's method (with REXCH correlation structure) also supported this hypothesis. Other covariate effects were found to lack any significant data evidence under any model.

Under Model 1, we found that there is significant evidence that patients with a worse record of past echocardiograms (that is, with smaller Y value in the previous hospital visit) are at higher risk for future hospital visits. This is evident from the interval estimates that γ is below 0 for all three correlation structures and under any model of $f(t)$. We also found that the interval estimates of the regression parameter η of the effect of frailty W on Y are below 0 for Model 1 (under all the correlation structures), and that the association between the risk of hospital visits and the progression of heart-wall thickness has significant support from the data. It is also evident that a patient with higher than average frailty (a weaker patient) is at higher risk of hospital visits and has a higher rate of decay in heart-wall thickness even after the adjustment for dose–effect. We have also used CPO and CCPO values to perform model selection. As shown in Table 5, it is again clear that Model 1 is superior to Model 0 and the REXCH correlation structure comes out to be the winner. In general, the value of the LPML measure is found to increase for switching from models with a fixed correlation structure (FAR1) to models with subject-specific correlation structure (RAR1 and REXCH), as well as for switching from Model 0 to Model 1 under the same correlation structure and $f(t)$. This indicates that there is ample data evidence to support a substantial amount of patient variability with regard to association between heart-wall thicknesses measured at different time points. Some patients' course of heart-wall thickness over time is more predictable than those of other

patients. It is also clear that Model 1 produces narrower credible intervals and slightly higher LPML values than Model 0 for any correlation structure, however, Model 1 with REXCH correlation structure comes out to be the winner in terms of the largest LPML. This may suggest that the correlation structure between the echocardiogram values measured at two time points r and s has a significant amount of patient to patient variability; however, the AR(1) structure is unable to model the monotone relationship of ρ_{irs} with the distance $|r - s|$ between two time points. The results do not show significant support for using the partially linear model (PLM) for $\mu_i(t|\mathbf{x}_i(t))$ with nonparametric $f(t)$ in this data example. This is obvious in Figure 2, because the estimates of the $f(t)$ by BNP and Wang's method show trend very similar to that of the estimated $f(t)$ using the BP method. All three methods indicate that the heart-wall thickness begins to plateau after about $t = 10$. The PLM estimate of Wang et al. (2005) indicates a noticeable improvement after $t = 15$ and this is not very realistic (clinically).

7. DISCUSSION

In this article we considered Bayesian estimation of regression parameters in models for longitudinal data when the follow-up times are random and dependent on the history of previous outcomes and follow-up times. We have shown that Bayesian estimation of the regression parameters can handle both a model where the distribution of follow-up times is ignorable for estimation of longitudinal response and a more general model when the follow-up process is informative about the longitudinal response. For the ignorable follow-up model, the entire joint distribution of the longitudinal responses and the covariance must be correctly specified to avoid bias. We have found that, particularly for the time-dependent covariate of the longitudinal outcome, the width and coverage probability of the Bayesian interval estimate using the partially linear joint model are much better than the interval estimate of Wang et al. (2005) using the partially linear model with ignorable follow-up assumption.

For the follow-up time process, we presented an extension of the nonhomogeneous mixed Poisson process (NHMPP) given unobservable frailty random effects. As discussed in Oakes (1992), the advantages of the NHMPP model for recurrent events data include the identifiability of the frailty distribution from observed data. For NHMPP model, the marginal risk of new (say, k th) events (after integrating out the unobservable w_i) depends on the past history of events $\mathcal{U}_{ik}$ only through the summary of total number of past events ($k - 1$) and current calendar time t , that is, $h_{ik}(t|\mathcal{U}_{ik}) = h_{ik}(t)$. An extension of the NHMPP model is the serially correlated gamma frailty model of Henderson and Shimakura (2003), where W_i is modeled as $W_i(t)$, which is time varying and piecewise-constant within a pre-specified grid. However, the joint density of the piecewise-constant $W_i(t)$ is not identifiable from observed data (implying a possible dependence of model estimates on the assumed frailty density); even for $\gamma = 0$, the $h_{ik}(t|\mathcal{U}_{ik})$ is not a function of k and t (making it hard to present a simple prediction model for T_{ik} given history). Please note that the work of Henderson and Shimakura presents the composite likelihood method to analyze only discrete panel count data (contrary to the continuous time recurrent clinic visits we are dealing with). Extension of the composite likelihood method to analyze data based on actual clinic visit times is not trivial and is a topic for future research.

There are other applications of our Model 1 where the focus will be on joint modeling of the recurrent event times (say, emergency room or clinic visits due to some intervention procedure) and longitudinal responses measured at recurrent even times, when both variables are of interest. This is particularly useful in various quality of life studies where the times of emergency room (or clinic) visits as well as the responses to quality of life questionnaires collected at clinic visits are important measures of a patient's quality of life experience over time. To analyze and model such examples, our methodologies can be directly applied with minor changes, such as including a covariate effect in the survival model of follow-up times. One important competing model in this situation incorporates the frailty nonhomogeneous

Poisson counting process model of Oakes (1992) and Sinha (1993) instead of the multivariate survival model of (2.10) for follow-up times. Developing semiparametric Bayesian methods for such models and their applications to data examples are the subjects of the authors' ongoing research projects.

In this article, for the longitudinal outcome process, we also introduced novel Bayesian implementation of the partially linear regression model and we extended this partially linear model to accommodate an informative follow-up process. We presented methods that can present joint models and analysis of longitudinal and follow-up processes. The simulation study demonstrates that it is more prudent to use a partially linear longitudinal model with informative follow-up process for reliable interval estimates of the regression parameters and mean response unless a formal model diagnostic method suggests otherwise. For model diagnostics, we introduced a novel generalization of a cross-validated Bayesian procedure, and this CCPO procedure can find applications in wide ranges of situations to compare a model with noninformative monitoring or missingness versus a model with informative missingness. However, the details about wide ranging applications of the CCPO method and methods of calibrating this measure are still under development. Other Bayesian model assessments are also available, for example, Kullback–Leibler distance or the L measures of Ibrahim et al. (2001a), which deserve to be explored for application in longitudinal data with outcome-dependent follow-up times.

Acknowledgements

The authors thank the referees, the associate editor, and the editor for valuable comments that contributed to substantial improvement of the earlier version of this article. The research of Lipsitz and Sinha's was supported by R01 CA6922 and R01 AI060373. Mallick's research was supported by R01 CA104620 and CA57030.

References

- Berry SM, Carroll RJ, Ruppert D. Bayesian Smoothing and Regression Splines for Measurement Error Problems. *Journal of the American Statistical Association* 2002;97:160–169.
- Chen HY. A Note on the Prospective Analysis of Outcome-Dependent Samples. *Journal of the Royal Statistical Society, Ser. B* 2003;65:575–584.
- Eubank, RL. *Nonparametric Regression and Spline Smoothing*. 2. New York: Springer-Verlag; 1999.
- Gelfand, AE.; Dey, DK.; Chang, H. Model Determination Using Predictive Distributions With Implementation via Sampling-Based Methods (with discussion). In: Bernardo, JM.; Berger, JO.; Dawid, AP.; Smith, AFM., editors. *Bayesian Statistics*. 4. Oxford: Oxford University Press; 1992. p. 147–167.
- Gelman, A. Inference and Monitoring Convergence. In: Gilks, DSW.; Richardson, S., editors. *Markov Chain Monte Carlo in Practice*. London: Chapman & Hall; 1996. p. 131–140.
- Geman S, Geman D. Stochastic Relaxation, Gibbs Distributions, and the Bayesian Restoration of Images. *IEEE Transactions on Pattern Analysis and Machine Intelligence* 1984;6:721–741.
- Henderson R, Shimakura S. A Serially Correlated Gamma Frailty Model for Longitudinal Count Data. *Biometrika* 2003;90:355–366.
- Hougaard, P. *Analysis of Multivariate Survival Data*. New York: Springer-Verlag; 2000.
- Hu Z, Wang N, Carroll RJ. Profile-Kernel versus Backfitting in the Partially Linear Models for Longitudinal/Clustered Data. *Biometrika* 2004;91:251–262.
- Ibrahim J, Chen M, Sinha D. Criterion Based Methods for Bayesian Model Assessment. *Statistica Sinica* 2001a;11:419–443.
- Ibrahim, J.; Chen, M.; Sinha, D. *Bayesian Survival Analysis*. New York: Springer-Verlag; 2001b.
- Klein, JP.; Moeschberger, ML. *Survival Analysis: Techniques for Censored and Truncated Data*. New York: Springer-Verlag; 1997.
- Liang K-Y, Zeger SL. Longitudinal Data Analysis Using Generalized Linear Models. *Biometrika* 1986;73:13–22.

- Lin X, Carroll RJ. Semiparametric Estimation in General Repeated Measures Problems. *Journal of the Royal Statistical Society, Ser. B* 2006;68:69–88.
- Lipshultz SE, Lipsitz SR, Mone SM, Goorin AM, Sallan SE, Sanders SP, Orav EJ, Gelber RD, Colan SD. Female Sex and Higher Drug Dose as Risk Factors for Late Cardiotoxic Effects of Doxorubicin Therapy for Childhood Cancer. *The New England Journal of Medicine* 1995;332:1738–1743. [PubMed: 7760889]
- Lipsitz SR, Fitzmaurice GM, Ibrahim JG, Gelber R, Lipshultz S. Parameter Estimation in Longitudinal Studies With Outcome-Dependent Follow-up. *Biometrics* 2002;58:621–630. [PubMed: 12229997]
- Little and Rubin (2002), ???.
- Oakes, D. Frailty Models for Multiple Event Times. In: Klein, JP.; Goel, P., editors. *Survival Analysis: State of the Art*. 4. Dordrecht: Kluwer Academic; 1992. p. 371-379.
- Opsomer J, Wang Y, Yang Y. Nonparametric Regression With Correlated Errors. *Statistical Science* 2001;16:134–153.
- Severini TA, Staniswalis JG. Quasi-Likelihood Estimation in Semiparametric Models. *Journal of the American Statistical Association* 1994;89:501–511.
- Sinha D. Semiparametric Bayesian Analysis of Multiple Event Time Data. *Journal of the American Statistical Association* 1993;88:979–983.
- Vaupel J, Manton K, Stallard E. The Impact of Heterogeneity in Individual Frailty on the Dynamic Mortality. *Demography* 1979;16:439–454. [PubMed: 510638]
- Wang N, Carroll RJ, Lin X. Efficient Semiparametric Marginal Estimation for Longitudinal/Clustered Data. *Journal of the American Statistical Association* 2005;100:147–157.

Figure 1.
Plot of the true $f(t)$ and estimated $f(t)$ for the simulation study (—, BNP; ····, true; ---, BP).

Figure 2. Plots of the estimated function fitted by BP, BNP, and Wang's method to analyze the cardiotoxicity data (—, BNP; ·····, BP; ---, Wang).

Table 1

Models with different correlation structures

Model	Correlation structure
RAR1	$\rho_{irs} = \rho_i^{ r-s }$
FAR1	$\rho_{irs} = \rho^{ r-s }$ for all i
IND	$\rho_{irs} = 0$
REXCH	$\rho_{irs} = \rho_i$

Table 2
True coverage probability of the 95% interval estimate of the regression parameter

Model of $f(t)$	Method	Model used	True correlation structure		
			RARI	FARI	REXCH
Quadratic	MLE	M0, FARI	.81	.90	.86
		M0, FEXCH	.79	.87	.88
	Bayes	M0, RARI	.92	.92	.88
		M0, FARI	.83	.92	.90
		M0, REXCH	.84	.87	.96
Nonparametric		M1, RARI	.96	.94	.96
		M1, FARI	.92	.94	.88
		M1, REXCH	.92	.92	.96
		M0, RARI	.92	.91	.90
		M0, FARI	.90	.93	.91
	Wang	M0, REXCH	.92	.92	.93
		M0, RARI	.92	.90	.91
		M0, FARI	.91	.93	.92
		M0, REXCH	.91	.90	.93
		M1, RARI	.94	.92	.90
		M1, FARI	.90	.94	.94
		M1, REXCH	.92	.94	.96

NOTE: M0 means Model 0; M1 means Model 1. Simulation results are from 500 datasets of 50 subjects; the data were simulated under Model 1.

Table 3Coverage probability and interval lengths for the regression coefficient of the time-varying $\{x_i(t)\}$

Method	Coverage probability	Width ratio
BNP	.94	1
BP	.91	85.1%
Wang	.88	81.4%

Table 4
Estimated regression parameters in cardiotoxicity data

Method	CORR	Dose	S*D	γ
Model 0				
MLE	FAR1	(-.66, -.15)	(-.65, .10)	
	FEXCH	(-.64, -.15)	(-.65, .23)	
Wang	REXCH	(-.79, -.06)	(-.77, .08)	
BP	RAR1	(-.61, -.19)	(-.63, -.01)	
	FAR1	(-.64, -.20)	(-.61, .08)	
	REXCH	(-.61, -.18)	(-.75, -.02)	
BNP	RAR1	(-.61, -.18)	(-.65, .01)	
	FAR1	(-.50, -.19)	(-.51, .08)	
	REXCH	(-.61, -.09)	(-.76, .00)	
Model 1				
BP	RAR1	(-.61, -.13)	(-.59, .03)	(-.81, -.19)
	FAR1	(-.54, -.20)	(-.43, .09)	(-.76, -.24)
	REXCH	(-.58, -.07)	(-.62, .06)	(-.84, -.12) ^a
BNP	RAR1	(-.62, -.16)	(-.61, .06)	(-.80, -.11)
	FAR1	(-.55, -.05)	(-.40, .14)	(-1.55, -.26)
	REXCH	(-.57, -.20)	(-.55, .06)	(-.87, -.21) ^b

^a Model with second best LPML.

^b Model with best LPML.

Table 5

$LPML = \frac{1}{n} \sum_i \log(CCPO_i)$ for the cardiotoxicity data

Method	Model	Correlation structure		
		FAR1	RAR1	REXCH
BP	Model 0	-8.4	-8.5	-7.2
	Model 1	-7.1	-7.0	-6.9 ^a
BNP	Model 0	-7.6	-7.5	-7.2
	Model 1	-7.1	-6.9	-6.8 ^b

^aThe second best model according to the maximum LPML measure.

^bThe best model according to the maximum LPML measure.