bertus, episcopus Lisiacensis", for an estate in Gloucester, with which, we may infer, he was invested subsequently to his installation, while the gift of the queen was made to him previously to that, unless there were two priests, "Gislebertus" by Christian name, among the royal followers, one of whom held only clerical, and the other both clerical and medical, position in the retinue.

Nigellus* was a layman, I believe, and his name I now propose as that of the first surgeon in English military history. He stands as the Baron Larrey of his day, following the fortunes of the Conqueror of England. He appears in *Domesday Book* as tenant in capite of estates in Hants, Wiltshire, Hereford, and Shropshire, with which he must have been invested in reward of his services in the capacity of *medicus*, and, it may be inferred, in the special branch surgery, which it was not lawful for ecclesiastics to practise. His first two possessions in Hants and Wilts may have been given immediately after the Conquest in the first distribution of lands; but the last two must have been of later acquisition, as the subjugation of that part of England was effected at a later date, and the Conqueror could not have conveyed to his followers any lands of which he was not the actual lord by conquest, and by disposition of their Saxon owners.

It may be inferred that, as soon as William had completed the successive subjugation of the provinces, the lands were distributed among those who were present, according to the estimate of services rendered, and that these estates in Hereford and Shropshire were the reward of Nigellus on the conquest of the Welsh Marches, and that possession was given by "seizin", or on the spot. Nigellus was possessor of other estates, which, not being of regal gift, must have been by feof of the great barons, who subdivided their large grants from the king among vassals on military tenure. These estates must have been the reward of professional services to the donors; and Nigellus, not unmindful of the church, although himself a layman, endowed the church of Monteburgh with an estate of this kind, situated in Somersetshire.

From these unquestionable data, it may be inferred that the profession was well represented and well rewarded at the conquest of England; and in them we possess the historical proof of the value placed on military services by one of the greatest generals and sovereigns in universal history, whose descendants still possess, after the lapse of eight centuries, the throne he acquired, now giving laws to the widest empire the world has known.

After the completion of the great event on which all the subsequent history of England turns, "we know", as Hallam tells us in his *History of Europe during the Middle Ages*, "that the Conqueror distributed this kingdom into about sixty thousand parcels of nearly equal value, from each of which the services of a soldier were due. He may possibly have been the inventor of this politic arrangement". The service due to the king was limited to forty days in the year, after which it was continued at the expense of the crown. As military service was thus rendered feudal and but of short duration, medical services, it may be inferred, were of the same nature, and if any were required they were provided by the great barons for their feudal contingents. Doubtless the Norman kings were attended by their selected professors of the art; but as the privy expenses of the courts of these kings have not come down to us, nothing whatever is known of them. As, however, in their next appearance in our history they continued to be of continental extraction, so it may be inferred they were so in the blank interval.

[To be continued.]

THE PATHOLOGY OF SICK-HEADACHE.

By P. W. LATHAM, M.D., F.R.C.P.,

Physician to Addenbrooke's Hospital; formerly Assistant-Physician to the Westminster Hospital, etc.

DR. ANSTIE, in his paper in the *JOURNAL* of January 18th, gives no reason for his belief "that the sympathetic phenomena in migraine are mere secondary matters," but states that the views of Du Bois Reymond and Möllendorff "are rejected upon grounds which will be found stated at length in the chapters on 'Pathology' and on 'Complications' in his book on *Neuralgia*." I have carefully read over these chapters, and unfortunately fail to see the force of his reasoning; in fact, Dr. Anstie himself must have felt it inconclusive, for he says (pp. 120-121): "Nor, indeed, should I greatly care if it were finally de-

cided that migraine and clonus should be separated from the true trigeminal neuralgia, provided the following points were well impressed on the minds of practitioners." Now Dr. Anstie says "the essential seat of every true neuralgia is the posterior root of the spinal nerve in which the pain is felt; and that the essential condition of the tissue of that nerve-root is atrophy, which is usually non-inflammatory in origin" (p. 110). In his last paper in the *Practitioner* for January (p. 31), he defines migrainous pain as "atrophic molecular irritation in the trigeminal root;" and in his work (p. 163) he claims for his theory "that the whole argument shall be taken together, for it is a case of cumulative proof; every link must be weighed and tested before the remarkable strength of the chain can be felt." Let us test the first link in this strong chain by an examination of the prodromata. One which often presents itself is a glimmering commencing on one side or other of the field of vision, and, as I pointed out in my former paper, there is during this glimmering stage in many cases, a tingling in some portion of the body, as the face, or the side of the tongue, or the arm, etc., and the tingling is on the *same side* of the body as that on which the glimmering is first seen. When the head begins to ache, the pain is felt on the side or temple *opposite* to that on which the glimmering first appeared.

The "hackneyed" Tissot describes a very interesting case of this nature. "I was consulted by an officer in the Austrian service, aged 32, whose migraine was marked by characteristic nervous symptoms. 'From the age of nine years' (such are his own words) 'I suffered about every second month, sometimes more frequently, from migraine, though I have also been quite free from an attack for a year. It began in the eyes, and without previous warning. Everything suddenly appeared cloudy, more so, however, on one side than the other, like some one who has been steadily gazing at the sun. This lasted, perhaps, ten minutes, then an arm and a leg on the same side went to sleep. Sometimes one side would be affected, sometimes the other. There was a feeling of tingling or formication, which was also experienced in the mouth and tongue, and for the time being I had difficulty in speaking. These sensations continued from ten minutes to a quarter of an hour; then the pain in the head began, but in the temple only, where it continued very severe for seven or eight hours. After vomiting, I felt relief.'" (*Œuvres de M. Tissot*, 1790, tome 13, p. 112.)

Tissot omits to mention whether or not in this case the dimness of vision commenced on the opposite side to that in which the pain in the head was developed. In all the cases, however, which have come under my own observation, this is the fact. The pain in the head is always developed on the side opposite to that portion of the field of vision which is most obscured.

Only last Saturday (January 18th) a case came under my notice at Addenbrooke's Hospital which showed this in a very satisfactory manner. The patient had charge of a stationary steam-engine; and on the previous day, "while I was occupied," he said, "in cleaning my engine, a dimness came over my right eye, and everything to the right seemed indistinct. It was just the same as if I had been looking at the sun, except that the mistiness seemed all rolling about. If I closed my left eye I could see very little indeed; and though the mistiness and movement were visible when the left eye was open, I was then able to see things much better than when it was shut. After this had continued from ten minutes to a quarter of an hour, I felt a pain in the left temple and over the left eye, which soon became very sharp." He described further the character of the pain, nausea, etc., which usually accompany sick-headache.

If, then, we have disturbance in the field of vision, together with tingling along nerves distributed to the side of the body *opposite* to where the pain in the head is subsequently developed, how is it to be accounted for?

According to the view which I have advanced, we have at this stage excitement of the sympathetic, and contraction of the blood-vessels of the side opposite to where the tingling is felt; atrophic molecular irritation of the roots, or an anæmic condition of the vessels about them, seems so far to be a satisfactory explanation. But in the next stage, with paralysis of the sympathetic, the headache is developed, and the tingling of the nerves on one side of the body ceases, and is replaced by aching and painful impressions in the opposite side of the head. Now, according to Dr. Anstie, it is this stage—the headache or migrainous pain—which depends upon atrophic molecular irritation. If his theory be correct, and migraine be what is ordinarily called a neuralgia, we shall then be forced to believe what is, I think, unique, that in this form of *neuralgia* "atrophic molecular irritation" should first linger about one set of nerves, and then transfer its affections to the roots of corresponding nerves on the opposite side. How this may be explained, I leave Dr. Anstie to show. On the other hand, it is not difficult to understand that, if the vessels of the brain on the aching side are dilated

* The Rev. R. W. Eyton, in his *Antiquities of Shropshire* (the model of county history and topography), states, vol. x, p. 1, that Nigellus was a clerk and physician. He does not state that he was a priest in orders, and he appears to base his opinion of his being "a clerk" solely on the fact that the estates conferred on him in Shropshire had been previously in possession of Spirtes, an eminent Anglo-Saxon ecclesiastic, who was exiled by Edward the Confessor. He asserts that Nigellus was physician to Count Roger de Montgomery in 1086 A.D.

and so produce increased pressure on the nerves, painful sensations will be experienced along the course of the nerves; at all events, the pain, as Möllendorff shows, is lessened by compressing the carotid, and this would hardly remove atrophic molecular irritation.

Since the appearance of my paper in the JOURNAL of January 6th, Dr. Wilks has kindly referred me to his lecture on the subject of sick-headache which appeared in the *Medical Times and Gazette* of January 2nd, 1869. In this lecture Dr. Wilks has, without, I believe, being aware of Möllendorff's views, attributed the headache to the same conditions, and almost in the same terms, as the latter had done in 1867. "The vaso-motor nerve on one side is for the time paralysed, the vessels of the head dilate, more blood is sent to it, and thus the increased heat, throbbing, and pain which the patient has to suffer until the tone of the nerve is restored."

Dr. Wilks mentions one important point which Möllendorff has omitted; namely, that the pupil is contracted. He afterwards goes on to say, "the most important question to solve is the immediate cause of the function of the sympathetic being thus in abeyance." On this point, too, Möllendorff offers no suggestion.

I will now endeavour to show, more clearly than I have done on former occasions, that the suspension of the function of this nerve is dependent upon an antecedent excitement, which excitement takes place from defective tone and controlling power in the cerebro-spinal system. I base my proof upon the following propositions.

1. In the cerebro-spinal system are included fibres possessing an inhibitory power over the sympathetic system; or, in other words, in those organs in which it is possible to operate separately on the fibres proceeding from the sympathetic and from the cerebro-spinal system, the section or paralysis of one kind of fibres produces the same results as galvanism or excitement of the other kind of fibres. This is illustrated by the following extract. "All glands are provided with sympathetic nerves, and many, if not all, possess others derived from the cerebro-spinal nervous system. The experiments just referred to show that the quantity of a secretion is differently affected by the section or irritation of these two sets of nerves. Thus, irritation of the pneumogastric nerves increases the quantity of the gastric juice, whilst irritation of the sympathetic nerves diminishes or arrests it. Again, division of the sympathetic nerves of the submaxillary gland increases the flow of saliva, but irritation of the distal cut portion of the nerve diminishes it; on the other hand, section of the cerebral nerve diminishes, whilst a similar irritation of the undivided cerebral nerve causes an increase of, the secretion. Since, in the former case, the small arteries of the gland contract and the supply of blood is diminished, whilst in the latter those vessels dilate and more blood is distributed to the gland, the diminution or augmentation of the secretion accords in either case with differences in the quantity of blood conveyed to the gland, and the influence of the nervous system in regulating the quantity of the secretion is indirectly manifested by the dilatation or contraction of the coats of the small arteries." (Marshall's *Physiology*, vol. 2, 1867, p. 349.)

2. Violent mental emotions produce, in particular organs of some individuals, the same effect as would result from stimulation of the sympathetic, or section of the cerebro-spinal nerve. "Violent emotion may suspend the salivary secretion, as is shown by the well known test often resorted to in India for the discovery of a thief among the servants of a family—that of compelling all the parties to hold a certain quantity of rice in the mouth during a few minutes—the offender being generally distinguished by the comparative dryness of his mouthful at the end of the experiment That the secretion of gastric fluid is entirely suspended by powerful mental emotion seems almost certain, from the well known influence which this has in dissipating the appetite for food, and in suspending the digestive process when in active operation." (Carpenter's *Physiology*, edited by Power, 1869, p. 811.)

3. That remedies which produce contraction of the minute arteries, and which are considered as exciters of the sympathetic ganglia, will, when vision is affected by them, cause similar phenomena to those described by Dr. Hubert Airy (*Philosophical Transactions*, 1870) under the term *Teichopsia*.

To illustrate this proposition, I shall take the action of digitalis. Whatever may be the primary action of this drug in small doses, most observers agree that in full doses it acts as a stimulant to the cardiac and other ganglia of the sympathetic system. Traube's most recent view is that it acts on the vaso-motor centre. According with this, various observers have noticed the contraction of the minute arteries under its administration. Dr. Dickinson found digitalis useful in menorrhagia from its contractile effect on these vessels. The small vessels of the frog's foot have been observed to contract by Drs. Fothergill and Malan on the application of an infusion of digitalis. Stadion,

Sanders, and others refer to one fact which has an important bearing on my present theory—the production, namely, under the action of digitalis of a glimmering before the eyes. Now, it occurred to me that, if the glimmering which sometimes precedes sick-headache be due to contraction of the cerebral vessels resulting from excited action of the sympathetic, and if digitalis excite the sympathetic and so sometimes produce disturbance of vision, the appearances described under the two conditions ought to correspond with each other. I have been at considerable pains to find descriptions of the glimmering resulting from the administration of digitalis; and in those which I have found I have been struck by their remarkable correspondence with the descriptions of the appearances preceding sick-headache which I have recorded in my paper of March 23, 1872.

Dr. T. L. Brunton (*On Digitalis*, 1868, p. 46) says:—"The derangement of sight which I noticed was of two kinds—1st, a general mistiness of objects, such as is seen before fainting; and, 2nd, a large bright spot advancing before me, which sometimes resembled a ring showing prismatic colours faintly."

Purkinje (*Rust's Magazin*, Band xx, 1825, s. 236) describes the appearances much more fully. The phenomenon presented itself, he says, "in the middle of the field of vision, as a roundish spot, faintly luminous, appearing and disappearing, and around it and concentric with it several similar light and dark waves possessing like movements were observable."

After a larger dose of digitalis these waves appeared to spread from a faintly luminous quatrefoil or cinquefoil in the centre of vision—an appearance evidently not very unlike to Dr. Hubert Airy's serrated outline in *teichopsia*. As the glimmering declined, it was observed only on one side of the field of vision in curvilinear outline (*loc. cit.*, fig. 41).

"The appearance and disappearance of internal movement in these figures is best compared to the undulations seen in a broad vessel of water, the surface of which has been put into rapid motion at one or more points" (*loc. cit.*, p. 243).

"Besides this, on closing the eyes and gently rubbing them I could produce a tremulous phosphorence."

"This can be best represented by making similar figures on a flat surface in the dark with a solution of phosphorus in oil, and then as soon as the phosphorescence begins to disappear, reproducing it by a gentle movement in the air; in this way tremulousness and increase and diminution of the light, with its disappearance and reappearance, are produced, similar to what are observed in the subjective phenomena" (*loc. cit.*, Taf. 4, fig. 38-42).

The deductions which I draw from these propositions are the following. 1. If violent mental emotion stimulate the sympathetic, it does so by withdrawing the inhibitory influence of the cerebro-spinal system, and what violent emotion produces rapidly, long continued mental effort or fatigue will accomplish more slowly, but as surely. 2. The prodromata of sick-headache are due to excitement of the sympathetic; and this is proved to be the case, by the same phenomena appearing when vision is affected, as are produced by the action of digitalis. 3. It is a physiological axiom that excitement is followed by depression; and consequently excitement of the sympathetic is the cause of the "function of the nerve being temporarily in abeyance," and the headache associated with this condition.

It may be urged that the phenomena due to the sympathetic vary very much in the attacks of sick-headache in different individuals. This only supports my argument. The phenomena produced by digitalis in different individuals vary quite as much.

The abdominal complications likewise strongly support my view. The somewhat enlarged liver, the faint tinge of jaundice, etc., which occur sometimes in adults, are precisely those conditions which would result from section or paralysis of the splanchnic nerves.

The coldness of the hands and feet, too, whilst the head is hot and throbbing, is another corroboration. The fact was first discovered by Ludwig and Lovén, that when the action of the vaso-motor centre is suspended in the part supplied by the nerve, and in those which immediately adjoin it, so that their vessels become dilated, contraction of the vessels in other parts of the body is at the same time produced. The blood-pressure is thus increased generally, and produces in the locally dilated vessels a very rapid stream of blood. (See Dr. T. L. Brunton's Lectures, *BRITISH MEDICAL JOURNAL*, 1871, vol. i, p. 583.)

As the centre of the vaso-motor centre seems to be in the medulla oblongata, I so far agree as to the locality of the malady; but, as the phenomena of migraine can be explained by a series of changes which may be verified by physiological experiment, Dr. Anstie must bring forward stronger arguments than he has hitherto done before I can believe in (what even the microscope will not disclose) "atrophic molecular irritation of the trigeminus root" being the cause of sick-headache.