taneous ecchymosis formed just below the left ham; it was about the size of a sixpence, and somewhat tender. The knee continued steadily to diminish in size during the observations.

The foregoing observations show no diminution in the amount of the urea; on the contrary, the amount is above the natural average. The observations of Grandidier (*Die Hämophilie*, Leipzig, 1855, p. 36) and of Schliemann (*De Dispositione ad Hæmorrhagernic. hæredit.*, Diss. Inaug. Würzburg, 1831, p. 19) showed some diminution of the urea, in one case being 1.8 per cent., only a little if anything below natural; in the other, .5 per cent., a considerable diminution. My own observations are confirmed by those of Flüger, who found no diminution in the urea in one of his cases. (Schmidt's *Fährb.*, Band cxvii, p. 330.)

My friend, Dr. Norman Moore of St. Bartholomew's Hospital, has drawn my attention to a passage in Heberden's *Commentaries*, which he considers contains an allusion to hæmophilia. The passage may be found at page 340 of the edition published at London in 1802, in chapter 78, *De Purpureis Maculis*, and is as follows.

"In quodam puero sic affecto, si modo digitus leviter cuti imprimeretur, continuo sanguis ex vasis vicinis effundebatur, et suggillatio, tanquam in collis, fieret.

"Simul cum his maculis vidi tumores in artubus inferioribus oriri ejusdem coloris ac reliquæ cutis, et sine dolore, nisi cum membrum moveretur. Hi tumores post decem dies subsederunt, at notæ purpureæ paucos adhuc dies superfuert.

"Puerum quinquennem hujus modi tumores et dolores pessimè habuerunt. Penis adeo intumuerat, at urina ægrè reddi posset. Ad hæc, tormina interdum erant, cum vomitu, et simul stercora lineis cruentis notabantur; urina quoque sanguine colorabatur. Dolore crura occupante, æger non poterat ingredi, et brevi post doloris accessum crura punctiunculis cruentis repleta erant. Post intermissiones paucorum dierum, tumores, et notæ cruentæ reverti solebant. Hæc mala diu puerum fatigabant. Quæ alvum leniter moverunt, videbantur aliquantulum juvare. In decocto corticis Peruviani parum, aut nihil erat auxilii."

It thus appears that, saving the traumatic hæmorrhages, this great observer had noticed all the most striking features in hæmophilia—the traumatic and spontaneous ecchymoses, the swellings of the joints, their liability to recur, and the improvement following aperients. I believe it is the first time that this interesting passage has been pointed out.

CASE OF HYDROPHOBIA.

By THOMAS PARTRIDGE, L.K.Q.C.P., M.R.C.S.E.

THE question raised by Dr. Burder, as to the cause, etc., of the disease known as hydrophobia, is my excuse for sending to the JOURNAL an imperfect report of a case that occurred in my practice when residing in Birmingham.

Mr. J. S., aged 39, druggist, of a stout plethoric habit and excitable temperament, sent for me at 11.30 P.M. on the Thursday before Good Friday, 1859. I found him in bed, greatly excited, and complaining of constrictive spasms in the anterior diaphragmatic region and a tingling sensation of the right forefinger. The pulse was quick and tremulous, the skin covered with a cold sweat; the pupils were dilated, and he had a wild despairing expression of countenance. He said, "I am going to have hydrophobia." His history was that, in June 1858, just 323 days previously, he was bitten on the forefinger of the right hand by his own terrier dog. The wound bled but little. He sucked it and applied poultices for a day or two. The wound healed slowly. He continued his usual avocations, and also read with me for the Apothecaries' Hall examination, which he passed in January. On the Tuesday preceding his death, while playing at quoits, he complained of a peculiar sensation in the right shoulder and forearm, which he attributed to rheumatism; it caused him to discontinue the game. The following day he felt very unwell, from loss of appetite, nausea, flatulence, and depression of spirits. These symptoms increased until the Thursday evening, when the sensation in the finger became worse, and then, for the first time, he suspected the cause and sent for me. On my approaching him, he threw up his hands, begged me not to go near him, and became very excited. He was unable to swallow any fluid, the attempt bringing on spasm of the glottis. I gave a quarter of a grain of morphia placed on the tongue every hour. After three doses he appeared calmer and inclined to sleep. I left about 4 A.M., requesting his wife to send for two other medical friends should he be worse. On Good Friday morning, at twelve o'clock, I visited him again, and found the symptoms greatly increased in severity. He was perfectly sensible, complaining of violent spasms, continually hawking and spitting a viscid frothy mucus, which at last he had to pull away from the

mouth with his fingers. Mr. Goodall, then house-surgeon to the General Hospital, administered chloroform by the advice of Dr. Bell Fletcher and Mr. A. Baker. Large doses of cannabis Indica were also given without relief. He sank exhausted at half-past five P.M. An inquest was held, but no *post mortem* examination was made.

The long stage of incubation, if it may be so called, and the fact of the dog having bitten several other persons without ill effects ensuing, will tend to support Dr. Burder's suggestions. I may add that his wife stated the dog appeared strange in manner several days previously, starting up and snapping, running under the chairs and tables in a peculiar way; and, with the belief common amongst the public, she poisoned the animal, thinking by so doing the person bitten would not be affected. I may also say that several of those who were with him during the attack complained of shooting pains in the arms and fingers for weeks afterwards.

REPORTS

OF

MEDICAL AND SURGICAL PRACTICE IN THE HOSPITALS OF GREAT BRITAIN.

REPORT ON THE ADMINISTRATION OF ETHER.

[Continued from p. 116.]

STAMFORD INFIRMARY.

DR. NEWMAN, surgeon to the Infirmary, writes as follows. Ether has been administered in two cases requiring operation, which have recently occurred under my care in this infirmary. I am indebted to my friend Mr. Gibbings, house-surgeon, who gave the ether, for the following notes; and in the conclusions I also concur.

In the first case, that of F., aged 33, a healthy woman, having a growth from the abdominal wall requiring removal, the ether was given on a large sponge covered with a towel. The objections to this mode of administration seem to be that the drug is diffused to a disagreeable extent through the room, and that it is wastefully employed—as much as six ounces being required in this instance.

In the second case, M., aged 66, an epithelial growth was excised from the hand. A cone of pasteboard, eight inches deep, was cut so that the wide end would fit the face fairly well, and a sponge was fixed close to the small upper opening. Two ounces of ether were sufficient to produce complete anæsthesia, and an ounce was afterwards needed to keep up the effect. This man would have probably been a bad subject for chloroform. He had well-marked arcus, largely dilated cutaneous capillaries about the face and neck, and some indications of fatty heart. The heart-sounds were normal in rhythm, but feeble and heard over a large space. He had the reputation of drinking to excess.

The following points seem worthy of notice. 1. The time required to produce insensibility to pain was not longer with the ether than with chloroform as usually administered.—2. Recovery from the anæsthetic was much more rapid in the former than in the latter instance.—3. The stage of excitement was more marked with ether.—4. The pulse became more full and steady than before; and that this was maintained throughout the administration. This was especially to be noted in the second instance.—5. In neither case was there any subsequent sickness, or other disagreeable symptom, with the exception of some slight headache in the second instance.

A simple and easily managed inhaler would be of very material advantage.

SIR PATRICK DUN'S HOSPITAL, DUBLIN.

CASE OF CONCUSSION.

(Under the care of Dr. MOORE.)

ON August 26th last, William F., aged 30, when intoxicated, fell out of a cart on the back of his head. He remained about twenty minutes unconscious, and when he came to himself, as he termed it, he was unable to move, and had to be carried home. For nine days he remained in a generally paralytic state, with loss of motion and sensation in both upper and lower extremities, so that he could not move either fingers or toes; he had slight priapism, and was unable to pass urine. After the ninth day, he felt "pricking sensations" in his feet and hands. On admission to the hospital on the first week in October last, he complained of headache and of a dull aching sensation about the nape of the neck, and as far down as the middle dorsal region, especially when he remained for any length of time in the upright position. He had regained to a great extent the power of the lower extremities, but the