labour cannot be abolished, and thus the changes engendered thereby be obviated, some good at least can be achieved by substituting some other form of labour, some other method of obtaining a livelihood, for this which has induced disease in healthy organs, and can, if continued, but aggravate the morbid processes which it has inaugurated. That under favourable circumstances these morbid processes may be retarded, is almost if not quite certain. In the case of voluntary efforts, which can be abandoned when becoming distinctly harmful, this can be really seen; in other cases where the struggle for existence necessitates a continuance of the labour, the unfortunate consequences of continued strain upon the circulatory organs are painfully apparent.

A general recognition of these facts would lead to some more equitable and fair remuneration of the men exposed to disease from mechanical strain. The rate of wages should be such that the man can, if so inclined, make provision for himself when this form of labour must be abandoned, and some easier occupation substituted in its stead, and also for his family when he is taken away, as he almost certainly will be, before he reaches what we term the allotted span of existence.

In addition to the changes in the heart and its valves, let us see what are the effects upon the blood-vessels themselves of this continued strain. While we have seen that aortic valvulitis is induced by increased arterial recoil from augmented distension, it is impossible to overlook the effect of this distension upon the walls of the arteries themselves. Under these very circumstances just mentioned, we find that atheroma is induced in the arterial coats, the same parenchymatous inflammation, in fact, which in the valves constitutes valvulitis. It has long been known that atheroma is most frequent at points of greatest strain or tension, the arch of the aorta, especially the outside of the curve, at points of junction of vessels, at points of flexion, etc.

It is also well known that the pulmonary artery and its branches become atheromatous when subjected to increased strain in mitral disease. It is also well known that the walls of a vein become atheromatous when an artery opens into it. The lining membrane of the auricle becomes firm, thickened, and rigid, behind a diseased mitral valve, when the auricular wall is subjected to great and increased distension. Under these circumstances the auricular wall is sufficiently rigid not to collapse when cut into, and its internal appearance is not unlike that of a dilated and atheromatous aorta. Arteries are also liable to atheroma in proportion as they are exposed suddenly and immediately to great distension, as in the cases of the renal and coronary arteries, both short vessels coming off at once from the great arterial trunk.

General atheroma is common in aortic regurgitation where the vessels are suddenly and extensively distended by a large mass of blood being thrown at each stroke into the arterial system; and, that too, under the augmented pressure of a highly hypertrophied ventricle, for in aortic regurgitation the ventricular cavity is enlarged as well as the walls hypertrophied. So, too, in the hammerman and in the gouty person the elastic tubes which connect the two muscular ends of the circulation are liable to become atheromatous, from the increased distension to which they are subjected. We have, then, atheromatous vessels and aortic valvulitis found together, a combination, as we know clinically, common enough.

This view of the causal association betwixt atheroma and strain is not original, and is commonly held and taught in Germany, while in this country it has been enunciated by Dr. Clifford Allbutt, as well as by that accomplished pathologist, Dr. Moxon. (On the Nature of Atheroma in the Arteries. *Guy's Hospital Reports*.) That certain constitutional conditions may favour the development of atheroma is probable; but there is every reason to agree with Dr. Moxon's conclusions, viz.—“1. That what is called atheroma of arteries is sub-inflammation of various degrees, of which the lower degrees end in fatty degeneration of the coats, along with the inflammatory products; and 2. That the determining cause of the occurrence of this change is mechanical strain.”

If, then, mechanical strain leads to atheroma and its terrible consequences, aneurism and apoplexy, if it leads to valvulitis, and to such changes as seriously affect the muscular walls in consequence, and, more remotely, to all the fearful consequences of muscular failure, no apology is necessary for bringing the subject of mechanical strain in relation to the circulatory organs before your notice.

We are all clinically well aware of the association betwixt valvular disease and atheroma, and dyscrasial conditions, as gout, syphilis, and habitual drunkenness. But though we have been conscious of the fact, little light has ever been thrown upon the relationship, and it is extremely doubtful if these conditions in themselves do more than render the subjects of them more liable to changes which are less readily induced in healthy persons. It would appear that these conditions, tending as they do to induce prematurely those tissue-changes now known as senile

degeneration, favour valvulitis from its usual provoking cause, and not from any special action belonging to themselves. In gout we have seen that the exciting cause, strain, is aggravated; and in habitual drunkenness renal disease is so common as to leave no room for surprise that aortic valvulitis is also frequent, without any favouring conditions due to the dyscrasia.

The occurrence of atheroma and aortic valvulitis in syphilis is not so apparent, though the connection betwixt syphilis and the vocal chords—parts also exposed to tension—is not without importance, and throws some light, at least, on the present subject.

The effect of mechanical strain is also interesting, in regard to the forms of heart-disease in the two sexes. Not possessing any statistics, it is necessary to fall back on general impressions—distinct general impressions, however. From my experience, the conclusion has been forced upon me, that the greater portion of heart-disease, actual organic heart-disease, in women, is rather a yielding of the heart-walls, a form of weak heart, than valvular disease, especially disease of the aortic valves. Where the latter occurs, it is usually in elderly women, with chronic renal disease, and then we cannot expect sex to afford much protection. In man there is a decided tendency to aortic disease, with its accompanying changes.

It is difficult to divest oneself of a strong suspicion that this is not so much due to mere sex, but rather to the pursuits of each sex. The occupations of man render him more liable to strain on the circulatory organs, and its consequences, valvulitis and atheroma. Probably these changes in the subjects of syphilis are much more common in the male than the female, due to their different pursuits. Females are, however, subject to aortic valvulitis, when exposed to mechanical strain. In Vienna, where much of the coarser work is done by women, Bamberger, in fifty cases of aortic regurgitation, found no fewer than twelve cases in women. This also throws some light on the effects of mechanical strain.

What determines valvulitis in the mitral valve, commonly the subject of such change, we cannot yet say. We know that the mitral valve is very liable to be affected in acute rheumatism, and, according to the Germans, in scarlatina, though to a less extent; but as yet these are ultimate facts of which no satisfactory explanation has been forthcoming. Virchow's view, that the frequency of mitral valvulitis is due to violent closure of this valve by the muscular left ventricle, seems highly probable.

Finally, we may sum up the foregoing in several propositions.

1. Changes in the right heart are induced by increased strain when the mitral valve is diseased.
2. Mitral valvulitis often results from aortic disease, in consequence of the mitral valve being forcibly closed by an hypertrophied ventricle.
3. Aortic valvulitis, as well as atheroma, is intimately associated with mechanical strain.
4. Certain dyscrasial conditions, in which these affections are common, merely favour the occurrence of such changes.
5. Women are much less subject to aortic valvulitis than men are, and this is due to their pursuits rather than to their sex.
6. The importance of mechanical strain in the production of disease in the circulatory organs is scarcely yet sufficiently appreciated.

In conclusion, no attempt is being made here to give mechanical strain any undue prominence in the causation of disease in the organs of the circulation, but merely to point out such indications of its importance as the present state of our knowledge will warrant. Nor can it be supposed that it is possible, in a brief summary like this, to bring forward evidence to substantiate each conclusion. The conclusions are only what appear to me to be indicated by my investigations and reflections. Such as they are, they are given to the profession in the hope that they may attract such an amount of attention as may substantiate them, or explain them away. Whichever happens, we may hope that the truth will be ultimately elicited. Should they be substantiated, a direct impetus will be given to the treatment, both preventive and palliative, of a large class of most important diseases.

DESCRIPTION OF A NEW DOUBLE CURRENT INHALER FOR ADMINISTERING ETHER.

By J. T. CLOVER, F.R.C.S. Eng.

THE opinions of medical men on the merits of ether and chloroform are so various, that I think I need not apologise for attempting to reconcile them. There are some surgeons who do not hesitate to express their opinion that chloroform is an uncertain agent; that it is dangerous in some cases, and in others not; and that we cannot always discover beforehand whether it will prove so. To such men, ether, which very

rarely destroys life by its immediate effect, must be welcome ; and they will speak lightly of the choking sensations and headache, and even of the vomiting, which ether excites. Others, who have found chloroform safe, and believe they can account for the fatal cases by circumstances which may be avoided, must object to an agent which is less readily applied, and more complained of by the patient and bystanders than chloroform is ; but this will not account for such differing reports on the occurrence of vomiting. One surgeon reports eleven cases of vomiting out of twelve ether cases, and another says that only one patient vomited out of a much larger number to whom he gave it. I think that the amount of the anæsthetic which entered the blood, the way it was given, and the time it was kept up, must have differed widely, and would go far to account for the different results. I believe chloroform is safe if due precautions are taken, and that ether is unsafe without them ; but inasmuch as ether destroys animal life less quickly, it does not need so much care in using it.

To insure safety in producing the complete quietude of a patient by chloroform, we ought, besides watching the effects produced, to use an apparatus which will render it impossible for a patient to have within his chest as much as four per cent. of the vapour of chloroform ; and this can only be effected by measuring both the air and the chloroform by a special apparatus. The inhalation of ether is made less irritating by the same apparatus ; but I do not now use it for ether, except in operations about the mouth, as I have succeeded in supplying the ether-vapour with sufficiently uniform dilution by the inhaler which I am about to describe. It consists of a face-piece without any valves ; a metal box measuring six inches by four, and five deep ; and an elastic tube, five-eighths of an inch in diameter, to connect them. The box is either suspended by a ribbon from the administrator's neck, or placed upon the patient's bed. Inside the box is a tube of very thin copper, which conveys the expired air through it, and is then provided with a valve which opens only during expiration. This tube is broad enough to extend across the box, and undulating, in order to present a large surface, which is covered with cloth to absorb the ether. Plates of metal are so disposed as to direct the current of air, which enters through a valve during inspiration, over the surface of the tube. The following diagram represents a perpendicular section of the inhaler.

The flexible tube leading to the face-piece joins it at B. The part of the vessel containing ether-vapour is marked E. When the patient inhales, air enters at A and follows the course marked by the arrows ; when he exhales, the current in the flexible tube is reversed, and enters the metallic tube marked R, and escapes at the valve S. The opening for supplying and removing the ether is at C.

It will be observed that the part where the greatest cold is produced, by the current of fresh air passing over the ether, corresponds with the termination of the exhaling tube, and will condense much of the ether-vapour breathed back by the patient ; also, that the current of air towards the patient goes over a surface which has been warmed by the patient's breath as it passes outwards through the tube. A sliding shutter at O regulates the admission of air, without passing through the ether-chamber ; but the egress of the patient's breath at the same opening, in expiration, is prevented by a valve. A similar valve and shutter are placed at the end of the flexible tube, near the

face-piece. Both should be open at first and closed gradually, that coughing and struggling against the pungency of the ether may be avoided.

After supplying about six or eight ounces of ether, the inhaler should be shaken, to diffuse the liquid over the cloth which covers the exhaling tube. At the conclusion of the operation, a considerable quantity of ether and some water will be found condensed inside this tube, and may be poured out at S by inclining the inhaler. The ether thus saved can be rectified and used again. As, under ordinary circumstances, it would have been diffused through the room, it is a measure of the extent to which the odour of ether about the house is prevented.

The chief advantage of this apparatus is, however, in the comparative uniformity of mixture with air which it effects, and which, whilst securing complete quietude of the patient, prevents the unpleasant results of an over-strong dose. Although this double current-inhaler could be modified so as to make it useful for other anæsthetics, it is for ether that I have now introduced it. I still recommend chloroform to be given by means of a bag containing a known proportion in every cubic inch of air.

The apparatus can be obtained of Messrs. Mayer and Meltzer, of Great Portland Street, or of Messrs. Coxeter.

ON THE ADMINISTRATION OF CHLOROFORM.*

By HENRY MARSHALL, M.D., F.R.S.Ed., F.R.C.S.Ed.,
Consulting-Surgeon to the Bristol General Hospital.

It is now nearly a quarter of a century since the late Sir James Simpson discovered the anæsthetic properties of chloroform ; and since then, up till a very recent period, it has in this country, and indeed throughout Europe, been considered so superior to any other anæsthetic, as to be practically the only one used. There has, however, of late, been an uneasy feeling current that its dangers have been underestimated, and a re-action against its use has in many quarters set in.

Believing the present panic to be quite uncalled for, I venture to bring the subject of chloroform administration once more before you. My contention is, that in careful and competent hands the danger to life is so slight as hardly more requiring to be taken into account than the risk run in the performance of some minor surgical operation, which, though usually perfectly safe, may, under unusual and untoward circumstances, at times prove fatal. It is a fact full of significance, that those who have had the largest experience in giving chloroform have had the greatest immunity from fatal results, a strong *primæ facie* argument against unavoidable death being at all common. Mr. Syme's practice gives 7,500 cases of chloroform administration without a single fatal result ; and Mr. Clover, with an equally large experience, has been alike uniformly successful. Dr. Snow computed he had given it in upwards of 4000 cases, with one death while under the influence of chloroform, but very doubtful if in any way due to it. Dr. Jones has given it in upwards of 6000 cases, with one death ; Mr. Lister in some 4000 cases, with no death. So also the experience of particular hospitals goes far to show that, where it is most extensively used, "accidents" are least frequently met with. This is notably the case in the Edinburgh Infirmary, where chloroform was first publicly administered, and since then has been given with a frequency unknown in other hospitals ; and I am glad to add the Bristol General Hospital bears similar testimony.

My own experience of chloroform dates almost from the time of its introduction, having administered it early in the year 1848. A few years subsequently, I gave it with extreme frequency for the late Mr. Syme, in his private and public practice, and then learned that perfect confidence in its safety as an anæsthetic which subsequent experience has only tended to confirm. At that period, it was no uncommon occurrence for patients to come to Edinburgh for the express purpose of having operations performed under the influence of chloroform, after being refused it in London and other parts of England, on the score of heart-disease and other disorders, which were deemed incompatible with its safe administration. Acting under Mr. Syme's direction, I gave it in these cases with the same confidence as if no disease were present ; his maxim being, that a case suitable for surgical operation was suitable for chloroform, an empty stomach being the only condition insisted on. The same principles as were then enunciated by Mr. Syme continue to be taught by his distinguished successor, Mr. Lister, and are successfully practised by innumerable pupils in all parts of the world.

The mode of administration which we adopt is extremely simple. The

* Read before the Bath and Bristol Branch.