uric acid. In some cases, which I have had special opportunities of watching closely, the outbreak of a fit of convulsion had been preceded by more than usual accumulation of phosphatic matter and of uric acid. It is interesting to remember that uric acid crystals are not seldom found in the urine and in the blood in the albuminuria of pregnancy. I am not aware of any distinct evidence in support of the conjecture that uric acid, as such, is the exciting cause of convulsion; and I am not prepared to accept the doctrine of Frerichs, that the cause is ammonia resulting from the decomposition of urea. I venture to submit—pretending to no recondite skill in humoral chemistry—that, until more precise correlative chemical and clinical investigations have been made, it is wiser to be content with the general conclusion that the poisonous convulsion-provoking element cannot be specified; but that it is to be found amongst the products that ought to be excreted through the agency of the lungs and glandular system.

In rather a large proportion of cases, including not a few in which the climacteric has not been reached, this loading of the urine with phosphates and uric acid is greatly occasioned, or increased, by the habit of resorting to stimulants. In these cases, vomiting is a frequent complication; and, in subjects not specially prone to epilepsy, constitutes the chief nervous disorder. Occasionally, albuminuria even is caused by the combination of alcoholism and phosphatic accumulation, and it may persist so long as to give ground for concluding that it depends upon permanent Bright's disease; but I have seen it vanish rapidly when alcohol was cut off, and when the glandular system was set to work, the nervous symptoms subsiding simultaneously.

In this class of cases may be easily traced the influence of the mind, of the emotions, in the evolution of convulsion; or perhaps it might be more correct in some cases to say that the poisoned blood takes effect first upon the brain, inducing disorder of the intellect, and that thus it becomes a more ready irritant of the spinal cord. In many cases of hysteria and of epilepsy the fit is preceded for several hours, for a day or more, by a strange alteration in the mind. The patient exhibits unwonted excitement, passion, suspicion; is irrepressibly loquacious, perhaps violent in action; complains of intense headache; she is, in fact, for the time beside herself. It is quite certain that perception is disordered, and the faculty of comparison suspended. The apparent untruthfulness of patients of this class is often a source of pain to those about them. In some cases, this untruthfulness is real. There is no saying more questionable than the toper's maxim, "In vino veritas". Alcohol is too often the enemy of truth; and this is never more clearly seen than in the conduct of those who have become the slaves of drink. But I refer to this subject for the purpose of offering a different explanation, which I am sure applies to many cases: the untruthfulness is apparent. There is untruth in reference to facts; but there may be no untruth if reference be made to the patient's own impressions. Under the mental perturbation of an impending fit—a state compounded of vertigo and delirium—the senses are subject to illusions, perception is distorted, and the false impressions are often crystallised as delusions, and so indelibly engraven on the memory. Something similar, I am certain, often occurs in persons whose faculties are impaired under the combined influence of nervous disease and the abuse of stimulants. I have seen persons who, on recovering from the disease and the associated alcoholism, have no longer shown any tendency to falsehood.

HYDRATE OF CHLORAL IN TETANUS.—Dr. Coryllos of Patras relates, in the *Allgemeine Wiener Med. Zeitung* for January 14th, the case of a lady aged 70, who, on August 27th, 1872, wounded the sole of her right foot with an iron nail. The injury caused her little inconvenience until September 12th, when spasms of the limb set in, accompanied with trismus. The next day, Dr. Coryllos found the patient complaining of painful tetanic spasms of the right leg, affecting chiefly the gastrocnemius, and of severe pain along the course of the crural nerve. The puncture made by the nail was visible; but there was no swelling, and scarcely any discharge from the wound. She could not open her mouth easily, and had frequent difficulty in swallowing. The cervical muscles were contracted; but those of the remainder of the body were not affected. She was ordered to take every two hours a tablespoonful of a mixture containing a drachm of chloral in two ounces of a mixture of equal parts of distilled water and orange syrup. The affected limb was rubbed three times a day with mercurial ointment and extract of belladonna. Two days later, as she remained in the same state, the dose of chloral was doubled; and the next day an eighth of a grain of acetate of morphia was given every two hours, alternately with the chloral. From this time she improved, and was well a month after being first seen. The morphia was given for four days only, four grains in all being used. The chloral was continued for fourteen days, in which time she took eighteen drachms.

CLINICAL LECTURES ON MENTAL AND CEREBRAL DISEASES.

By J. CRICHTON BROWNE, M.D., F.R.S.E.,

Medical Director, West Riding Asylum; Lecturer on Mental Diseases to the Leeds School of Medicine; etc.

IV.—CANCER OF THE BRAIN.—(Concluded.)

ARE there any symptoms, from the presence or combination of which, cancer of the brain may be inferred to exist? If you dip into the literature of the subject, you will find contradictory answers to that question. Andral, who analysed forty-three cases, observed by himself or others, says, that there are no characteristic symptoms at all, whereas Guislain, who made a careful study of organic lesions of the encephalon, maintains, that special pathognomonic signs may be detected in every case. MM. Sauze and Aubanel suggest that cerebral cancer is often confounded with general paralysis, while Dr. P. Berthier writes an able monograph, *De la Folie Cancéreuse*. Without acknowledging any such absurdity as cancerous insanity—we might as well speak of tubercular jaundice, or fibroid leucorrhœa—I am inclined to believe that, with every heterologous growth in the brain, we have certain definite symptoms, in a certain definite succession. I admit that, in a certain number of cases, these symptoms are not sufficiently pronounced and isolated from other symptoms to justify a differential diagnosis during life; but I feel satisfied that, in a considerable proportion of instances, they are adequate to conduct us to a trustworthy conclusion, without any aid from the confessions of the *post mortem* room. When a patient, of cancerous dyscrasia, and with an open sore, say on the breast, becomes depressed and demented, slowly loses power on one side, complains of acute intracranial pain, and suffers from convulsive attacks, we have no hesitation in asserting, that a cancerous deposit has taken place in the brain. Cancer of the brain can, therefore, be diagnosed during life. The question is, can its existence be ascertained without the clue afforded by the open sore? Is there anything in the symptoms which points to cancer more directly than to other coarse lesions of the supreme nerve centre? In considering that question, it is, I think, necessary to divide all the symptoms which have been observed and described in cases of cerebral cancer into two great groups: the necessary and the contingent; the universal and the particular. The position of the tumour in the brain-mass, its rate of increase, the degree of irritation which it excites, or of softening which it induces, imply, of course, an infinite variety of what may be termed secondary or accidental symptoms, which will vary in every individual case. But the nature of the tumour, the cachexia in which it originates, the intracranial pressure caused by its growth, will be expressed in primary and essential symptoms, which will be identical in all cases. Putting aside the contingent, let us apply ourselves, for a few minutes, to the necessary symptoms. And first, as to the mental symptoms. You know that the cancerous cachexia has a peculiar mental, as well as bodily complexion associated with it: the emotions are dark and sallow, as well as the skin; a captious temper and a despondent tone of feeling are indeed as indicative of that change in the blood, or in nutrition, in which the cachexia consists, as any of the other somewhat vague signs by which it is supposed to manifest itself. When insanity occurs in persons of strongly cancerous cachexia, it is almost invariably of the melancholic type; and when the cancerous cachexia is strongly developed in an already insane person, melancholia is most frequently superinduced. Some of you will be able to recall to mind, as examples of these statements, Selina H., lately in Ward 25, who, in the course of a mammary cancer, gradually passed from a native sombreness of disposition to abject misery, leading to repeated suicidal attempts, and shaping itself into delusions, such as that her tongue was removed, so that she could never speak again; and Ralph W., who died in 18 Ward, and who, from being a cheerful, excitable old man, became wretched, and querulous and forlorn, when attacked by malignant disease of the pancreas. In cancerous marasmus, when anæmia and emaciation, and exhaustion, add their evil influence to that of the cachexia, the despondency proper to the cachexia is intensified, and is accompanied by a sense of mental debility that ultimately converts it into a settled despair. Few scenes more harrowing can, I think, be witnessed, than the death-bed of a patient who is consciously sinking under malignant disease, and who is writhing under physical and mental anguish.

In cancer of the brain, however, such a depth of despondency is rarely reached, and that is, first, because cancerous marasmus is not often attained, the tumour killing by its position before exhaustion is induced; and secondly, because the feelings are deadened and obliterated by the progress of the tumour. A moderate degree of melancholia, due to the cachectic state, and to interference with the cerebral blood-supply by the presence of the tumour of the cranium, is the first necessary mental symptom of cancer of the brain. That symptom has been present in the most marked manner in the four cases of cancer of the brain which have come under my observation. It is casually noted in many cases which I find recorded, and where it is not referred to it is fair to deduce, from the meagreness of the records, that it, like many other phenomena, had been overlooked. When a patient is seeking advice for what is presumed to be a bodily ailment, he will not, unless questioned, afford any information as to his mental experience, and, least of all, will he obtrude on notice a moderate degree of depression of spirits, which he does not himself regard as morbid, and perhaps connects with some real circumstance or event. Hence, moderate depression in cancer of the brain may well have escaped detection in many cases in which it existed.

The second reason which I gave for the moderate degree of the melancholia—which I have told you I look upon as the first necessary symptom of cancer of the brain—was because the feelings become blunted and obliterated as the malady advances. That statement contains the key to the second necessary or universal symptom, which is progressive mental weakness. I need not tell you that the effect of great pressure on the brain, as by depression of bone or lodgment of a large foreign body, is total suspension of its functions and unconsciousness; nor need I remind you that that power of subsequent accommodation, which the brain manifests when the pressure exerted upon it is slight in amount, is only displayed to a limited extent. A material amount of pressure cannot be exerted upon the brain without interfering with its action, and a steadily augmenting pressure will inevitably entail a steadily diminishing freedom of mental power. The brain must have room in which to work; and if that room be circumscribed, so will be its performances. We see this in hypertrophy of the brain, in apoplexy, in arachnoid cysts—and we may see it, if we look for it, in cancer of the brain. Whenever the magnitude of the tumour is such as to cause displacement or pressure, the mind becomes slow and laborious in action, like a heart beating in a water-logged pericardium. When its dimensions expand, the mind is more and more crippled and confined; memory fails, judgment is dethroned; attention can be fixed only by a grievous effort. At last, when the conquering tumour still further asserts itself, the mind is abolished, deep coma and then death ensue.

The case which formed the text of this lecture clearly exemplifies the presence of progressive mental weakness as causes of the cancer, and so did the three other cases which I have seen. Progressive fatuity I find also hinted at, or expressly designated, in many recorded cases of cancer of the brain. In all the ten cases of this disease, which Dr. John W. Ogle has reported with his usual faithful minuteness in the *Journal of Mental Science*, there were distinct evidences of failure of mental power. It was therefore with astonishment that at the end of his valuable observations I came upon these words, "Neither was there (in any case) during life (?) anything of the nature of mental imbecility, or any symptom of the various phases or forms of insanity." It seems to me that in every case there was indubitable mental imbecility, or rather progressive mental weakness, which is the second necessary and universal mental symptom of cancer of the brain, and which, with a moderate degree of melancholia, is the only necessary and universal mental symptom upon which I shall now insist.

Next, as to the necessary and universal bodily symptoms. Most important of these is pain in the head, severe and persistent, and confined to one side or region. This pain, although always present, is liable to exacerbations at irregular intervals, when it sometimes becomes excruciating, and compels moans and cries. Commencing with the first deposit of cancer, it continues with its growth, until sentency is impaired and destroyed by the incursion of paralysis or coma. As long as it remains it is aggravated by strenuous bodily or mental exercise, by excess in wine, or by startling impressions upon the senses. How this pain is produced, we can only speculate. We have been taught that the brain-substance is absolutely insensible; and more recently we have been assured by Sappey that there are sensitive *nervi nervorum*. It may be by irritation of these *nervi nervorum*, or of the central expansions of the afferent nerves in the brain-mass, which is not perhaps so utterly insensitive as has been asserted, or by reference from a displaced or compressed centre of sensibility, that the pain of cerebral cancer is produced. But, however that may be, its existence is invariable, and its character almost unique.

But besides pain, there are other necessary bodily symptoms; and of

these, slowly gravescent paralysis is the most significant. The first establishment of the paralysis is often sudden—after an attack of convulsions, or a brief period of unconsciousness—and is probably due to some movement of the tumour, to some extravasation of blood, or to some yielding of brain-tissue; but when once inaugurated, it proceeds onwards, compromising more and more the strength of the side which it affects, for it is always hemiplegic in character. In this it is unlike the paralysis due to clot, which is generally mitigated for a time after the formation of the clot when it is contracting, or is at least stationary. Also, it is unlike the paralysis of clot in this, that without any fresh seizure, slowly, not suddenly, it takes possession, in a mild degree, of the side of the body opposite to that on which it was first manifested. As the tumour swells, it compresses not merely that hemisphere in which it is located, but that upon which it has no hold, and thus secondarily impairs its ingredient power.

The third necessary bodily symptom to which we shall advert, is muscular tremor or convulsion at some stage of the malady. Either at its outset, or when its advance is hastened by the rupture of a vessel, or when the motor centres are mechanically disturbed, or when the tumour becomes an eccentric irritation to these motor centres, a state of muscular agitation, confined to a few short convulsions, or spread over continued tremblings, coarser than those of delirium tremens, and lasting for days or even weeks, betokens that the lower motor centres are in some way involved.

The fourth necessary bodily symptom, never absent at some part of the progress of the disease, is perversion or enfeeblement of one or more of the special senses. The sight becomes dim and impaired, or is altogether lost, most commonly in one eye; or is perverted so that illusions or hallucinations are seen; or hearing becomes dull, or strange noises in the ears are heard, or intolerable tastes and smells assail the palate and nose, or distressing itchings torment some part of the skin. Diverse in character though these experiences are, they have this in common, that a disturbance of sensation is the essential constituent in each, and that that disturbance is referable to a centric change; and such a disturbance is always encountered when the centric change is due to cancer. Reflection will convince you that it is impossible for a tumour to grow to any size within the skull without interfering with the origins of the efferent nerves, directly or indirectly, by drawing them into its fatal meshes, by cutting off their nutrient supplies, by kindling neuritis, or occasioning atrophy; and experience will confirm the conclusion to which reflection carries you. In all tumours of the brain modifications of sensation occur, and, *à fortiori*, in cancerous tumours of the brain. In every case, by patient investigation, they may, I believe, be brought to light. But exploring the central nervous system requires as much perseverance, and toil, and perspicacity, as exploring Central Africa; so it may well be that such symptoms have occasionally eluded observation.

The fifth and last necessary and universal bodily symptom in cancer of the brain is the cancerous cachexia, or that morbid constitutional state of the blood and tissues in which malignant growths occur, and in which several outward and recognisable manifestations are usually not wanting. The dark muddy complexion, wasted body, feeble muscles, morose manner, and gloomy countenance of the victim of any serious organic disease, at once create a suspicion that it is of malignant nature, and are sometimes so characteristic as to convert a suspicion into a certainty. When cancer of the brain is secondary, these outward signs are for the most part markedly present; and even when it is primary, they are, I should say, invariably exhibited to such an extent as to insure their recognition by a keen and practised eye.

Having completed the enumeration of the necessary and universal symptoms of cancer of the brain, let us recapitulate them. Given a case, in which you have the following symptoms: 1. Depression of spirits; 2. Progressive mental weakness; 3. Severe pain in one region of the head; 4. Slowly gravescent hemiplegic paralysis; 5. Epileptiform seizures; 6. Failure of one or more of the senses; 7. The cancerous cachexia—what is the disease? Only one reply can be given to that question—namely, cancer of the brain. The first six symptoms point unmistakably to tumour of the brain, and the seventh points to the nature of the tumour. I allow, however, that the difficulty lies just in the seventh symptom. There ought not to be insuperable difficulty in any case of cerebral cancer in determining that there is a tumour of the brain, but there may be an almost insurmountable difficulty in deciding that it is cancerous. To relieve us from this difficulty, we must look to our pathologists for earlier and more precise indications of the cancerous cachexia in the blood or secretions, or in the outward configuration. When these are obtained, we shall have less hesitation in diagnosing cancer of the brain, and in pronouncing when an intercranial growth, proved to exist, is not tubercular, or syphilitic, or fibro-plastic, or gliomatous, but cancerous. Pray remember, however, we are not

left without aids to diagnosis beyond these which I have yet mentioned. The case which I have supposed is of the worst description as regards diagnosis. In actual practice you will find not merely the seven cardinal symptoms upon which we have dwelt so long, but other minor or secondary symptoms, which, though subject to great variations, often furnish valuable corroborative testimony. These I can only name. They include delusions of every dye—aphasia, altered affections, sudden impulses, morbid appetites, ptosis, muscular twitchings or rigidity, staggering gait, glandular enlargements, vomiting, interrupted circulation, and menstrual irregularities. These, or some of these, may complicate, but more often simplify, diagnosis, which is again facilitated by the absence of other symptoms distinctive of other organic diseases, and by considerations as to the age, history, and habits of the person affected. To show you how contingent are interlarded with necessary symptoms, I shall briefly describe a case to you. Maria S., a widow, 54 years of age, was received into this asylum on the 4th May, 1869. She had then been out of health for some years, never having altogether shaken off the grief occasioned by her husband's death, until the close of 1868, when she became silly and childish, and suffered from a fit, and horrible pains in her head. In the spring following, she had several fits and attacks of transient excitement. She also, by imperceptible degrees, lost the use of her left side; and in consequence of this, as she persisted in moving about, suffered several falls. When brought here, she was a pale, sallow, anæmic-looking woman—so thin, as to be described in the case-book as a living skeleton. She was in a fatuous state. Though garrulous and anxious to talk, especially about her own illness, no reliable information could be got from her, as her memory was utterly fallacious. She could not recall her husband's Christian name, and was oblivious as to all measurements of time, and names of days, seasons, and places. She was often unable to find the word which she wanted, and seemed to introduce into the sentence for which that word was required any other word that occurred to her at random. Her expression was careworn, and also singular, as she had exophthalmos and blindness of the right eye, and ptosis of the right eyelid. The pupil of the left eye was much contracted, and its margin was irregular. The mouth was drawn to the left; and the tongue, when put out, pointed to the left. The sensibility of the right side of the face was much diminished. There was almost complete loss of power in the left arm and leg; there was partial loss of power, with tremor, in the right arm and leg; there was a systolic murmur at the base of the heart, and a thick white fur on the tongue. During her brief sojourn in the asylum, Maria S. became more and more fatuous; her appetite for food being at some times voracious, and at other times altogether absent, so that she had to be fed. The paralysis of the right side increased, uncontrollable vomiting and diarrhoea came on, coma supervened, and then came death on the 15th June. At the necropsy, the brain was found flattened and compressed, and at its base was a cancerous tumour. This took origin in the right temporo-sphenoidal lobe, and extended inwards, being divided by a neck into two parts, each about the size of a small walnut; an outer part, fibrous, containing cysts and fluid contents; and an inner part, soft and pulpy, and of a pink colour, variegated by deep red blotches, and with a delicate fibrous matrix. The outer part was embedded in the temporo-sphenoidal lobe, and the inner lay in a sort of excavation, which had been formed by the absorption of portions of the body of the sphenoid bone, the orbital plate, and the pituitary body, and by displacement of the posterior, orbital, and surrounding gyri. It compressed the roots of the right olfactory nerve, the right optic tract and nerve, and the right fifth nerve.

The prognosis in cerebral cancer, when it is diagnosed, is of course as gloomy as can be: nothing but death, speedy death, can be predicted. It is not, perhaps, utopian to hope that, with the progress of therapeutics, some means may be discovered of resolving or controlling malignant growths. Distinguished and sober-minded surgeons have entertained that hope. At present, however, we must be content with smoothing the pathway to the grave, and with retarding, if that may be, the passage of our patient along that miserable thorny road. If the nature and position of the tumour could be satisfactorily made out during life, its growth might perchance be slackened or arrested by frequent faradisation; change of climate might also be beneficial. It is curious that, while in tubercle that remedy has been, and is, most fashionable and successful, in cancer it should never have received a fair trial; and yet cancer appears to be highly susceptible to the influence of climate. Abounding in Europe, it is rare in Egypt, Algiers, Senegal, and Arabia: even in England its distribution is partial, as Mr. Haviland has conclusively proved. Haunting low lying grounds through which large rivers, prone to overflow their banks, descend to the sea, it eschews dry and elevated districts. Surely the progress and propagation of cancer might be sometimes checked by a resort to those climates which are

least favourable to its growth. One of the great objects in cancer of the brain, as in cancer of any other part, is to relieve pain; and that, after all, is best accomplished by the employment of opium, or some of its preparations or alkaloids. The hypodermic injection of morphia is an inestimable boon; nepenthe is a benefactor—it can at least confer an euthanasia. Cannabis Indica acts well; and so, under certain circumstances, do chlorodyne, chloral, and chloroform—the latter being, of course, used with extreme caution. When convulsions occur, bromide of potassium, in combination with tincture of sumbul, is beneficial; and, when delirium and excitement have to be combated, ergot may be had recourse to, or alcohol freely administered, for the delirium is sometimes the expression of exhaustion.

EXPERIMENTAL RESEARCHES IN CEREBRAL PHYSIOLOGY AND PATHOLOGY.

By DAVID FERRIER, M.D.,

Professor of Forensic Medicine in King's College; Junior Physician to the West London Hospital.

PRELIMINARY NOTICE.

THE opportunity kindly afforded me by Dr. Crichton Browne, of experimenting on over thirty guinea-pigs, rabbits, cats, and dogs, in the pathological laboratory of the West Riding Asylum, Wakefield, has enabled me to arrive at certain results and conclusions which seem worthy of a brief preliminary notice, pending the publication of details of method, experiments, and illustrations, in the West Riding Asylum Reports.

The following is a summary of the more important conclusions.

1. The anterior portions of the cerebral hemisphere are the chief centres of voluntary motion and the active outward manifestation of intelligence.
2. The individual convolutions are separate and distinct centres; and in certain definite groups of convolutions (to some extent indicated by the researches of Fritsch and Hitzig), and in corresponding regions of non-convoluted brains, are localised the centres for the various movements of the eyelids, the face, the mouth, the ear, the neck, the hand, foot, and tail. Striking differences corresponding with the habits of the animal are to be found in the differentiation of the centres. Thus the centres for the tail in dogs, the paw in cats, and the lips and mouth in rabbits, are highly differentiated and pronounced.
3. The action of the hemispheres is in general crossed; but certain movements of the mouth, tongue, and neck, are bilaterally co-ordinated from each cerebral hemisphere.
4. The proximate causes of the different epilepsies are, as Dr. Hughlings Jackson supposes, "discharging lesions" of the different centres in the cerebral hemispheres. The affection may be limited artificially to one muscle, or group of muscles, or may be made to involve all the muscles represented in the cerebral hemispheres, with foaming at the mouth, biting of the tongue, and loss of consciousness. When induced artificially in animals, the affection as a rule first invades the muscles most in voluntary use, in striking harmony with the clinical observations of Dr. Hughlings Jackson.
5. Chorea is of the same nature as epilepsy, dependent on momentary discharging lesions of the individual cerebral centres. In this respect, Dr. Hughlings Jackson's views are again experimentally confirmed.
6. The corpora striata have crossed action, and are centres for the muscles of the opposite side of the body. Powerful irritation of one causes rigid pleurosthotonus, the flexors predominating over the extensors.
7. The optic thalamus, fornix, hippocampus major, and the convolutions grouped around it, have no motor signification.
8. The optic lobes or corpora quadrigemina, besides being concerned with vision and the movements of the iris, are centres for the extensor muscles of the head, trunk, and legs. Irritation of these centres causes rigid opisthotonus.
9. The cerebellum is the co-ordinating centre for the muscles of the eyeball. Each separate lobule (in rabbits) is a distinct centre for special alterations of the optic axes.
10. On the integrity of these centres depends the maintenance of the equilibrium of the body.
11. Nystagmus, or oscillation of the eyeballs, is an epileptiform affection of the cerebellar oculo-motorial centres.
12. These results explain many hitherto obscure symptoms of cerebral disease, and enable us to localise with greater certainty many forms of cerebral lesion.