LECTURES

ON THE

PATHOLOGY, DIAGNOSIS, AND TREATMENT OF BRIGHT'S DISEASE.

By GEORGE JOHNSON, M.D., F.R.S.,
Physician to King's College Hospital; Professor of Medicine in King's
College, London; etc.

LECTURE VI.

Albuminuria not associated with what is commonly understood by Bright's Disease.—I. *Passive Congestion of the Kidney with Albuminuria, the Result of Impeded Venous Circulation.*—II. *Albuminuria and Hematuria from Embolism in the Minute Blood-Vessels of the Kidney.*—III. *Puerperal Albuminuria—Four Classes of Cases.*—IV. *Atrophy and Suppurative Inflammation of the Kidney from Retention of Urine—Symptoms and Diagnosis.*—V. *Acute Cystitis resembling Acute Bright's Disease—Symptoms, Diagnosis, and Treatment.*

BEFORE I go on to discuss the treatment of Bright's disease, which I shall do in my next lecture, I wish to direct your attention to certain cases of albuminuria resulting from various causes, but not associated with what is commonly understood as Bright's disease. My object in referring to these cases now, is to give you some hints which may assist you to distinguish them from each other and from cases of actual Bright's disease. My remarks upon these cases of albuminuria not associated with Bright's disease will have reference mainly to pathology and diagnosis, with only an occasional suggestion on the important subject of treatment.

I. *Passive Congestion of the Kidney with Albuminuria, the Result of Impeded Venous Circulation.*—There is a class of cases in which albuminuria results from passive congestion of the kidney consequent on some impediment to the return of blood through the systemic veins. The causes of this impediment are diverse. Valvular disease of the heart is one of the most frequent of them. Degeneration and consequent weakness of the muscular walls of the heart is a not uncommon cause. The impediment may originate in the lungs, as a result of emphysema with bronchitis, of extensive pneumonic consolidation, or of compression of one or both lungs by a copious liquid effusion into the pleura. Again, a dropsical effusion in the cavity of the peritoneum, the result of cirrhosis or other obstructive disease of the liver, may so compress the vena cava and impede the return of venous blood, as to cause passive renal congestion and albuminuria. In the advanced stages of pregnancy, the pressure of the uterus on the veins may cause passive renal congestion and albuminuria; but I shall presently refer more particularly to albuminuria in connexion with pregnancy. In general, the diagnosis of each of these causes of impeded circulation is not difficult. Then, as to the effect upon the urine and the kidney: the urine becomes scanty in proportion to the degree of venous congestion and the consequent tardiness of the blood-stream through the kidney. The secretion is usually high coloured, of high specific gravity, often turbid with urates, and more or less impregnated with albumen.

The mechanism of albuminuria consequent on the passive engorgement of the kidney which results from an impeded return of blood through the veins, may be illustrated by reference to Fig. 1, Lecture I. When, in consequence of an obstruction at the heart, the systemic veins become overfull, the distension of the renal vein, acting backwards through the intertubular capillaries, causes engorgement of the Malpighian capillaries, and a consequent transudation of serum through their walls. This serous transudation, mingling with the urine, renders it albuminous. Small hyaline and granular casts may be seen when the turbid urine has been cleared by warmth or by dilution with water. The Malpighian capillaries are sometimes ruptured by over-distension; the urine is then blood-tinged, and blood-casts are visible. The secondary character of the renal complication is usually apparent from the history of these cases; and the diagnosis may sometimes be confirmed by the fact that, when the circulation has been relieved by rest in bed, by hydragogues, by puncturing the legs, or by tapping the abdomen, the albumen disappears from the urine, to return, perhaps, when the circulation again becomes more embarrassed within the chest or by the reaccumulation of liquid in the peritoneum. I have seen this happen again and again during the progress of the same case. The first effect of passive congestion upon the kidney is to cause more

or less enlargement with some induration of the gland. The ultimate result of long continued congestion is atrophy and contraction, the surface of the kidney becoming uneven and finely granular as the wasting process goes on. The explanation of the phenomena is not difficult. For the due performance of its secretory function, and for the maintenance of its nutrition, it is essential that the blood move freely through the gland. An impeded return of blood through the veins involves, as a necessary consequence, a partial blood-stasis, and, as a result of this, a scanty secretion of urine, with impaired nutrition and atrophy of the gland. On microscopic examination of the kidneys, some of the tubes may be seen to be opaque with disintegrated epithelium and fibrine, some denuded, and in various stages of atrophy and contraction. When atrophy of the kidney has been a result of passive congestion consequent on a mechanical hindrance of the circulation, I have never found the muscular walls of the minute renal arteries hypertrophied.

II. *Albuminuria and Hematuria from Embolism in the Minute Blood-Vessels of the Kidney.*—There is yet another mode in which valvular disease of the heart may for a time render the urine albuminous and even bloody. You are aware that, when one of the valves of the heart has its surface roughened by inflammation or by senile degenerative changes, a very common result is a deposit of fibrine upon the roughened surface; and further, that these fibrinous deposits, having no organic union with the valve beneath, are very liable to become detached by the current of blood, and then to obstruct the vessels in any organ to which they may chance to be conveyed. One result of the mechanical plugging of blood-vessels is the formation of so-called "fibrinous deposits" in the kidney. The portion of kidney which is the seat of recent obstruction is raised above the level of the surrounding renal tissue; it is firm, anæmic, and of a yellowish white colour, with an intensely red injected margin. The older deposits are softer than the surrounding tissue, appear shrunk and depressed, and have not the red margin. In a still more advanced stage, the appearance of a deposit entirely passes away, and a depressed cicatrix is left.

On a microscopical examination of a recent fibrinous patch, the tubes in the seat of the deposit appear opaque from containing fibrinous coagula. Some tubes contain oil. Many of the intertubular capillaries contain fibrinous coagula; while others contain oil-globules, which are clustered in the form of rings surrounding the tubes. Granular coagula and oil-globules may also be seen in some of the Malpighian capillaries and the afferent arteries. The coagula in the vessels are more clearly seen after the tissues have been rendered transparent by dilute acetic acid. In the red vascular zone which surrounds the recent deposit, the Malpighian and the intertubular capillaries are seen to be injected and gorged with blood. The probable explanation of the phenomena is, that a soft fibrinous mass from a cardiac valve is arrested in a branch of a renal artery; and there it becomes disintegrated, and the fragments are carried on into the intertubular capillaries. The circulation through these vessels is arrested; there is, consequently, a backward engorgement of the Malpighian capillaries, and an albuminous and fibrinous effusion into the uriniferous tubes. The capillaries at the margin of the obstructed patch are greatly distended by the diverted current of blood, and in consequence the urine may become albuminous and even blood-tinged, hyaline and blood casts being visible under the microscope. Subsequently, the exudation into the tubes, the epithelium of the tubes, and the fibrinous coagula within the obstructed blood-vessels, undergo a fatty transformation, and all trace of the normal structure disappears. The fatty matter at length becomes absorbed, and a depressed cicatrix remains on the surface of the kidney. Two or more pale fibrinous patches of different dates may sometimes be found in the same kidney. In some instances, the fatty matter which results from the transformation of the tissues and the fibrinous coagula does not become absorbed, but remains encysted. This is the explanation of the cysts which are sometimes found filled with a thick dark liquid, composed of oil free and in cells, with which often plates of cholesteroline are mingled.

The diagnosis of embolism in the renal vessels is usually uncertain. We may suspect the occurrence when, with the physical signs of aortic or mitral disease, without great impediment of the general circulation, the urine suddenly becomes albuminous or bloody. In some cases, extensive embolism in one or both kidneys has been attended with severe lumbar pains, a scanty secretion of urine, and vomiting; but, when the obstructed portions of kidney are small, there may be no symptoms to indicate the occurrence of renal embolism.

III. *Puerperal Albuminuria.*—Since the time that Dr. Lever, in the *Guy's Hospital Reports* (1843), published the fact that puerperal convulsions are in a large proportion of cases associated with albuminuria, the subject of albuminuria in connexion with pregnancy has excited much interest. Later observations have established the essential accuracy of Dr. Lever's observations; but they have also shown that the

connexion between puerperal convulsions and albuminuria is not constant. Convulsions may occur without albuminuria; and, on the other hand, albuminuria in pregnant women may be unassociated with convulsions. The few remarks which I propose to address to you on this subject will have reference mainly to the subject of puerperal albuminuria, and only incidentally to the association of puerperal convulsions with albuminuria. With reference to the subject of puerperal convulsions, I advise you to read carefully Dr. Barnes's very able and interesting Lumleian Lectures *On the Convulsive Diseases of Women*. I have seen more or less of a considerable number of cases of albuminuria associated with pregnancy; and, looking over my notes of these cases, I find that they arrange themselves in four classes, each having in some respects a different history and pathology.

1. Women known to be suffering from chronic Bright's disease may become pregnant, pass through all the stages of pregnancy and parturition, and even suckle their infants, without accident or complication. A lady whom I saw some years since with my friend Dr. S. H. Steel of Abergavenny, while suffering from chronic Bright's disease supervening upon an acute attack which resulted from exposure to cold, twice became pregnant, each time had an uncomplicated labour, and gave birth to two healthy children, each of which she suckled for about a year, not only without detriment, but apparently with benefit to her health. In 1849, I first saw with the late Dr. Tanner a woman who during her tenth pregnancy had general dropsy with albuminuria. The labour was natural; the child was healthy, and nursed for a time. After that, while the urine continued to be albuminous, she twice became pregnant, and had uncomplicated labours. She ultimately died with contracted granular fat kidneys. It is probable that the existence of albuminuria from any cause increases the risk of puerperal convulsions at the time when to the exalted reflex excitability of the nervous system in the parturient woman there is superadded the disturbing element of violent and general muscular contraction; but cases like those which I have cited show that, when there is a free secretion of urine, with absence of uræmic symptoms, labour may be unattended with any serious complication.

2. In a second class of cases, during the latter months of pregnancy, there is more or less general oedema, with headache and other nervous symptoms, not unfrequently culminating in convulsions, which may recur again and again. The urine is scanty, high coloured, often turbid with urates, of high specific gravity, and contains a large amount of albumen. On microscopic examination, it is found to contain small hyaline casts, with a few granular casts, but few or no epithelial casts. After delivery, the urine quickly becomes copious, of pale colour, of low specific gravity, and within forty-eight hours the albumen may have entirely disappeared. Such a case I saw in the year 1857 with Dr. Greenhalgh and Mr. Peter Marshall. The most probable explanation of this class of cases is, that the pressure of the gravid uterus on the vena cava causes gradually increasing passive engorgement of the kidney, albuminuria, a scanty secretion of urine, dropsy, and at length uræmic convulsions. The rapid disappearance of the albuminuria and the other symptoms after the emptying of the uterus is explicable on no other theory than that of passive renal congestion consequent on mechanical pressure. Cases of this class are more common in primipare, and probably for the reason that in first pregnancies the abdominal walls are less yielding; there is, therefore, greater tension and greater pressure on the large venous trunks than during subsequent pregnancies, when the abdominal walls are more flaccid.

3. There is a third class of cases, in which the theory of mechanical pressure is not admissible. I allude to those cases in which albuminuria comes on at an early period of pregnancy, before the uterus has attained sufficient size and weight to interfere mechanically with the circulation through the kidney. In these cases there is evidence of acute desquamative nephritis. The urine is not only scanty and highly albuminous, but often blood-tinged, and contains epithelial and blood casts. Such a case I have in the hospital at the present time (March 1873). When albuminuria sets in during the progress of pregnancy, it is very apt to lead on to convulsions, to retinal hæmorrhage and albuminuric retinitis, with serious defect of vision. The renal symptoms may gradually pass away after delivery; if so, they may or may not return with the next pregnancy. In other cases, the albuminuria is persistent; the urine is of pale colour, of low specific gravity, and deposits small hyaline and granular casts. Ultimately, uræmic symptoms occur; and, after death, the kidneys are found either contracted and granular or large and pale. A painful case of this kind I saw not long since with Mr. Cadge of Norwich. The most probable explanation of this class of cases is that which refers the renal disease to some previous blood-change. Obviously, pregnant women are exposed to the ordinary exciting causes of renal disease; and acute Bright's disease

originating during pregnancy may result from exposure to cold and wet, from excessive eating and drinking, or from some zymotic blood-poison. But, in addition to these more common causes of albuminuria, it is probable that, connected with the evolution of the uterus and the development and growth of the foetus, there may sometimes be associated abnormal blood-changes, resulting in renal disease with albuminuria. In addition to other indications of the occasional occurrence of morbid states of blood in pregnant women, I may refer to those cases in which puerperal chorea is associated with acute endocarditis and fibrinous deposits in the mitral or aortic valves; the chorea being, in all probability, a result of capillary embolism in the region of one or both corpora striata.

4. There is yet a fourth class of cases, of which I have seen and noted several examples. I refer now to cases in which albuminuria and other symptoms of renal disease appear for the first time soon after delivery. Within a day or two after delivery, or after an interval of several days, sometimes after imprudent exposure to cold, a rigor occurs, and is followed by febrile symptoms. The urine is soon found to be scanty, with all the characters indicative of acute desquamative nephritis. There may be general dropsy, with or without uræmic nervous symptoms, such as headache and convulsions. The renal symptoms, after a period varying from a few weeks to several months, may gradually and entirely pass away; or the disease may become chronic, and result in a large white kidney. In such cases as this, the renal symptoms may with confidence be referred to the blood-contamination consequent on the absorption of morbid materials from the interior of the uterus after parturition. These cases are pathologically allied to, and sometimes associated with, a form of septicæmic puerperal fever. An interesting case of this kind, about which I was consulted, has been published by Messrs. Melland and Windsor of Manchester in the BRITISH MEDICAL JOURNAL (Sept. 12th, 1857). Here, too, I would suggest the probability that, when the foetus dies and is retained *in utero* until decomposition is commenced, there may sometimes be an absorption of foul gases and liquids, which in one case may give rise to the phenomena of *ante partum* phlegmasia dolens, as in a case which we have recently had in the hospital; and, in other instances, acute desquamative nephritis may result from this source of blood-infection.

It will be obvious that the distinction between the four classes of cases of albuminuria in connexion with pregnancy, which I have here briefly indicated, is of considerable practical importance, inasmuch as upon an exact diagnosis depends, not only the prognosis, but the treatment, of each case of puerperal albuminuria.

One hint I may give you with reference to the expediency of allowing a woman suffering from albuminuria to nurse her infant. The case of Dr. Steel's patient, to which I just now referred, shows that, when there are no symptoms of blood-poisoning, the mere fact of albuminuria does not prevent a woman from being a good nurse; but, on the other hand, when albuminuria is a result of a recent blood-infection, the mother's milk may become contaminated, and act as a poison to her infant. In one case about which I was consulted, a lady had acute renal disease, resulting probably from *post partum* absorption from the interior of the uterus. She recovered after a long illness complicated with pelvic cellulitis and abscess; but her infant, after taking the breast for five weeks, became feverish, and died with symptoms of pyæmia at the age of six weeks. It seemed probable that the infant's illness was a result of infection through the milk; the infection being not uræmic, but septicæmic—a consequence of the absorption of noxious uterine discharges. In cases similar to this, the mother should not be permitted to nurse her infant.

[To be concluded.]

OBSTETRIC MEMORANDA.

REPEATED ABNORMAL PRESENTATIONS.

THE following memorandum on this topic may be deemed worthy of record. A woman died here some years ago who had seven pregnancies, of which five terminated in "cross-births", one in cephalic presentation, and the other in an abortion at three or four months. No instruments were used at any time. A midwife attended all the labours, and appears to have turned on each occasion, and said there was "a bone in the way"; she was certified, and had a wide reputation for skill. The labour with cephalic presentation lasted four days, and resulted in the birth of a female (now aged 17), whose head was "pulpy", and who was apparently dead; her intellect is good, and she has not had any fits, but she is "nervous." All the other children were still-born.

STANLEY HAYNES, M.D., Malvern.