cess in an infant five weeks old, who had severe convulsions for nearly a fortnight. He was under its influence for nearly twenty-four hours, allowing him to wake to take food, and he emerged from the sleep perfectly well, and with no return of the fits."

"Dr. Williamson of Manchester has recorded the case of an infant six weeks old whom he kept under the influence of chloroform for sixty hours, sixteen ounces of chloroform having been used with complete success."

The following is a brief account of my own cases.

CASE I.—On March 10th, 1872, I was called to see a child aged two years, whom I found in severe convulsions of more than half an hour's duration. It appeared that, about two hours previously to the commencement of the fit, the child had eaten a hearty dinner of pork and fruit-pie, and that it had been sick once. The contractions of the muscles of the left side of the face and body were very severe: the face was very pale. I had the child placed in a warm bath, while a messenger was despatched to my house for chloroform. The bath not having had any beneficial effect, I proceeded to administer chloroform; the child was speedily brought under its influence, and the convulsions gradually disappeared. The child was placed in bed, and, shortly after its recovery from the anæsthetic effect of the chloroform, it fell into a calm sleep, which lasted two or three hours. A grain of calomel was then administered, followed by a dose of castor-oil; and on the following morning I found the child running about the house, and to all appearances well. The bowels had acted freely, and there had been no return of convulsions.

CASE II.—On September 2nd, 1872, I was called upon to see a child aged eighteen months, who had been in strong convulsions for five hours. I found it with strong interrupted contractions of the flexors of the limbs of the left side, and twitching of the muscles of the face. She was cutting some of her molar teeth. The friends had placed her in a warm bath before sending for me, but without benefit. I immediately administered chloroform, and the child passed imperceptibly into a comfortable sleep, from which she did not awake for three or four hours, when she seemed cheerful and intelligent, and without any sign of convulsion. A grain of calomel was then administered, followed by castor-oil, and the gums were lanced. On the following day the child seemed comparatively well; it had passed a good night, without any return of convulsions. The bowels not having acted freely, the castor-oil was repeated, and was followed by a motion resembling dirty green paint. On the third day the child was convalescent.

ON THE TREATMENT OF ULCERS OF THE LEG.

By J. GORDON BLACK, M.D. Lond.,

Surgeon to the Hospital for Sick Children, Newcastle-upon-Tyne.

THE perusal of the valuable reports which have appeared in the JOURNAL on the treatment of ulcers, at the various London hospitals, induces me to offer a few remarks, in the belief that good will accrue from the further ventilation of the subject.

It seems pretty generally admitted, that the treatment of ulcers of the leg, in the out-patient room, is unsatisfactory and disheartening. Whilst some instance the intemperate habits, the poorly fed and over-worked condition of the patients, to account for this want of success, I am more inclined to blame a wide-spread belief in the profession, as expressed by Mr. Lawson, of the Middlesex Hospital, that "for the effectual treatment of all ulcers of the leg, absolute rest of the limb is the first element." Having for some time past been in the habit of curing cases of this kind without requiring the patient to neglect his ordinary duties for a single day, I certainly cannot hold such a belief. If it be possible to cure a large ulcer of the leg without rest, and in quite as short a time, to say the least, as would be required to heal the same by recourse to the horizontal position, what becomes of the theory that absolute rest is necessary?

Moreover, during the time that the patient remains in bed, the circulation through the limb is rendered more efficient, and healing of the ulcer ensues; but no sooner are ordinary duties resumed, than the old conditions recur, bringing back with them the inevitable ulcer. A cure, therefore, under the absolute rest system, can scarcely be alleged, because it is not permanent; whereas, if an ulcer be healed without rest, it is clear that a cure has been effected, provided similar therapeutic conditions are maintained. This latter provision cannot, of course, be observed under the former plan of management.

Knowing, as I do, the easy practicability of healing ulcers without rest, I cannot think it proper to admit such ailments to the wards of a

hospital. To do so, seems wasting the funds of the institution, no less than the time and labour of the sufferer. At the same time, it is not creditable to surgery that such patients should be neglected, or given to understand that their weary and loathsome malady is incurable.

The plan which I adopt is practically the same as that recommended by Baynton, nearly eighty years ago, but with the important modification of using it antiseptically. Baynton's strapping has long been recognised as valuable and effective, and is described by Mr. Erichsen under the head of "Indolent ulcer." Such management, however, taxed too severely the time and patience of the surgeon, for its successful practice. Unless the plasters were very frequently removed (Mr. Erichsen says every forty-eight hours, at least), the pent-up discharge became very offensive, causing the dressings to be disagreeable, both to doctor and patient.

In order to avoid these disadvantages, I now warm the plasters by passing them through hot water, to which a little solution of carbolic acid has been added. The sore having been washed clean by the patient, is then saturated with weak solution of carbolic acid, and the straps, first treated as described, applied. The pieces of plaster (stout emplastrum saponis), should be two inches broad, and long enough to overlap four inches, after passing completely round the limb. They should be applied after the manner of a "Scot's dressing," from about three inches below the lowest diseased surface, to about the same distance above the highest. In their adjustment I think it most important to use no compression, but simply to lay them down evenly, so as to fit the limb accurately, and leave no creases in the plaster. Should pain be produced, the strap has been improperly applied, and must be at once removed. The bandaging of the limb, lightly and carefully, from the toes to the knee, finishes the dressing, which latter need not occupy more than ten minutes altogether. The patient may be told to return at the end of a week, when, on removal, the plasters will show only a slight moisture, instead of the profuse and offensive discharge seen when no antiseptic is used.

The advantages of the above plan of treatment are briefly these. It is cleanly; it saves the time and labour of the surgeon, for the dressings need rarely be changed oftener than once a week, and occupy only a few minutes. And finally, whilst the healing process is conducted with a minimum of pain and discomfort to the patient, he is in no way restricted from pursuing his ordinary occupations.

In cases of irritable ulcer, with much pain, Baynton recommended the sufferer to remove the bandage occasionally, and pour cold water upon the strapping for a few minutes, afterwards drying lightly with a soft towel, and reapplying the bandage. The plan is an excellent one, and usually very grateful to the patient's feelings.

Instead of employing carbolic acid, another good antiseptic may be used, namely, sulphurous acid. This is easily applied by playing upon the ulcer and surrounding diseased skin with a Dewar's spray-apparatus. The plasters may then be adjusted, after passing them through hot water, simply. A little smarting ensues, which, however, soon passes off. The effect of the sulphurous acid, in checking discharge and mal-odour, is quite as good as that of the carbolic, whilst its application is perhaps less troublesome and disagreeable. The sulphurous acid is especially suitable to ulcers of moderate size.

Baynton's strapping, especially when used antiseptically, may be employed for nearly all kinds of ulcers. The surface of a weak, indolent, or inflamed ulcer, speedily assumes a healthy appearance, without the preliminary use of astringent, soothing, or other lotions, being necessary. The most irritable sore may be stripped if care, and no compression, be used. Occasionally, however, it may be found advantageous to substitute linen or calico for the plaster straps.

For varicose ulcers, no treatment could be better. The horny edges of the "callous" variety quickly disappear, without recourse to such a dangerous excitant as blistering, which may easily set up unmanageable inflammation in the old or infirm. In eight or ten dressings, even very extensive ulcers may be healed by strapping, so that the cases must be few where skin grafting is really needed.

The administration of medicines internally is unnecessary. In most cases opium may be given to relieve pain, but the healing process goes on steadily, without such assistance.

I observe that Mr. Nourse, of Brighton, has used strapping and bandaging with great success. The plan, I feel sure, only requires more extensive employment to be better appreciated. For the frequent failure of the ordinary treatment by lotions, etc., patients are often blamed, because they do not strictly obey the instructions given. This neglect is, however, due rather to the well-known feebleness of such remedies, than to any lack of pains or inconvenience on the part of the sufferers. On the other hand, the interest which the latter manifest in carrying out directions under the treatment by strapping, is by no means the least recommendation of this method.