fused her food, tore her hair, pinched her skin, destroyed her clothing, was violent to those around her in a marked degree, cunning and malicious, foul in her language, and raved and fumed with passion, rendering it often necessary, during the paroxysms of excitement, to hold her, or to place her in the padded room for safety's sake. She was sleepless and noisy in the night time, and dirty in her habits. She entertained, also, some of the more common delusions, viz., that she was not a woman, that she was dead, etc., and was malevolent to her husband, whom she threatened with hideous punishments. She caused great anxiety on account of her proclivity to elude the surveillance of those in charge of her, in her endeavours to administer her own destruction. She lost flesh, became thin and wiry, and her hands and feet were cold and clammy. She was subjected to several modes of treatment, principally for the purpose of subduing the excitement. At different times she was packed and repacked. She underwent courses of the cold shower-bath several times. Chloral was administered in doses of from forty to sixty grains, night after night, without producing the least effect; at other times the drug was tried in the day time, in twenty-grain doses every two hours, but its administration proved equally futile; one day, as much as one hundred and twenty grains were given in this way, yet no rest or sleep was procured. Bromide of potassium was then administered, in half-drachm doses every three hours, but only to prove as great a failure as its predecessor. The same salt, in combination with cannabis Indica, was given with the same result. About thirteen months after her admission, hæmatoma appeared in the left ear, and followed the usual course, which has been already described. Although inactive, the swelling has shrunk but very little, and at present a very considerable cartilaginous tumour exists, involving the antihelix, concha, anti-tragus, and tragus, in an unshapely irregular mass. The meatus is quite patent, and hearing equally acute, as on the unaffected side.

The case was considered hopeless until August, 1872, when the excitement became gradually less, her conversation more coherent and rational, her sleep restored, and her habits improved. These conditions continued to develop a return of reason; and in October it was noted that she conducted herself with propriety, could comprehend, and act correctly; the most prominent symptom left being her inability to identify individuals, as she mistook persons before her for some of the actors in her previous history. About the middle of November, the remaining cloudiness of her perceptions cleared up, and she was then considered perfectly sane. A great improvement in her physical condition had also taken place. Continuing perfectly well, with her intellect seemingly as good as it had been originally, her memory vigorous, and moral sense restored, and in good health and condition, she was removed from the institution on the 6th February, 1873, cured.

Seeing that the system had sustained no shock, that no strong emotional affection had been perceived, and that no intercurrent disease had substituted itself, it is impossible to assign any particular agent which influenced the patient's condition and brought about her recovery, beyond systematic moral discipline.

RUPTURE OF THE JEJUNUM, FROM A FALL, IN A GIRL ELEVEN YEARS OF AGE.

By E. HOLLAND, M.D.Lond.

H. P., described as a delicate, lively, and "flyaway" child, fell, at 4 P.M., and struck the umbilical region forcibly against the edge of the stairs. She uttered a sharp cry, got up, walked upstairs alone, vomited the contents of the stomach untinged with blood, complained that she could not draw her breath, and lay down and slept for half-an-hour. On awaking, she got up, walked about, talked freely, and, though seeming faint, made no complaint of pain. At 9 P.M., she went to bed, without supper, and slept soundly till 4 A.M., when she awoke and asked for water. Afterwards, she slept till 6 A.M., when she awoke and again asked for water. At 8 A.M. she walked in her night-dress to an adjoining room, and, without complaint of pain, asked for water and ice. At 12, she walked alone down stairs and lay down on some chairs, and, complaining a little of pain, had some hot flannels applied, but requested them to be removed on account of the "weight". At 4 P.M., for the first time, she asked for something to eat, and, after taking a few mouthfuls of an egg pudding, she vomited, fell back pale, and died without movement.

At the necropsy, thirty-six hours after death, the abdomen was found to be distended with flatus and fluid, and the umbilical region discoloured by bruising. The peritoneal cavity was filled with a sero-flaky fluid, deeply tinged with bile. The peritoneum generally was minutely injected, and more or less universally covered with soft lymph. The jejunum was ruptured for nearly half of its circumference, twelve

inches from the pylorus, and presented a bruised appearance on each side of the rupture. On cutting open the gut, it was found healthy, and its mucous membrane uraised by extravasated blood in the neighbourhood of the rupture. There were no adhesions externally around the rupture, no thickening of the edges, and the latter dovetailed when laid side by side.

COMMENTS.—The interesting features of this case are—first, the slight cause, which can only have its results explained in the delicacy of fibre resulting from youth, and a state of health below par; secondly, the apparent absence of any severe suffering or shock, so that the friends did not consider her ill enough to require medical attendance until the fatal syncope set in, twenty-four hours after the inquiry; thirdly, the sudden termination, with vomiting, after taking food.

SELECTIONS FROM JOURNALS.

MIDWIFERY.

ACTION OF ERGOT ON THE BLADDER.—In an article in the *Centralblatt für die Medizinischen Wissenschaften*, May 24, Dr. Wernich of Berlin directs attention to the distension of the bladder which has very frequently been observed in cases of poisoning by ergot. The distension is attributed to the action of the drug on the vesical sphincter; and practical application of it had been made in the treatment of paralysis of the sphincter vesicæ after typhus, by Oppolzer; in the nocturnal enuresis of children and senile incontinence of urine dependent on simple weakness of the sphincter, by Clarus; and in paralysis of the sphincter in cases of paraplegia, by Barbier, Arnal, Monneret, and Brown-Séquard. The point on which Dr. Wernich especially insists is, that this action of ergot may produce an impediment to labour when it is given for the purpose of assisting this process. He relates two observations. In one, a woman pregnant for the fifth time, was in labour with twins. The pains having ceased for eight hours, and the os uteri being fully dilated, the first child was removed with ease; immediately after this, the bladder was emptied. To obtain the expulsion of the second child, ergot was given. In the course of three hours, severe labour-pains set in, but no progress was made. On revisiting the patient, Dr. Wernich found the bladder enormously distended, and drew off the urine. Immediately afterwards, a moderate pain brought the head below the brim of the pelvis, and labour was soon completed. In the second case, that of a primipara, the head remained deep in the pelvis, and the pains had ceased for some hours. The os uteri was fully dilated. Two doses of powdered ergot soon brought on pains; but, at the end of two hours, these had produced no effect. The bladder was found to be much distended; on passing the catheter, a very large quantity of clear urine escaped. The head of the child was at once propelled beyond the brim by the uterine contractions, and, as the woman was much prostrated, delivery was easily completed by the forceps. Dr. Wernich hence advises a careful examination of the bladder in all cases of labour where ergot is given, and the use of the catheter whenever the employment of the drug for some time is not followed by the expected result.

THERAPEUTICS.

ACTION OF COLD WATER ON THE SPLEEN.—Dr. F. Mosler (*Virchow's Archiv*, 1873, part 1) has arrived at the following conclusions from experiments on the action of water on the exposed spleens of animals. 1. The immediate contact of water with the normal spleen produces a visible contraction of the organ, varying in degree with the temperature of the water and the duration of the application. 2. In a less degree, cold water exerts the same action on the spleen through the intestinal walls. The effect of a cold douche is greater than that of the application of cold compresses or pieces of ice; probably the mechanical influence plays a part here. The action of water is inferior to that of quinine in causing contraction of the spleen. 3. Cold water also produces diminution in the size of splenic tumours, both acute and chronic. 4. The febrile paroxysm in ague may be arrested by cold douches applied after Fleury's method. 5. The cold douche does not supersede the use of quinine either in recent or in chronic cases of intermittent fever. 6. The therapeutic action of the cold douche in intermittent fever is not complete. It does not prevent relapses nor the formation of splenic tumours. 7. The splenic tumour in typhus is reduced in size by the use of cold water. 8. Much good is to be expected from a combination of the application of cold over the spleen, either in the form of ice or of the cold douche, with the administration of quinine.