LECTURES ON THE VARIETIES IN THE MUSCLES OF MAN.

Delivered at the Royal College of Surgeons of England.

By G. M. HUMPHRY, M.D., F.R.S.,

Lecturer on Anatomy and Physiology in the College; Professor of Anatomy in the University of Cambridge; etc.

LECTURE I.—Monday, June 2, 1873.

FEW subjects are more fertile in interest and instruction than that of variety in animal and vegetable forms; and few would tend more, if rightly apprehended, to throw light upon some of the great questions which perplex the morphologist and the physiologist. The varieties in any one structure or groups of structures are similar in nature to those in any other; and we cannot but feel that they are also similar to, and are held together by bonds of unity in the same manner as, those differences which constitute the peculiarities of the several groups, classes, orders, and species of the animal kingdom. All these, in their several degrees, are illustrations of variety—of the great principle of variety blended with, yet subordinate to, uniformity.

I endeavoured, to some extent, to point this out in the three lectures which I had last year the honour to give in this place*: to show, that is, that the muscles of man are but varieties from a simple pattern, which we find to be displayed in its greatest simplicity in some of the lower animal forms. Not that these are devoid of variety; for in these simple forms there is some variation from or modification of the simple pattern, adapting each for the particular place it has to occupy in the animal kingdom. I propose, in the three lectures I have now to give, to show that the varieties, or abnormalities as they are sometimes called, in the muscular system of man, are but modifications of the simple pattern, though differing from that modification which, being the more usual, we regard as the normal one. That this is so, may in most instances be clearly traced. To ascertain, however, the causes of the varieties, or the laws which govern them, is more than I can pretend to do. As I have just hinted, I make no doubt that they are of the same nature as the causes and laws which govern the diversities of the different animals; and, as in the case of these, I think it may be shown that they have a relation to utility which may be expressed in the following way. Taking the normal standard as the most perfect or most useful, those deviations from it are the most frequent which will least interfere with the perfection or usefulness of the muscle concerned; and, as a sequence of this, the muscles which have the most definitely specialised action, or rather those in which the definitely specialised action is most important to the economy, are the least frequently absent, and are, on the whole, the least liable to variety.

The materials which have been collected by the persevering labours of Wood in particular, and also of Meckel, Gruber, Henle, McWhinnie, Macalister, Turner, and many others, are amply sufficient to enable us to arrive at some generalisations; and my inability to accomplish much in this way will be due less to a deficiency of facts than to want of power to anneal them well together into good and consistent theory.

It has been remarked, and observation confirms the remark, that no two things are alike, and that variety is the order of Nature. It has been asserted, and the assertion has been questioned, that variety is an essential element in material form and quality. A little reflection will, I think, show that it must necessarily be so. No two particles can be precisely alike, because they cannot occupy the same space, and cannot, therefore, have the same surroundings. Those near the circumference of a sphere, for instance, must differ from those near the centre, owing to the differences of attractive forces operating upon them. The difference thus initiated is capable of indefinite multiplication; it will be manifested most clearly in the bodies which are formed by the greatest aggregation of particles; and it will be in increasing ratio in proportion to the complex nature and relation of the bodies. Hence it becomes apparent, even to the unaided eye, that no two blades of grass and no two muscles are alike; and we perceive that, in the nature of things, they cannot be alike. It is not, however, with this subtle kind of variety that I propose to occupy your attention; nor, on the other hand, with those varieties which come under the head of malformation of the body or monstrosity, but rather with the varie-

ties of muscles usually understood as such, and comprising duplicity, anomalies of attachment or connexion with one another, etc. There may be no bold barrier-line separating them from differences of size and form on the one hand, and from monstrosities on the other; but, practically, they are sufficiently distinct from both, and are recognised to be so; and it will be found that, as a general rule, they do not markedly interfere with the function of the muscle or muscles concerned.

Before entering upon the consideration of the individual varieties, I will make a few reflections which have arisen in the course of the investigation, and to which I shall from time to time have occasion to refer.

1. We may, as has just been remarked, fairly assume that the ordinary standard of muscular anatomy in man, as in each other animal, is the best, the one most fitted to the movements which have to be performed. Hence the varieties in and deviations from the standard are not changes for the better, but, for the most part, are for the worse, though the variety may not be attended with any manifest deterioration or imperfection of function. The perfection of the standard is attained by the requisite segmentation of muscles from the simple primitive mass, and by the concentration of the fibres, by means of tendons, upon the parts where the force may be most advantageously exerted. And the varieties are due, in most instances, to imperfection of segmentation, or to a want of proper concentration of fibres, which sometimes presents itself in the form of too great segmentation or in too great a range of attachment.

The principle of subdivision, with, at the same time, that of concentration—of subdivision for the purpose of executing varied movements, and of concentration for the purpose of executing them in the most favourable manner and with the least expenditure of force—is carried to its utmost in the upper limb and, especially, in the hand. Here, therefore, we may expect to find the greatest frequency and the greatest number of varieties. In the trunk, where the opposite conditions occur, where specialisation is at its minimum, and where there is little difference in the disposition of the muscles in different animals, varieties are least frequent and fewest.

2. An imperfection of segmentation is, as we should expect, most frequently observed in the case of muscles the fibres of which—the fibres, that is, of two or more adjacent muscles—are parallel, and which are in the habit of acting simultaneously or nearly so, such as the pectoralis major and the deltoid, and more particularly the two radial extensors of the wrist; also the peroneus tertius and the extensor digitorum pedis: whereas the abdominal muscles, the fibres of which take different directions, and the peronei which act upon the different parts of the foot and in different modes, rarely present examples of imperfect segmentation from one another.

An imperfection in concentration is illustrated in the simplest manner by a want of due limitation of the attachment of a muscle, as when the coraco-brachialis extends down the humerus to the inner condyle, or when the biceps has an additional origin from the humerus, or the peronei extend upon the toes. This may be attended with additional cleavage, leading to further subdivision of muscles; as when a third radial extensor of the carpus is formed, or when a segment of the tibialis anticus is inserted into the first phalanx of the hallux, or when an additional stylo-glossus or stylo-hyoid is present. Sometimes supernumerary muscles are found, which are scarcely the result of wider extension and additional cleavage of the ordinary muscles, but still are to be referred to the persistence and development of some part of the embryonic or primitive muscular mass, which usually becomes absorbed, and makes way for the fuller growth of the regular muscles. Hence the presence of these supernumeraries is often associated with imperfect development of the regulars; or they may be due to what may be called a hypertrophic development of the muscular system, in which case the presence of supernumerary muscles is associated, not with a lower, but with a higher, amount of growth of the regular muscles. Accordingly, in several of the instances in which supernumerary muscles were found, it was observed that the general muscular development was unusually great. The supernumerary muscles, if not the mere segments from or the reduplications of the ordinary muscles, are usually the obvious adjuncts to, or substitutes for, them. There are, however, some instances in which this can scarcely be said to be the case.

3. The muscles which are most frequently wanting are, on the whole, those which are most easily spared—the muscles, that is, for which substitutes are most easily found, or which do not play any very definite or important part in the movements of the body. Such are the pyramidalis abdominis, the psoas parvus, the palmaris longus and the plantaris, the extensor primii internodii pollicis, the peroneus tertius, the omo-hyoid, the stylo-hyoid, and the gemellus superior. It will be

* See BRITISH MEDICAL JOURNAL, 1872, vol. i, pp. 657 and 685; vol. ii, pp. 4, 33, 57, and 85.

remarked, that of these some are, as it were, lingering remnants of muscles which are more developed in other animals, and which seem to be present in man rather because they are part of the animal muscular system, than because they serve any special purpose in him. Some, on the contrary, are developments peculiar to him or nearly so, are initiated in him, and seem scarcely to have established themselves as stable elements of the animal muscular system; whereas others do not admit of being classed under either of these heads.

4. With regard to the correlation of muscular varieties, it is well known that several are often present in the same subject, and that they are often associated with varieties in the other structures, as in the blood-vessels and nerves, and even in the bones.* Moreover, the same kind of variety is likely to be repeated in the same person; that is to say, extensions of attachment, additional cleavages, or supernumerary muscles, are likely to be associated, or to appear at different parts of one subject. And the same is true of the variations in which the contrary tendency—viz., deficiency of development—is the leading feature.† To this, however, there are many exceptions; it being no uncommon thing for deficiency in one part to be associated with excess in another.‡ I have already mentioned that the presence of a supernumerary muscle is sometimes associated with a deficiency in an adjacent muscle, the function of which it shares or performs.

Moreover, the varieties are, as a rule, symmetrical. At least, in rather more than half the instances this is found to be the case. It is, however, to be remarked, that corresponding varieties are not commonly found in the upper and lower limbs, or on the opposite (flexor and extensor) surfaces of the same limb. In other words, the varieties do not usually affect the serially homologous muscles in the upper and lower limbs of the same person, or the antagonistically corresponding muscles in the same limb. Thus, in the tables given by Wood, there are seven instances of varieties of the *lumbrales pedis*, and in only one was there any variety in the *lumbrales manus*. In that case, the third *lumbralis* was double in both hands and in both feet. In the majority of the cases in which the slip from the extensor carpi ulnaris to the extensor tendon of the little finger was present, the homologous slip from the peroneus tertius to the extensor tendon of the little toe was absent. Occasionally, this slip passed to the metacarpal or to the metatarsal bone, instead of to the extensor tendon; but I do not find any instance in which the like anomaly occurred in the upper and lower limbs of the same person. Frequent as are the varieties in the flexors of the digits in the upper and in the lower limbs, an absence of the superficial flexor to the fifth digit afforded the only instance of a similar abnormality in these muscles in the two limbs; and I do not note any instance of a corresponding variety in the interossei of the two limbs. We are the more surprised at the frequent want of correspondence in the varieties in the muscles of the two limbs, forasmuch as we observe that the foot and the hand in the same persons are often the seat of similar deformities from excess, deficiency, or variation in development of the digits.§ We learn from this that we must not look to varieties as a source of much information respecting the homologous disposition of the muscles in the two limbs. I do not mean to say that they are altogether fruitless in this respect; for we shall find that the varieties in a muscle in one limb, upper or lower, are not unfrequently reminders of the usual disposition of the serially homologous muscle in the other limb.

I have searched in vain to make out any other correlation of varieties of muscles beyond those which I have mentioned. Thus, the discovery of a variety in a particular muscle suggests to us that there are, probably, other varieties in the muscular, the vascular, the nervous, and the other systems of the body; that these varieties in the muscles of other parts are probably of the like nature to that in the one observed, and are probably symmetrical; but we cannot infer that any particular muscle—scarcely even the serially homologous muscle in the other limb—is likely to be affected.

It appears that males and females, and the two sides of the body, are about equally liable to be the subject of varieties. We scarcely yet know in what class of persons they are most likely to appear, or whether they are more common in any one class than in others. We should expect to meet with them in those persons who present any obvious de-

formity or defect of development; and this receives some support from Mr. Carver's case of the idiot before mentioned. Judging from the dissection of the Bushwoman by Flower and Murie,* and of the negro mentioned in *Guy's Hospital Reports*, vol. xiv, we may infer that they are at least as common in the other races of mankind as in the European; and, judging from the differing account of the dissections of members of the same species of several of the lower animals, especially of the Monkeys and Apes, by different anatomists, we may infer that varieties are as common in them as in Man; and it may be remarked that in animals as well as in Man the muscles not uncommonly differ on the two sides of the body.

We might, perhaps, on the whole, have anticipated that varieties would be most frequent in the higher orders of animals, because in them the departure from the simple type is greatest for the purpose of executing the more complex movements. There is not, however, sufficient evidence to enable us to form an opinion on this point; and we must remember that though in the higher animals the muscular dispositions are more complex, yet the special functions of the several parts of the muscular system are more definite; and this would, to some extent at any rate, counterbalance the tendency to variety which the greater complexity might engender. It seems, indeed, to be shown by the dissections of different animals, including Man, that there is less fixity, more license of development, in the muscular than in the other structures of the body, except perhaps the blood-vessels. This accords with the fact, that the variations in the muscles in the different classes are greater than those of the other structures; are greater, I think, than would have been expected from merely taking the variety of movements into account; are greater, that is, than would appear to be necessary. For instance, there seems to be no reason for the varieties in the flexor muscles of the digits being so great in the different classes of animals. We learn from this two things—first, that the tendency to variety in the formative processes is greater than we should have anticipated from a superficial observation of animal form; and that it is limited or restrained only, as remarked in my *Observations on Myology*, p. 178, “by the high controlling forces which insure the requisite subservience to utility”. Secondly, the varieties in the several members of each class are more numerous in those structures which vary most in the different classes of the animal kingdom. I have already mentioned that the muscles vary most in those parts of each animal in which the varieties are greatest in the different animals, as the limbs, and, more particularly, the distal parts of the limb.

Moreover, the varieties themselves are very variable; that is, an unusual muscular factor, such as an additional head of the biceps brachii, differs a good deal in the different instances in which it is found. Still, there are lines, or “directions”, as they have been called by Wood, which the variation most frequently takes; and these are so often in the direction of muscles normally existing in the lower animals, as to be strongly suggestive of a common origin of the muscular system in the several members of the animal kingdom. All that we can say, however, with any degree of certainty is, that they are illustrations and results of the similarity of the forces which are in operation in the evolution of each of the several animals. The forces which evolve the coraco-brachialis of a Man and of an Echidna are similar; the materials are similar, and the surroundings are similar. It is probable, therefore, that the variations in the human coraco-brachial will present a resemblance more or less close to that muscle in the Echidna; and it does not necessarily follow that the muscle in the one instance has been evolved from that in the other, and that the variation is consequently a reversion. It may be asked, and with some reason, why, then, does not the coraco-brachial of the Echidna sometimes vary in the direction of that of Man? Perhaps it does so. At present we know too little of the varieties in animals to draw inferences from them. Still, we should judge it to be more probable that the higher and more nicely adjusted frame of Man would vary in the direction of the lower animals, than that the reverse should be the case. We should also expect that an animal would be more likely to vary in the direction of another near to it in the animal scale, and resembling it, than of one more remote and dissimilar, because the forces which evolve the two are more alike in the one instance than in the other. Unquestionably the variations in Man are not unfrequently reminders of the anatomy of the Ape; though somewhat too much has been made of this, for these variations are often reminders of the anatomy of other animals as well as of the Ape. Often the variations are reminders of the anatomy of other animals, and not of the Ape; sometimes of animals far distant in the zoological scale. Often, too, they are *sui generis*, and not reminders of the anatomy of any other animal.

* See case of an idiot, by Mr. Carver, in the *Journal of Anatomy*, iii, 257, in which abnormalities were presented by the bones of the skull, as well as by the muscles, vessels, and nerves.

† In each of the thirteen varieties recorded by Meckel (*Arch.*, v, 115), as occurring in the same subject, the anomaly was an excess of attachment, a supernumerary muscle, a preternatural cleavage, or an extension as of the *psaos parvus* to the femur. Moreover, all the muscles in this subject were unusually strong.

‡ See Wood, *Proceedings of the Royal Society*, June 6th, 1844, p. 299. Hallett found the peroneus tertius absent in a case of many anomalies by excess. Meckel found the biceps in one arm with three heads, and in the opposite arm with only one.

§ For examples of this, see Dr. J. W. Ogle's paper on Hereditary Transmission of Structural Peculiarities, *British and Foreign Medico-Chirurgical Review*, April 1872.

* *Journal of Anatomy*, i, 189. Wood found very few irregularities in a Negro and in a Lascar dissected by him.—*Proceedings of the Royal Society*, 1865, p. 386.

Much, therefore, as we may be tempted to adduce variations in the muscular system as additional arguments in favour of the evolution of man and animals from a common origin, it is perhaps more philosophical to refrain for the present, and to leave the mind unfettered by the bias attendant upon the consideration of this great subject, which, in the present epoch of scientific inquiry, tends rather too much to engross the thoughts, and which sometimes casts too strong a shadow upon the work, and too obviously obscures the view, of the investigators of Nature.

MUSCLES OF ABDOMEN.

I will consider, first, the varieties in the muscles of the trunk, commencing with those of the abdomen. They are few and infrequent, in accordance with the simple disposition of these muscles, with their comparatively slight deviation from the primitive, unstratified, or partially stratified, transversely segmented layer, which we find in the Cryptobranch, and with the little variation in structure, associated with little variation in function, which they present in different animals.

The most noteworthy varieties are the occasional persistence of inscriptions in the obliquus externus in the neighbourhood of the lower ribs, and contiguous with their cartilages. In one of these inscriptions a piece of cartilage was found by Henle. It may seem rather remarkable that such inscriptions are not more frequently met with in this and the other abdominal muscles; but it must be remembered that in the lower animals the disappearance of the transverse septa, of which the inscriptions are remnants, is usually associated with the stratification of the muscular mass, and with the difference in the direction of the component fibres of the several strata.

Additional slips are occasionally present, segmented from the normal muscles. Such a one is described in *Guy's Hospital Reports*, vol. xvi, p. 149, passing from the cartilage of the twelfth costal cartilage to the ilium between the obliquus externus and the obliquus internus. The most frequent varieties in the abdominal wall are presented by the two muscles which are most easily spared, and which are, therefore, least impressed into regularity by the constraining influence of utility—viz., the pyramidalis and the psoas parvus. Of these, the first is often (in one case out of four) absent on one or both sides, its place being usually supplied by an increased breadth in the rectus. It, moreover, varies in size, and is sometimes double. The psoas parvus is absent in one or both sides in nearly half the number of subjects examined. It has been found in some instances extending beyond the ordinary range of its origin. It is sometimes divided longitudinally into two muscles; and in some instances it extends over the pubes in company with the psoas magnus, and blended with it,* or inserted separately from it into the line descending from the trochanter minor to the linea aspera.† These two—psoas magnus and psoas parvus—it may be remembered, are both parts of, what I have called in my lectures last year, the internal or transversalis stratum of the abdominal wall. The psoas parvus not infrequently preponderates in lower animals, especially in some of the Carnivora, in which the movements of the pelvis upon the spine are free and strong. In Man, the femoral, or psoas magnus, portion nearly, often quite, pushes the pelvic, or psoas parvus, portion out of the field. The latter is not, however, always thus stunted by its greater neighbour; for in some instances, as just said, it encroaches upon the territory of the psoas magnus, extending with it to the femur, and being inserted with it, or near it, into the linea aspera.

It may be remarked that the pyramidalis is very inconstant in Mammals, being more frequently absent than present; as, indeed, is the spine of the pubes, with which, or with the marsupial bone, it is often associated.‡ Any inferences which might be drawn from this in relation to its frequent absence in Man, are, however, weakened by the fact that the other muscle—the psoas parvus—is very generally present in Mammals, and often attains to considerable size. In Birds, both are absent.

The absence of the transversalis abdominis has been noted in one instance by Macalister. Considering the remarkable disposition of the oblique and transverse muscles with regard to the rectus, and that it differs from the disposition which prevails in many animals, among others the Chimpanzee, in which the internal oblique passes entirely upon the superficial aspect of the rectus, we should have expected to find varieties here to be rather frequent. This, however, appears not to be the case; at any rate, they have not attracted attention. Neither do I find notice of varieties in the external oblique, the diaphragm, or the intercostals, which are of sufficient importance to occupy our time.

[To be concluded.]

CLINICAL LECTURE

ON

CHRONIC BASILAR MENINGITIS.

Delivered at the Belle Vue Hospital Medical College, New York.

By WILLIAM A. HAMMOND, M.D.,

Professor of Diseases of the Mind and Nervous System, and of Clinical Medicine.

THE patient whom I have now the opportunity of showing you, first made her appearance at this clinique in the winter of 1871-72, about a year ago. At that time she was suffering from deep-seated pains in the head, vertigo, and paralysis of the third nerve on the left side, as evidenced by ptosis, dilatation of the pupil, and external strabismus, the latter condition producing diplopia. Conjoined with these symptoms, there was slight but decided paralysis of the muscles of the face, arm, and leg of the opposite side, together with cutaneous anesthesia.

The history of the case was not that of cerebral hæmorrhage or of embolism, for inquiry showed that the symptoms had been of very gradual development, a fact not consistent with the existence of either of the lesions mentioned. I was disposed to think that the affection was an aneurism of the left posterior communicating or posterior cerebral artery, both of which, as you know, are in very intimate topographical relation with the crus cerebri; a tumour of some other kind, involving the left crus, or pressing upon it; or chronic basilar meningitis with effusion, implicating the same part. I regarded the latter as the most probable morbid condition, for the reason mainly that the symptoms were not so intense as regarded paralysis and pain as are those which result from either of the other diseases named.

The explanation of the phenomena observed in this case some of you will doubtless recollect; but, as the patient illustrates important physiological and pathological facts, and as most of you have not seen her before, it will not be out of place for me briefly to touch upon the relation of the observed symptoms to the supposed pathological condition.

In the first place there were ptosis, external strabismus, and dilated pupil of the left side. These circumstances indicated the involution of the third pair of nerves in the lesion, probably of the same side, even if there were no accompanying symptoms but with these latter, with very considerable certainty.

The third pair of nerves has its apparent origin in the crura cerebri—the right nerve from the right crus, and the left nerve from the left crus. If, however, the fibres be followed out by minute dissection, as has been done by Vulpian, they are seen to be arranged into three groups. Of these, the middle and posterior decussate, after passing entirely through the crus, while the anterior group passes forward to the optic thalamus, in which ganglion the fibres are lost. None of the fibres of origin originate in the crus, and this latter may be entirely dissected away, and the third nerve be left intact.

Now, if Vulpian be right in his view, that a considerable number of the fibres of the third nerve decussate, any disease of the brain affecting these fibres must be manifested by paralysis of the muscles supplied by the nerve of the opposite side; and, as the motor and sensory fibres of the spinal cord decussate below the point at which the third nerves decussate, the disease, if causing paralysis of other parts of the body, would induce this condition also on the opposite side, or, in other words, on the same side with the paralysis of the muscles supplied by the third nerve. But in the case before us the reverse is the fact, the muscles of the eye, and those of the face, arm, and leg, being affected on opposite sides. The disease, therefore, whatever its nature, must be confined entirely to the left side of the brain.

The left crus cerebri contains motor and sensory fibres coming from the right side of the body, below the medulla oblongata. It contains the fibres of the left third nerve; disease involving the left crus would, therefore, cause paralysis of the muscles supplied by the left third nerve, and of motion and sensation in the left half of the body. And this is exactly what existed in the patient before us a year ago.

As we have seen, the anterior group of the fibres of origin of the third nerve come from the optic thalamus of the same side, and a lesion affecting this ganglion would give rise to a similar condition so far as the paralysis is concerned. But it is difficult to conceive what morbid process could go on in the optic thalamus, and cause such a group of phenomena, so gradually developed as those which mark this case, and not at the same time produce more extensive disturbance. Taking all the facts into consideration, I thought the theory of basilar meningitis,

* Meckel's *Archiv*, v, 116.

† *Guy's Hospital Reports*, xiv.

‡ See remarks on this, *Journal of Anatomy*, vi, 312; and *Observations on Myology*, p. 124.