and, by the last statistics, it would appear that, during the past three years, at St. Bartholomew's, as low a rate as 18 per cent. has been obtained, and Mr. Callender, by the admirable method of treatment of stumps, which he has, for some time past, adopted, appears to be likely to reduce this rate of amputation-mortality to a considerably lower level still; in fact, to bring it within the limits of Simpson's returns from country and private practice.

Let it not be forgotten also that Guthrie's returns of primary amputations of all limbs, performed during the Peninsular war, show a mortality of only 1 in 12, or, in other words, for those amputations that are usually the most fatal, little more than 8 per cent.—a considerably lower rate than, according to Simpson, is obtained for all cases, inclusive of pathological cases in country practice. I believe that the fact is, that many amputations are performed in hospital practice, under circumstances both of injury and disease, in which consent would be refused by the patient and his friends in private. If we look at the table of amputations in this hospital, which I use as the basis for the observations I am making to you, I cannot but come to the conclusion that we have amputated rather freely, and that, under similar circumstances of injury and disease in private, some of the cases would have been left to die without amputation. However this may be, we have abundant evidence, independently of that furnished by Simpson's figures from the returns which I have already given you, of the results of par-turition, ovariectomy, and military amputations, to show that septic influences of the most destructive character are capable of being generated by the accumulation, under one roof, of people under certain conditions of natural process, of injury and of operation. It is to this condition so generated, and which might, so to speak, at any time be developed at will, grown, as it were, by the overcrowding of a ward with suppurating wounds, that we apply the term "hospitalism".

In the next lecture, we will proceed to the consideration of these conditions, their mode of production, and the influence they exercise on hospital mortality.

LETTSONIAN LECTURES

ON

SYPHILIS AS A CAUSE OF DISEASE OF THE NERVOUS SYSTEM.

Delivered before the Medical Society of London.

By W. H. BROADBENT, M.D., F.R.C.P.,

Physician and Lecturer on Medicine at St. Mary's Hospital.

LECTURE II.

GENTLEMEN,—In my last lecture, I considered the various features of syphilis which will bear on the consideration of this disease as it affects the nervous system, and more especially the nature of the morbid changes to which it gives rise, and the habits of locality and rate of progress of the morbid processes. It remains now for me to apply the general conclusions there given, and to trace the effects of syphilitic lesions in the different nerve-centres.

I showed, and illustrated by a few cases, the results when individual nerves or groups of nerves were involved. I did not attempt to distinguish between syphilitic neuralgia and neuralgia arising from other causes, and, so far as I am aware, no distinction exists, and the syphilitic origin is ascertained only by the syphilitic history. In paralysis of individual nerves, it is already a feature suggestive of a syphilitic source, when it is made out that the cause of the paralysis lies within the cranial or spinal cavity, and outside the nerve-centres.

Paralysis of the third nerve is so commonly a result of syphilitic disease about the base of the brain, that this is the first supposition which arises when a case presents itself, and it requires elimination before other hypotheses are entertained. The same might be said of the other cranial nerves, except the seventh; and, when more than one cranial nerve is affected, the presumption of syphilis increases. Corroborative evidence, if not at once yielded by history or co-existing lesions, would be sought in evidence of the constitutional condition produced by syphilis and in nocturnal pains. But I must proceed to that part of the subject which will engage our attention this evening.

Syphilitic Affections of the Spinal Cord.—The range of symptoms producible by disease of the cord is not very wide. The tendency of most is to paraplegia; and, to understand this, we have only to remember that, in some parts, the cord is not thicker than the little finger, and that consequently a limited area of inflammation, or a small tumour,

might involve a complete segment, which is all that is required to cut off the parts below from the volitional centres in the brain, and produce this form of paralysis.

Paraplegia is rather common as a result of syphilis (Bazin says that two-thirds of the cases are syphilitic), and, if taken early and treated energetically, it is usually curable. It is, therefore, very desirable that, if possible, the distinguishing features of syphilitic paraplegia should be pointed out. It has been given, as one of the characters of syphilitic paralysis generally, and of syphilitic paraplegia, that sensation is usually not impaired, and that reflex action persists; but this, if it were true as a matter of fact, which I shall show is not the case, would not constitute a peculiarity. It is the rule, in any form of disease of the nervous system, that the motor functions are more easily affected than the sensory. Even in nerves in which motor and sensory fibres are mixed up together, an injury, such as pressure, may destroy for some time the conductivity of motor impulses, while the only effect on the sensory functions is a temporary *dysæsthesia*. In experimenting on the cord, this difference has been found to be even more marked; division of the white fibres constituting the motor tract being followed by distinct and definite motor paralysis, while, in the case of the central grey matter which is the channel for sensations, a slight bridge suffices to convey upwards the sensory impressions; and, while there is any continuity of grey matter, *anæsthesia* is not produced. Again, it is the rule, in hemiplegia from cerebral lesion, that motion is more frequently and persistently impaired than sensation.

It is not in individual symptoms, nor in any arbitrary and unexplained difference in the phenomena, that we shall find the distinctive characters of syphilitic affections of any part of the nervous system, but in a comprehensive view of all the facts of the case.

It will not occupy much time, and will conduce to clearness, if I enumerate the principal diseases which affect the cord, with their leading characters.

Myelitis, inflammation of the substance of the cord, if general, gives rise to a progressive loss of motor-power from below upwards, with loss of sensation, preceded by tingling; the rectum and bladder, and their sphincters, sharing in the paralysis. There is little pain, at most a dull aching. Reflex action is abolished, and death occurs when the inflammation extends to the respiratory centres. Bed-sores come on early. A local myelitis, involving a segment of the cord, causes paralysis of the parts below, motor and sensory (or motor only for a time), with loss of control over the evacuations. Reflex action will persist, and may be exaggerated from concentration on the motor-cells of the anterior nerve-roots of impression, which normally would have been partly transmitted upwards to the brain; the paralysed limbs often start involuntarily. If there be pain, it is usually due, not to the myelitis, but to some extraneous cause, as, for example, to disease of the vertebræ.

Spinal meningitis is attended with pain, both along the spine and in the limbs, provoked by movements, especially such as involve motion of the spinal column. There are painful startings or tonic contractions of the limbs or muscles of the trunk, and sometimes severe tetanoid spasms; sensibility is usually intensified. Paraplegia comes on late and gradually, if at all. There is loss of power in the bladder, not usually in the rectum. Spinal congestion, according to Dr. Radcliffe, gives rise to sudden incomplete paraplegia, varying in degree, without loss of sensation or of control over the evacuations, but attended with aching in the back and limbs. There may, however, be a rapidly progressive and fatal paraplegia, which leaves evidences of congestion only.

Other diseases are sclerosis of the posterior columns and cornua, and of the anterior white columns and cornua, the former giving rise to locomotor ataxy, the latter to wasting palsy. There may, again, be softening of the cord, which, if non-inflammatory, is not to be distinguished from local chronic myelitis, or it may be the seat of tumours. Syphilis may give rise to myelitis or meningitis, or to the local sclerosis or gummatous tumour described by Dr. Moxon; but, before speaking of these, the question of the dependence of locomotor ataxy on syphilis may be considered. It has been stated that locomotor ataxy is a form of disease often set up in the cord by syphilis; but I find no support for this view in recorded cases, or in my own experience, and it is not the habit of syphilitic disease to follow, in its advance, any functional or structural arrangement in the part affected, or to confine itself to a particular vascular area. I have had under observation, for a long time, a woman suffering from ataxy, which I attribute to an attack of meningitis along the posterior aspect of the cord, possibly syphilitic; and it is easily conceivable that, in multiple syphilitic deposits in the cord, a sufficient number might be situated in the course of the posterior white column and grey matter to impair spinal co-ordination; but it is not probable that degeneration or sclerosis of a considerable length

of the posterior columns, invading other parts of the cord reluctantly, so to speak, will be due to syphilis.

So far as I know, there is still less reason to attribute atrophic paralysis to syphilis than locomotor ataxy.

Coming now to myelitis—the form most frequently seen in syphilis is subacute or chronic inflammation of a segment of the cord; but in my opinion acute general or local myelitis may be caused by syphilis, especially in the secondary stage.

This is a point which cannot easily be proved. There is nothing peculiar in acute myelitis to indicate its cause, and, the disease being rare, it must be long before a sufficient number of cases can be collected in which it is associated with syphilis, even by observers alive to the possibility, while in the secondary stage a *post mortem* diagnosis cannot be made so readily as in the tertiary period of the disease.

In 1859, I saw in quick succession two cases—one in the Hôpital du Midi under Ricord, the other under Trousseau in the Hôtel Dieu—of rapidly progressing paraplegia, terminating fatally. In both, there was general and advanced inflammation, and softening of the spinal cord. Both the patients were young, and suffering from secondary syphilis in an early stage. Ricord, with characteristic caution, refused to pronounce the myelitis syphilitic, though it was not the first case of the kind he had seen in his wards. Trousseau was more disposed to consider syphilis as the cause of the inflammation.

Of acute local myelitis, I give a case in which also the disease of the cord came on in the secondary period.

CASE I. Paraplegia, in Secondary Syphilis, Acute Myelitis of a Segment of Cord just above the Lumbar Enlargement: Death from Ulceration of Bladder and Extravasation of Urine.—A warehouseman, aged 28, was admitted under my care at St. Mary's Hospital February 23rd, 1872. His habits had been loose and irregular; he had had sores on the penis twice, eighteen months and twelve months before his admission, and had suffered from sore-throat and fissures in the palms of the hands. The inguinal glands were enlarged and matted together. It could not be made out which of the ulcers had given origin to the syphilis.

Eighteen days before he entered the hospital, he had stiffness in the right knee; next day in the left; no pain. Afterwards, pain came on in the lower limbs above and below the knees, with numbness; and, as he walked, he did not feel the feet when put to the ground, so that he tottered; a sensation also of pins and needles came in the feet. Three days after the accession of the symptoms, he was compelled to remain indoors; next day, to remain in bed; and on the day following, five days from the commencement of the illness, the sphincters were relaxed. He did not feel ill, but was thirsty. He had been taking warm-baths with mustard.

On admission, he was pale, but the expression was natural. The pupils were large. He complained only of loss of power and sensation in the lower limbs. The legs were absolutely powerless, and sensation was lost in them, except at the inner side of the thigh (obturator area), the insensibility extending on the abdomen nearly as high as the umbilicus. Reflex action was vigorous, and the legs were constantly kicking of themselves. There were incontinence of urine and feces, and small sores were seen on the sacrum and trochanters. Tenderness on pressure and percussion was found to be present over the three lower lumbar spines, but not in the skin here, or round the abdomen at the same level. Tongue moist, but coated. Pulse 90, full. Temperature 98.4 deg., both in groin and axilla, though the legs felt warmer to the hand. Urine turbid and ammoniacal.

The treatment ordered was as follows:—Iodide of potassium in doses of six grains; catheterism twice a day; protection of the abraded points by cotton-wool; a free supply of milk and beef-tea.

On the 28th, he said sensation was more distinct over the inner aspect of the thigh, and a pinch was felt at any part of the limb above the knee. The dose of iodide was increased to nine grains.

On March 2nd, the patient was feverish, and had no appetite. Temperature 101 deg.

On the 4th, he complained of sore-throat, and the tonsils and soft palate were seen to be inflamed and swollen; there was blood as well as pus in the urine, which was still ammoniacal.

On the 7th, the tonsillitis had subsided, but the temperature remained at about 100 deg., the same point as during the previous four days, and the thirst was excessive. The bladder was washed out with weak carboic acid lotion, as well as emptied. The fever continued, and on the 9th, was rather higher, the temperature being 101 deg. in the axilla, 101.5 deg. in the groin, but he said he had better sensation in the legs, and he had begun to feel the catheter for the first time. He was, however, weaker, and bark and acid were substituted for the iodide of potassium; wine also was given.

On the 10th, symptoms of perforation of the bladder set in—rigors,

sickness, severe pain in the abdomen, paroxysms of violent contraction of the abdominal muscles, and great exhaustion.

On the 11th, the face was expressive of collapse; the abdomen was tender; and the muscles were easily thrown into a state of rigid tension; there were no abdominal respiratory movements; no urine could be obtained. Death took place on the 12th.

On *post mortem* examination, fourteen hours after death, there was a remarkable amount of *post mortem* staining. In the abdomen, the effects of general peritonitis were seen; and in the bladder, a large ragged ulcer near the neck, which had penetrated the coats and allowed the urine to be extravasated. The liver was large, pale, and fatty; the kidneys healthy; the lumbar, mesenteric, and inguinal glands were large. A little old tubercle in the lungs; heart healthy. In the cord, about two and a half inches from its lower extremity, just above the lumbar enlargement, was a softened portion about half an inch in length. Above this part, the cord was firm, and the vessels were of the usual size; below it, the cord was firm, but the entire posterior aspect was covered by enlarged and tortuous vessels.

The paraplegia generally produced by syphilis is the result of disease slowly invading a segment of the spinal cord. The course of symptoms is such as might result either from a limited chronic myelitis, or from softening, or from tumour.

The age at which syphilitic paraplegia occurs, namely, in early adult life, or early middle age, excludes degenerative non-inflammatory softening. Embolic softening, so far as I know, only occurs in the cord in connection with a very general distribution of emboli, as in acute fatal chorea, or as in ulcerative endocarditis, when the effects produced elsewhere predominate. The absence of angular spinal curvature, and of the symptoms which disease of the bones or cartilages of the vertebral column produces before the cord is affected, excludes the most frequent cause of local myelitis not depending on syphilis; and we are left to decide, from the history or from collateral circumstances, between syphilis and rheumatism, or exposure to wet and cold, or injury.

In syphilis, the condition present is probably a small gumma, around which inflammatory softening is set up, and this explains the slow, halting, and unequal progress of the paralysis. Conversely, paraplegia, at first slight, remaining long at a given point, or advancing very slowly, then suddenly worse, mending a little again, perhaps spontaneously, or at least without specific treatment, but again increasing, and so pursuing its course to absolute loss of sensation and motion, while reflex action persists, should excite a suspicion of syphilitic mischief, and should lead to energetic treatment, whether corroborative evidence is forthcoming, or no.

While it was still uncertain whether syphilis could give rise to disease of the nervous system, and the purpose of inquiry was to establish a fact in medical science, no caution could be too great in accepting evidence on the point; but, the fact of causation once established, we are justified in going in advance of positive knowledge in our treatment.

[To be continued.]

THERAPEUTIC MEMORANDA.

EFFERVESCENT ASTRINGENT VOICE-LOZENGES.

CLERGYMEN, vocalists, actors, and public speakers, frequently suffer from dryness of the throat in commencing to use the voice. This dryness is very generally the result of nervousness, which arrests the secretion of saliva; and attention has already been drawn to the value of the simple effervescent lozenge of Mr. Cooper, of 26, Oxford Street, in allaying this trouble. Another cause, however, and one often found in conjunction with the foregoing, is relaxation of the uvula and mucous membrane of the pharynx; and this condition is always more or less present in the case of those who are constantly using the voice. For its relief an astringent is required; but most astringent lozenges produce dryness. The combination, however, of an astringent and effervescent saline, while bracing the mucous membrane, does not cause dryness, but, on the other hand, promotes a free secretion of saliva. Mr. Cooper has prepared some lozenges for me of this character, using the gum of eucalyptus as the astringent. By the addition of a small amount of cinnamon and clove, he has made it agreeable in taste, while it is quite free from any irritant stimulant, so frequently found in "voice-lozenges". Having prescribed these lozenges for some months with very considerable success, I am induced to think they may be found a useful addition to our remedies for some of the slighter throat-troubles for which one is constantly consulted in this climate.

LENNOX BROWN, Welbeck Street.