engaged in dissections, should be allowed to serve as a dresser in an hospital.

There is a remarkable resemblance, if not an actual similarity, between puerperal peritonitis and the erysipelatous inflammations of the peritoneum following operations on the organs contained in that cavity. The influence of contamination in the way that I mention is well illustrated by what occurred some years ago at the Vienna Hospital. Prior to 1846, the proportion of deaths amongst lying-in women in this hospital was one in ten. In order to check this frightful mortality, Dr. Semelweis required the students who attended these women to avoid making necropsies, and obliged them to wash their hands in a solution of chlorine before and after making vaginal examinations. The result was, that the mortality in 1848 fell to one in seventy-four. (Braidwood *On Pyæmia*, page 220.)

That erysipelas is often of epidemic origin, there can be no question; but the influence of any epidemic is immensely increased by an unhealthy condition of a ward from overcrowding. Of this I will give you an instance.

At the middle of November, 1872, ward I was filled by a very severe set of cases. There was one patient who had had both legs amputated below the knees; another whose leg had been amputated through the knee; another who had a compound fracture into the ankle-joint; one with extensive necrosis of the tibia, and another with extensive laceration of the fore-arm. And this condition of severe cases was kept up by two other primary amputations—one of the foot, the other of the thigh—these cases being admitted in December, but whilst the preceding ones were under treatment. There was no erysipelas in the ward until November 15th. On that day, the case of compound fracture into the ankle was attacked. He recovered, and the erysipelas had left him by the 21st. On this day, the patient with necrosis of the tibia had the sequestrum removed. He was a feeble old man. He was attacked by erysipelas on December 1st, was removed to the erysipelas ward, and died there on the 7th. On December 4th, the patient with scalp-wound was attacked, and was at once removed to the erysipelas ward. During this period, viz., on December 7th and on the 22nd, two of the other patients—the one with amputation through the knee-joint, and another—were seized with pyæmia. The patient who had been amputated through the knee, a healthy gardener, was the only one out of the twenty-three patients with acute traumatic pyæmia who recovered. During the same period, two of my patients in ward 5, both suffering from chronic wounds—one a fistula *in ano*, the other a sinus in the thigh—were attacked by erysipelas. To them, it is possible that the infection was conveyed by the house-surgeon or dressers; but, as other cases of erysipelas occurred in the hospital at the same time, and one was admitted from out of doors, it is possible that an epidemic influence existed.

In the next and concluding lecture, we will consider the means of prevention of septic disease, and of lessening the mortality in hospitals.

ERRATUM.—In Lecture II, page 97, column 2, line 38, after "general hospital practice," insert "in the adult."

CASE OF *POST PARTUM* HÆMORRHAGE TREATED BY THE INJECTION OF PERCHLORIDE OF IRON.

At the present time, when the intrauterine injection of the perchloride of iron in *post partum* hæmorrhage is on its trial, it is well that each individual case of its failure or success should be recorded.

In November 1873, I was called to the lying-in ward of this hospital to see a woman suffering from *post partum* hæmorrhage after the birth of her first child. The woman was very pale, but perfectly calm and collected. The uterus was found to be above the umbilicus, in a very soft and flaccid state. By the use of manual pressure, accompanied with cold application to the abdomen, the womb contracted, expelling a large quantity of blood and clots. This contraction continued for about five minutes, when it ceased, and the uterus was in the state in which I found it. The hand was now introduced, and, with the means I had previously used, contraction was the result, but to be followed in a much shorter period by relaxation. The woman's position was now serious; her pulse was quick, weak, and intermittent; her respiration was a series of sighs; she was very much agitated, and had great desire to be left alone. The injection of the perchloride of iron was now determined on. Two ounces of the strong solution were used in half a pint of iced water. The hæmorrhage immediately stopped; the uterus became corrugated on its internal surface, but remained uncontracted. Stimulants with a sedative were now administered. From this time, she progressed favourably, and was discharged at the usual period.

A. MACLEOD HAMILTON, M.D.,

Resident Medical Officer, Parish Infirmary, Liverpool.

LETTSONIAN LECTURES ON SYPHILIS AS A CAUSE OF DISEASE OF THE NERVOUS SYSTEM.

Delivered before the Medical Society of London.

By W. H. BROADBENT, M.D., F.R.C.P.,

Physician and Lecturer on Medicine at St. Mary's Hospital.

LECTURE II.—*Continued.*

CASE II. *Paraplegia, probably from a Syphilitic Nodule in the Middle Dorsal Region of the Spinal Cord.*—Ann F., aged 41, a widow, was under my care in St. Mary's Hospital from July 24th to December 2nd, 1872.

She had been married eleven years, and had had two early miscarriages, but no living children. There was no history of syphilitic symptoms, but one iris was found bound down by adhesions. Her husband had died early in 1871 from an illness, which I venture to pronounce syphilitic affection of the brain. In June, 1870, while in apparent health, he had an attack of convulsions, which was attributed to sun-stroke, though he was at work under cover. After this attack, he was never himself; he had frequent fits, and his mind became affected; at times also he lost the use of his limbs. He died in an asylum nine months after the first attack.

The patient had for three months been losing the use of her legs. First, the right failed; then this seemed to recover somewhat, and the left became affected. The weakness increased till she could scarcely move about; the bowels were confined, and the urine was not passed readily. On testing the power of the limbs as she lay in bed, the right leg was fairly strong, but the flexors at the back of the left thigh were almost powerless; the extensors of the knee still retained a good degree of strength. I may note in passing that, in partial paraplegia, the flexors are usually affected earlier and more severely than the extensors of the knee. The ankle was fixed and immovable, the foot arched, and the great toe turned in. There was no loss of sensation anywhere. When the patient tried to walk, the difficulty was greater than the degree of paralysis led me to expect; she could scarcely walk alone, and was very awkward and clumsy; the left foot shot out uncontrollably. She could stand as steadily with the eyes closed as when they were open.

Iodide of potassium was given in doses of four grains, and the continuous galvanic current was ordered. The case appeared to be slight, and I hoped for much from rest in bed and treatment. She had, however, come under observation at a moment when the disease was in active progress, and a few days later the legs were almost absolutely powerless, and sensation was lost, this aggravation of the symptoms having come on rather rapidly. She had fallen helplessly in trying to get out of bed, when unaware of her increased weakness.

I found her, on August 7th, quite unable to move either leg; the limbs, however, were not flaccid and soft, but, when handled and tested, they were firm and rather stiff from involuntary resistance of the muscles, and it was ascertained on inquiry that, before she had so completely lost power in them, they became rigid when the feet touched the ground. Reflex action took place when the soles were tickled, but it was not vigorous. There was violent involuntary starting of the lower extremities, both during asleep and while awake. The difficulty in sitting up was peculiar. As soon as she made the attempt, aiding herself with her hands, the legs contracted and resisted, and it was only after a severe effort that she succeeded, the lower extremities jerking all the time.

Sensation, which was severely tested, was almost totally lost on the right side as high as the lower edge of the ribs; on the left, up to the level of the mamma; but there was an imperfect kind of consciousness of the contact with the soles of the feet when they were tickled. The bowels were still confined; there was less complaint of difficulty in passing urine, because of the loss of sensation; but the urine was retained, and the catheter was required.

The dose of iodide of potassium was increased to twelve grains—at first, three times a day, then every four hours. The galvanism was discontinued. In the course of a few days, sensation began to return—first, in the left side, in which the anæsthesia had extended highest, and the recovery was general in the extremity and abdomen, and not progressive; while, on the right side, the return was at first progressive from above downwards. It was further noted that, while the difference in the degree of sensation on the two sides existed, reflex action was more vigorous in the limb in which sensation was most impaired.

On August 13th, she had more control over the evacuations, and she thought there was some return of sensation and power in the legs. She was still much troubled by the contraction of the legs at night, which was accompanied by cramp-like pains, and disturbed her sleep.

August 17th.—She had less starting of the limbs during sleep, and the feet could be distinctly moved. Sensation was returning. The difficulty in sitting up in bed was as great as ever. Immediately on the attempt being made, the muscles of the lower extremities contracted, drawing up the knees, which had to be forcibly pressed down before she could raise the body. This was an interesting phenomenon in relation to the functional mechanism of the spinal cord, and it contributed to the formation of a diagnosis. The urine was now passed naturally. The improvement continued up to August 31st. The startings had ceased, the toes and legs could be moved, and sensation was returning. Cod-liver oil and iron were now ordered, and, by mistake, the iodide was suspended. A general relapse occurred; sensation and power were again completely lost, and the painful nocturnal startings returned. I was out of town at the time, and a week elapsed before it was known that she was not taking the iodide of potassium. It was ordered in doses of fifteen grains three times a day on September 17th, and blisters were applied; after which, in a few days, amendment set in. I found her, on September 25th, a little better than when I left town at the end of August. The iodide of potassium was increased to twenty-four grains, and one-sixteenth of a grain of biniodide of mercury was given with quinine and extract of hop, in a pill, at night.

From this time, the improvement was continuous. A curious fact observed for some time, while sensation was still imperfect, was, that a sharp pain could be excited in the left side by pricking a particular part of the leg of the same side near the knee.

On October 12th, faradisation was ordered. On October 23rd, the patient was able to stand by supporting herself with a chair; on November 1st, to walk with assistance; on the 12th, for a few steps, alone. She was kept in the hospital till she could walk well, and she continued for some time after she left, on December 2nd, to take the iodide in doses which, on February 15th, had been gradually brought down to twelve grains three times a day. She was then walking without perceptible halt, and with great vigour, and, so far as I know, she continues well.

CASE III. *Paraplegia from Syphilitic Nodule or Local Myelitis: Recovery.*—Another case, strikingly similar, even in details, was under my care from March 28th to August 9th, 1873. The patient, a domestic servant, aged 29, had been suffering for four months from loss of power and sensation in the lower extremities, which had come on with pain in the back. She was unmarried, and I could obtain no evidence of syphilis, direct or indirect; but she had the peculiar sallow pallor which often accompanies the latter stages of the disease. I took this as sufficient warrant for the treatment I adopted; but I should not have considered myself justified in bringing the case before you as one of syphilitic disease of the spinal cord, had it not been that, after the patient's recovery from the paralysis, when suffering from a severe attack of pyrexia, she confessed to the sister of the ward that, many years before, she had had an illegitimate child, and subsequently had some illness, attributed to syphilis.

The pain in the back experienced at the outset of the illness was situated in the lower dorsal and upper lumbar region; it extended round the abdomen and down the legs, especially when she walked or stood. It lasted about a fortnight, and then the loss of power which had been coming on became suddenly worse, so that she was unable to rise from a chair, or to raise the limbs; sensation also was greatly impaired, and the feet and legs were very cold. When examined, a few days after her admission, she was unable to move either leg; sensation was greatly impaired in both extremities and over the lower part of the abdomen, but more on the right than on the left side. Reflex action was exaggerated, and the limbs jumped very much, especially during sleep. The catheter was required, and the bowels could only be relieved by means of enemata.

The treatment consisted in the administration of iodide of potassium in gradually increasing doses, and the result was complete recovery. The catamenia were absent till near the end of her stay in hospital; and at one time, early in July, a profuse purulent vaginal discharge came on. She was examined by Dr. Meadows, who found no abnormal condition of the uterus. Convalescence was also, for a few days, interrupted by a sharp attack of pyrexia, not accounted for by any local inflammation. During this, the temperature, habitually slightly below the normal standard, went up to 103.6 degs. Fahr. She walked out of the hospital on August 9th.

CASE IV. *Paraplegia; Meningo-Myelitis of Lower Extremity of Cord: Recovery.*—The patient, a groom, aged 43, came under my care on October 3rd, 1873, complaining of difficulty in holding his urine, and

weakness of the legs, with pain in the back. He had begun to suffer from intermittent pain in the back after a fall eight or nine months previously, which was increased or brought on by walking; he lately had lost power in the legs, and had been unable to retain his urine. He had had syphilis twenty years before.

It was found that the right leg was weaker than the left, the movements of this ankle being very imperfect, and flexion of the knee very feeble. The tests applied to ascertain the strength of the muscles gave pain. There was apparently cutaneous hyperæsthesia of the right leg. Reflex action was readily excited. He maintained the erect position with the eyes closed. A little pain was caused by percussion or pressure upon the last two lumbar vertebrae. There was at first improvement; less pain, more strength in the legs, and increasing control over the urine.

On October 17th, however, he felt cold and shivered; had great pain in both legs, especially at and below the knees, increased by movements. He was unable to walk. Pulse, 120; the temperature, previously rather below the normal, was now 100 degs. The pain became so severe, especially at night, that morphia injections were necessary, and the catheter was required on account of retention of urine. Later, the principal seat of pain was the left thigh, at first near the exit of the great sciatic nerve, but afterwards on the dorsum ilii and in the femur, in both of which situations it was found to be due to periostitis. The loss of power in the lower limbs did not increase, but the bladder continued to give trouble, and an abscess formed in or near the prostate, which, for a time, added to the patient's sufferings.

The treatment consisted in the administration of iodide of potassium and biniodide of mercury, together with the application of blisters over the seat of tenderness in the spine; and ultimately the patient recovered almost complete power in the legs and fair control over the urine.

CASE V. *Sensory Paraplegia; Loss of Control, but not of Power; Exaggerated Reflex Action.*—The following case is one of unusual interest, there being great impairment of sensation and exaggerated reflex action, with loss of control, but not of power. It might easily be taken for locomotor ataxy, but differs essentially from this affection in its symptoms, and, as I believe, in its pathology.

The patient, a cabman, aged 34, married, but childless, was admitted into St. Mary's Hospital, March 7th, 1873, for loss of power in the legs. He was a strong, stout, well-built man, with good colour, and large and firm muscles. He had, however, lost all his hair, not only from the scalp, but from the face, axillæ, and pubes; the eye-brows and eye-lashes also were gone. He had suffered from gonorrhœa ten years before, but had never had a sore on the penis; he had, however, had "measles" and a bad sore-throat fourteen years previously, which had lasted three months, and ample evidence of syphilis was afforded by large ulcerations of unmistakable character on both legs. The weakness of the lower extremities had been coming on for twelve months. He began to drag first the left foot, and then the right, and the knees and feet gradually became "numb." At times, they would be a little better after being in bed, or after sitting on his cab; they were worse in warm weather. He had been under the care of my colleague, Dr. Lawson, as out-patient for two months before his admission, and had been faradised with apparent advantage. For three weeks, however, he had been unable to walk.

On first seeing him, I told him to rise, so that I might have the opportunity of noting the degree of paralysis, and observing its character. He threw his legs off the bed without difficulty; but, when the feet were put to the ground and he endeavoured to stand, the attempt was completely defeated by violent spasmodic contraction of the legs, especially the right, which would have thrown him down, had he not been caught and assisted back to bed; the movements continued for some time afterwards. He explained that he could stand and walk better with his stockings on than when the feet were bare, and still better in strong boots. The difficulty obviously arose from excessive reflex action, which was more violent when the feet were in direct contact with the floor. Lying in bed, he could move every part of both lower extremities, and could resist forcibly the attempt to flex or extend any joint. The muscles were bulky and firm, and particularly well-nourished. He occasionally had startings of the limbs at night. Sensation was greatly impaired in every part of the thighs and legs, whether tested by compasses or by pricking and pinching. The insensibility to heat was shown by the fact that he had large blisters on the soles of both feet, caused by a hot bottle which he had not felt. He had, for three or four months, had difficulty in micturition; he had to wait some time before the flow began, and the urine came only in drops; it contained no albumen. Pulse and temperature normal.

To complete the account of the symptoms, it need only be added that, when he was able to stand and walk, he could do so equally well with the eyes shut or open, and he had no particular difficulty in turning.

He stood steadily with the eyes closed, when he could only walk with the aid of two sticks. In walking, the legs were not shot out or jerked up vaguely, as in locomotor ataxy, but were carried along stiffly and wide apart, the feet scraping the floor in consequence of want of flexibility in all the joints, which were kept rigid by tonic contraction of the muscles generally. The hand placed on the thigh or calf when he walked found them hard and firm, like those of an athlete.

The phenomena of this case are worthy of study and discussion. They were clearly not those of the condition known as locomotor ataxy, although they would be covered by the term. The mode of progression, the power of standing with closed eyes, the absence of pains in the limbs, exclude at once sclerosis of the posterior columns of the cord, which is the morbid change giving rise to locomotor ataxy; and there are other differences of equal significance. I cannot fully convey to you the precise grounds on which I came to the conclusion at which I have arrived, without reference to the functional mechanism of the spinal cord, as I understand it. The anterior grey columns consist demonstrably of the nuclei of the anterior or motor roots of the nerves fused together; the posterior grey columns in the same way of the nuclei of the posterior nerve-roots (this is inferred rather than distinctly proved). The antero-lateral white columns are the downward channel for motor impulses from the corpus striatum to the motor nerve nuclei; the central grey matter is the channel for sensory impressions upwards from the posterior nerve-nuclei to the thalamus; decussation of the fibres or cells constituting the sensory channel taking place one by one in the cord. Now, Brown-Séquard has shown that injury to the central grey matter will intercept sensory impressions without interfering with motor power, and this is exactly what had occurred in the case under consideration, and I concluded that a syphilitic tumour existed in the very axis of the cord. But the posterior and anterior nerve-nuclei together constitute the apparatus for reflex action, and there is in the cord an independent power of co-ordination, that is, of orderly combination of muscular movements for a given purpose, the mechanism for which consists in the linking together of the nerve-nuclei of different parts of the cord. It would have been difficult to say *à priori* whether such co-ordinating fibres would connect together anterior or posterior nerve-nuclei; but it has been found experimentally that it is through the posterior nuclei that co-ordination is effected, the fibres which effect it running in the posterior white columns. I am of opinion, however, that there is also an association of the anterior nerve-nuclei of different segments, as I find no other explanation of the peculiar arrangement of the fibres of the anterior white columns; and I think it serves some such purpose as the combination of the action of one fore-leg with the hind-leg of the opposite side in dogs, such association explaining the fact that injury to the corpus striatum does not occasion hemiplegia in dogs. Let me now, however compare this case, in which I suppose a segment of the central grey matter to be damaged, with locomotor ataxy, in which the posterior white columns are affected. In both we have impairment of sensation, with comparatively little loss of motor power, but with loss of control. In locomotor ataxy there is destruction of the fibres in the posterior white columns, which convey the impressions from one part of the cord to another, and bring about the delicate adjusting movements which insensibly maintain the balance of the body, and the patient is unable to stand with the eyes closed. My patient stood perfectly well. The loss of balancing power, then, in locomotor ataxy is not simply due to impaired sensation in the feet, as has been maintained by Dr. Clifford Allbutt. Again, in locomotor ataxy, motor impulses from the cerebrum for the act of walking arrive at certain nuclei in the lumbar swelling of the cord; in consequence of the disease in the posterior columns, these impulses are not diffused along them to the different segments of the cord which co-operate in the production of the harmonious movements of the limbs, but are concentrated upon the nuclei on which they impinge, throwing them into excessive and disorderly action, and causing the jerky gait. This was not the case with my patient, but, on the contrary, the impressions received by the feet in contact with the floor, instead of being partly transmitted to the brain for the communication of sensation, were, in consequence of the interruption in the upward channel, concentrated upon the cord, but distributed to different segments by the posterior columns, producing the excessive but co-ordinated reflex contraction of the muscles of the limbs which restrained or restricted their obedience to the will.

A small tumour, such as was indicated by the symptoms to be present in the axis of the cord, was just the kind of change to which syphilis might give rise, and which scarcely any other disease would produce. This is one of the examples, then, in which the symptoms alone would almost warrant a diagnosis of syphilis without any corroborative evidence.

The progress of the case was satisfactory. I began the administra-

tion of iodide of potassium in doses of ten grains, with carbonate of ammonia and sarsaparilla, carrying it eventually to twenty-five grains three times a day. In a few days, the nocturnal startings were less troublesome, then sensation began to return, and he passed urine better. He walked with the aid of two sticks on April 3rd, and on the 26th with only one, passing urine now quite readily; sensation also was much better, but not normal; the ulcers on the legs had healed. About this time faradisation was ordered.

He left the hospital on May 2nd, after a stay of less than two months. On the 27th, he drove a hansom to the Derby. He continued to take the iodide (fifteen grains) till July 24th, when tincture of iron with nux vomica was given instead. After a time, he gradually lost power, and, on September 3rd, had gone back considerably. The ulcers on the legs had also broken out again badly, and were extending. Iodide of potassium was therefore prescribed anew, and very shortly improvement again set in, which continues up to the present date. He is still faradised twice a week, as he thinks this does him good, and it aids the keeping him under observation. The relapse and improvement occurred independently of any change in this part of the treatment. He is now, against my advice, driving a four-wheeled cab, and has to assist in lifting and carrying luggage.

I was not so successful in a somewhat similar case which was under my care for four months, shortly before the case just related, namely, from September 2nd, 1872, to January 6th, 1873.

CASE VI.—The patient was a coachman, aged 30, who had had syphilis twelve months before he came to the hospital, and for seven months had been suffering from loss of power in the limbs, which had come on with pain in the loins. This at first had been induced by exertion, but had become constant, and the weakness was such that the patient could only walk by the aid of two sticks; his urine dribbled, and the bowels would be confined for three or four days, and then act suddenly without his control. His manner of walking was exactly that described in the previous case—the legs stiff and rather straddling, with all the muscles extremely firm; the joints unbending, the feet therefore scraping the ground. After walking, there was frequently most violent agitation of the legs, but especially of the right, and he would sit on his bed trying in vain to control the movements with both hands grasping the thigh. The patient had difficulty in standing with the feet together, tending to fall backwards, and, when the eyes were closed, he was still more unsteady on his legs.

In testing the muscles as he lay in bed, it was seen that he could execute any movement, and the limbs were remarkably well-nourished. He could resist flexion and extension of the ankle, but the muscles trembled; he resisted flexion of the knees powerfully, extension scarcely at all, the left leg being the stronger.

The treatment was by iodide of potassium, and, this failing to give marked results when carried to full doses, biniodide of mercury was given; blisters also were applied. He regained complete control over the bladder and rectum, and walked a little better, but needed two sticks, and the effort was usually followed by the violent shaking of the legs described; the amendment, therefore, was not sufficient to enable the patient to gain his livelihood.

Spinal meningitis is far less common than myelitis, and I have met with no acute case distinctly traceable to syphilis. A patient, however, was under my care in September last, suffering from severe pain in the back and limbs, made worse by movements, with feverishness, for which the cause assigned was a fall, in which the back was injured. The fall, however, had happened a month before, and he had since worked at his trade without pain or inconvenience. An explanation of the pain soon appeared in an outbreak of syphilitic eruption, and it was removed by a mercurial course. I have now (January 15th) in hospital a patient recovering from a subacute attack, who had previously suffered from paraplegia, and whose case I have already given. She left the hospital well on August 9th, and remained free from symptoms till when she began to suffer from rheumatism, as she supposed. There was, however, no swelling of the joints, or fever, and the pain was provoked by movements of the limbs or spine, and had the characters of pain caused by spinal meningitis. She began also to lose power again in the lower extremities. The pains were at once relieved by iodide of potassium.

One of the worst cases of syphilitic affection of the nervous system which I have seen, and which ultimately resulted in insanity, began with spinal meningitis. The patient was under the care of Mr. Walter Coulson, with whom I saw him, and the case is partly related in his treatise on syphilis. When I first saw him, in August 1866, he was complaining of rheumatism—obviously spinal meningitis—which completely crippled him. Every movement gave him pain, but especially the first efforts, after being for a time in one position; so that it was distressing to watch him rise from a chair. He carried himself stiffly,

was unable to stoop or to turn the head; in order to look round, he had to turn the whole body. There had previously been tertiary manifestations, and the meningitis yielded to iodide of potassium, leaving the patient, however, with a stoop; to it also was probably attributable sexual impotence, from which he was found to suffer afterwards.

[To be continued.]

SPINA BIFIDA CURED BY INJECTION.

By J. ROSS WATT, M.B.,

Consulting Surgeon to the Fever Hospital, Ayr, N.B.

THIS wood-cut, from a photograph, represents E. G., aged three years, as she was brought to me in April last, and therefore conveys a correct idea of the size and situation of a spina bifida tumour, which, wholly

covered by true skin, stood out with a prominence of nearly four inches, and measured thirteen and a half in circumference. It was very sensitive to the touch, became tense when the child cried, and palpation discovered a perfectly fluid condition of its contents. It was slightly pendulous when the child was erect, and gave her a ludicrous, although at present rather fashionable, appearance when dressed—the projection at her age being unlooked for.

Her mother stated that it was like a walnut at birth, and had increased gradually; that she walked at two years; was now active, although not firm on her legs, and easily knocked over; that she enjoyed excellent health, but was constantly annoyed by escapes from the bladder and rectum; and that she had always been advised to let it alone.

The treatment of this—my second successful case—was by the same method as that described in the JOURNAL of the 26th April last; viz., by injection of Dr. James Morton's solution of iodine and iodide of potassium in glycerine; and was briefly this. Two tentative tappings with a medium cannula, of respectively twelve and ten ounces of clear fluid, were made, with an interval of eight days, both being followed by an irritable and feverish condition of the child, and gradual refilling of the tumour to its original size. Ten days afterwards, other ten fluid-ounces were removed, and half a drachm of the solution injected. The opening was instantly closed on each occasion by a layer of lint soaked with flexile collodion. This last operation in a few days made the tumour tender, and caused the child to be feverish, restless, and extremely excitable, with decrease of appetite, milk and soups alone being taken. Ten days after this first injection, the tumour was very slightly less, and eight ounces were then removed, followed by injection of a similar quantity of the solution. This gave rise to even more severe irritative symptoms of the nature described; but a fortnight later showed the tumour only one-third of its original size, although still sensitive and painful when interfered with. At that time, about five ounces were removed, followed by a third injection of half a drachm; this caused cessation of any further formation of fluid, and a gradual absorption began, which, at the end of twelve weeks from the first interference, presented the appearance conveyed by the second wood-cut—a roughened, darkened, hardened, and thickened condition of skin, quite normally sensitive to the prick of a needle, replacing the tumour and closing the spinal aperture with an apparently gelatinous mass.

The treatment extended over nearly seven weeks, and the measured fluid removed amounted to forty-six ounces—a larger quantity, to my knowledge, than from any previously successful case. Half a drachm of the solution was deemed sufficient for each injection, owing to the large extent of sensitive surface to be dealt with; and the irritative symptoms developing therefrom quite justified the precaution taken.

This, my second case, with Dr. Morton's, make four successes, all that have been so treated; and, without appearing too confident of further success, the result is very hopeful and gratifying to the introducer of this method of treatment.

The patient is now very active on her feet (amusing herself, and standing as much consequent fatigue daily as her healthy companions), growing rapidly, and the sphincters are now almost wholly under control; cold sea-water baths being persevered with.*

THE PROPAGATION OF CHOLERA IN INDIA:

BEING A REPLY TO DR. MURRAY'S REVIEW OF DR. CUNINGHAM'S REPORT OF THE EPIDEMIC OF CHOLERA IN 1872.†

By J. M. CUNINGHAM, M.D.,

Sanitary Commissioner with the Government in India.‡

DR. MURRAY very kindly sent me his paper yesterday, so that I might be prepared to answer him at the meeting to-night; but I regret that, owing to a press of business during a short visit to England, I have not been able to do more than make a few rough notes on the subject. I shall first take up, as shortly as possible, the various points to which Dr. Murray has referred, and then discuss others of very great importance which he has left out. Before doing this, however, I may make one remark regarding the Report itself.

The great object I had in view was not to promulgate any opinions of my own, but to collect facts and to draw from them the practical conclusions as to the best mode of dealing with cholera which these facts warrant. The collected facts are the really important part of the Report. To the best of my belief, no fact of any importance has been misstated, nor have any which were ascertained been omitted. The Report has been widely circulated among medical officers in India, and no exception has been taken to the record. We pass on, then, to the interpretation to be put on the facts; and here Dr. Murray urges objections.

He first raises the question as to the comparative value of negative in contradistinction to what he calls positive evidence. I am glad he has raised this question, for the proper solution of it is a matter of very great moment in the cholera inquiry. What constitutes positive evidence? If a person, after having come into contact with a case of cholera, be affected with the disease, he claims this fact as positive evidence that cholera is contagious. But such reasoning is altogether fallacious. The mere coincidence of two events by no means shows that they stand to one another in the relation of cause and effect. The

* This case was shown at a meeting of the Ayrshire Medical Association.

† See BRITISH MEDICAL JOURNAL, January 17th, p. 71.

‡ Read before the Epidemiological Society of London.