LETT SOMIAN LECTURES

ON

SYPHILIS AS A CAUSE OF DISEASE OF THE NERVOUS SYSTEM.

Delivered before the Medical Society of London.

By W. H. BROADBENT, M.D., F.R.C.P.,
Physician and Lecturer on Medicine at St. Mary's Hospital.

LECTURE II.—*Concluded.*

Syphilitic Affections of the Medulla Oblongata and Pons.—The intracranial prolongation of the spinal cord forming the medulla oblongata and pons Varolii is complicated in its structure and function, not only by the separation of the nerve-nuclei from each other, in accordance with the specialisation of their functions, but by the change in the relations of the grey matter and tracts of white fibres with each other, the decussation of the motor tracts, and the connection of the cerebellum with the spinal axis which is here effected. The consequences of lesions here are, therefore, more varied, and the causes which may give rise to them are increased in number. These parts, for example, may be affected by disease in the cerebellum, or by aneurism of the basilar artery.

The increased dimensions of the spinal axis, moreover, permit unilateral limitation of a morbid change; and this is not uncommon, although the two halves are fused together, while unilateral mischief is rarely seen in the cord, which is almost completely divided longitudinally by the fissures. Possibly, however, the vascular membrane which dips into the fissures is more efficient in communicating a morbid change than the nervous structures.

The general results of disease in the medulla and pons are some interference with the sensory or motor tract, proceeding from the cord to the sensori-motor ganglia, together with disturbance or abolition of the function of one or more of the nerve-nuclei. It is seldom that we can come to the conclusion that there is disease here from sensory or motor paralysis of the limbs alone, or from affections of the nerves which have their origin in the intracranial part of the spinal axis alone, although this may be done occasionally when two nerves are simultaneously paralysed, the nuclei of which are conjoined or in close proximity, while the nerves take a different course after their emergence from the centre, as in the case of the sixth and seventh. The grouping of symptoms arising from the inclusion in a diseased area of several nerve-nuclei may be very varied; usually, the symptoms afford the means of making a very precise localisation of the lesion by the application of anatomical and physiological knowledge. Of course, disease here is attended with great danger to life, the slightest interference with the reflex centres of the respiratory and cardiac movements being fatal. Sudden death is, therefore, common. The nature of the morbid change will be arrived at by considerations such as those made use of in determining the probable cause of paraplegia. A slow, but irregular, progress of the affection will favour the hypothesis of syphilis; but the morbid changes resulting from syphilis do not specially affect the surface of the medulla and pons, according to Dr. Moxon's dictum, but rather the substance.

The progressive labio-glosso-laryngeal paralysis, which is the counterpart of locomotor ataxy, is not syphilitic; but it is quite possible for syphilitic change to involve the same parts of the medulla, and give rise to analogous or even identical symptoms. The course of the disease will, however, be different.

CASE I. Paralysis (Hemiplegic) of Left Face; Lateral Deviation of Eyes to Right; Impairment of Articulation and Deglutition; Loss of Sensation in Right Face; Sudden Death; Syphilitic Tumours in Pons and Medulla Oblongata.—William T., aged 46, was admitted into St. Mary's Hospital, under my care, on August 4th, 1871. He had been a soldier, and had served twenty years in India, enjoying while there good health; but, since his return, four years ago, he had suffered from his liver. He had syphilis in India. On his fore-arms and legs were numerous large scars, as of rupia; his complexion also was sallow, and his appearance that of syphilitic cachexia. On June 20th, while cutting grass, at 10.30 A.M., the sun being rather powerful, he said the sun struck the left side of his face; he felt queer, did not fall, but had to go indoors, and he was never able to work afterwards. He suffered from pain in the head, giddiness, double vision, and a feeling of cold-

ness down the right side. There was no vomiting. The face was drawn at the time, but had become worse since.

On admission, there was marked loss of power in the left side of the face; the wrinkles were smoothed out; the mouth was carried to the right; the distortion was always evident, but was worse when he opened the mouth, or tried to show the upper teeth. Both eyes were closed in the usual winking movements; he could not close the left eye alone. The uvula was drawn to the right, and this side of the palatine arch was lower than the left. He complained of difficulty in masticating his food. This was referred to the right side of the mouth. (No note of the condition of the masseters and temporals was made.)

Deglutition was embarrassed. The tongue was protruded straight, but his articulation was thick and indistinct, and the voice had a very nasal twang. There was no difficulty in the articulation of any particular letter as the alphabet was gone through. He had double vision, especially for distant objects, and a slight want of parallelism between the two eyes was evident. (In two different observations, this was described as internal strabismus of the left eye, and that the right was carried too far outwards.) A more noteworthy fact, however, was that there was lateral deviation of both eyes to the right. He could not bring the left outwards at all beyond the median plane of the orbit; the right he could bring slightly to the inner side of the median plane, but very little.

He said the right hand was dead, but there was no perceptible difference in power between the two hands when they were tested. He used the two legs equally well when in bed, but, when made to walk, he complained of giddiness, walked slowly, feebly, and unsteadily, staggering mostly to the right. He complained that, when shaving, the right side of the face felt like a board, while the left, in which the motor paralysis was present, felt natural; and it was found that a prick or pinch was much less felt on the right than on the left forehead, cheek, and chin. He said also that he felt less distinctly on the right than the left fore-arm; but the difference, if real, was very slight. Smell and taste were unimpaired, as also common sensation in the tongue; he had been a little deaf in the left ear before the attack. He had continuous severe pain in the head; sometimes all over, sometimes in the occipital and frontal regions, sometimes in the right half. He had a singing noise also in the head; he dozed rather than slept. The ophthalmoscopic signs were negative. There was no albuminuria; no heart-disease. The diagnosis given was syphilitic tumour in the pons and medulla.

He remained in the hospital nearly three weeks, taking iodide of potassium in doses gradually increased from six to sixteen grains. The headache was greatly relieved; he walked better and with less staggering, and his articulation was better; but the motor paralysis on the left side of the face, and the loss of sensation over the area of the right fifth, remained, as did also the lateral deflection of the eyes.

He insisted on leaving the hospital on August 23rd. During the night of August 30th, he got out of bed to smoke, and was found dead in his chair in the morning. The body was taken to the Marylebone Infirmary, where an inquest was held, and I unfortunately did not hear of his death till the examination had been made. I had the head re-opened, however, and carefully reinspected the brain. It was reported to me that the dura mater and arachnoid over the top of the hemispheres were adherent; the dura mater unduly vascular; the arachnoid opaque over the hemispheres, but not at the base. There was fluid in the ventricles, the parietes of which were soft, but apparently only by *post mortem* change. The only other morbid appearance found was that, when an incision was made along the median line of the pons and medulla, two small tumours were seen near the floor of the fourth ventricle. They were described as yellow in colour, like a corpus luteum, and of firm consistence, and they slipped easily out of the cavities in which they were contained. I did not see the tumours, but, in my re-examination of the parts, I found one cavity which would have held a pea in the substance of the pons, near the floor of the fourth ventricle, the lower margin of the cavity being well within the lower edge of the pons—the upper not quite reaching the centre of the pons—it was near the median line. Below this, and separated from it by nervous substance, was another somewhat larger cavity in the substance of the medulla, also near the floor of the ventricle. This cavity had not such well-defined walls as the other. A careful examination of both crura and of the hemispheres failed to reveal any other lesion.

There can be no doubt that the lower of these tumours was the cause of the dysphagia and of the sudden death, as it was in close proximity with, and probably involved in a greater or less degree, the reflex centres for deglutition, respiration, and the cardiac movements, the nuclei of the glosso-pharyngeal and pneumogastric nerves. It is unfortunate that I am not able to speak more positively with regard to the absence of other lesions, so as to be in a position to discuss more confidently the

relation of the upper tumour to the motor paralysis of the left face and of the eyes, and to the loss of sensation in the area of the right trigeminus. If the paralysis of the face and eyes had been due to lesion in the facial nucleus or nerve, and in the sixth nucleus or nerve of the same side, there would have been no difficulty in recognising one or other of the tumours as the probable cause, by involving the common nucleus of the sixth and facial nerves. But both the facial and the ocular paralysis were such as are met with in hemiplegia, and not such as are seen in Bell's paralysis, on the one hand, and in lesion of the sixth nerve, on the other; the action of the left orbicularis oculi was not affected, and there was lateral deviation of both eyes to the right, and not simply internal strabismus of the left. The lesion, then, was either in the right corpus striatum or in the course of the fibres passing from the left facial and sixth nerve-nuclei to the right corpus striatum, and almost certainly after their crossing to the right half of the pons.

I have not previously met with lateral deflection of the two eyes, except in hemiplegia, and usually in severe attacks. In my own experience, it has commonly been associated with lesion of the anterior part of the corpus striatum externum, and accordingly I examined most carefully these ganglia and the crura cerebri, but without finding any evidence of disease. I am inclined to consider the upper of the two tumours as the cause of the anæsthesia in the right side of the face, by encroaching on the nucleus or fibres of origin of the right trigeminus; and of the left facial and ocular paralysis, by cutting across the fibres between the common nucleus of these nerves and the right corpus striatum just as these fibres crossed the median line. I am the more disposed to take this view, on account of the persistence of the lateral deviation of the eyes, this being usually a fugitive condition when it is due to lesion in the corpus striatum, the reason for which I have explained elsewhere.

CASE II. Slight Right Hemiplegia of Gradual Access; Impaired Articulation; Peculiar Affection of Respiration; Sudden Death.—Ross C., a sweep, aged 39, became an out-patient at St. Mary's Hospital January 30th, 1865. He had gradually been losing power in the right limbs for a year, and the leg, hand, face, and tongue trembled. His speech was slow, and his utterance thick. The right pupil was smaller than the left. He was "very heavy for sleep," and would often fall asleep in his chair, and, while sleeping, his wife said he turned blue, and at length woke up suddenly, appearing bewildered, and often laughing. The articulatory defect was greater than was warranted by the degree of paralysis of the limbs, and was of a kind which suggested lesion of the medulla as the cause. The partial asphyxia which occurred during sleep also pointed to the medulla as the seat of some disease which made the reflex respiratory movements insufficient, and rendered supplemental voluntary respiration necessary. He had had syphilis, and iodide of potassium, in eight-grain doses, was given three times a day, and a Plummer's pill at night. He improved considerably, his speech became less thick, he regained almost full power in his right limbs, and was able to resume work. On March 9th, he paid his last visit to the hospital, and was remarked by his wife to be in very good spirits on account of his improvement. In the course of the day, he took beer to excess with a friend, in celebration of his recovery, went home, lay down on the bed, fell asleep, and was found dead.

I heard nothing of this till three or four weeks later. A *post mortem* examination had been made and an inquest held, the cause of death assigned being heart-disease. I afterwards saw the medical man who had made the examination, and found that the opinion given was an inference from the mode of death, and was not based on definite change in the heart. No morbid appearance had been found in the brain, but there had been no special examination of the pons and medulla.

In the following case, the diagnosis is less certain, both as to the seat and character of the disease; but there was a distinct history of syphilis.

CASE III. Left Hemiplegia with Tonic Spasm; Impaired Phonation and Articulation (Partial Paralysis of the Eighth and Ninth Nerves).—The patient, a schoolmaster, aged 32, came under my care at St. Mary's Hospital, October 15th, 1871, suffering from a peculiar form of hemiplegia, with impairment of articulation and other symptoms, which will be described in detail. The hemiplegia was on the left side, and was accompanied with a degree of rigidity of the affected limbs; these had not wasted at all, but were firm and well-nourished. He had considerable power and range of movement in the left arm, could carry the hand to the head, and use a fork for eating; but all movements were slow, vague, and wanting in precision, and extension of the elbow especially was difficult, on account of tonic contraction of the biceps. The act of walking was very clumsy and slow, and he often stumbled. The left leg was stiff, the toe turned in, and scraping along the floor, the knee unbending, the hips raised. He could not raise the foot to a chair, or flex the knee; the ankle also had little motion. There was no marked evidence of unilateral paralysis of the face, but the lower

lip had a tendency to hang, did not act properly in articulation, and could not be made to assist in producing a whistle. There had previously been dribbling of the saliva. The tongue also was large and unyielding, and not well under command. The speech was very peculiar. Phonation, the laryngeal part in speech, was imperfect, and the voice was squeaky, forced, and unnatural, requiring great effort; but he could produce a loud, deep *Ah!* In speaking, the mouth was expanded greatly laterally and the jaw depressed much more than usual, while the tongue was seen to be almost passive. His enunciation was slow and laboured, the words being brought out one by one, each with a distinct effort. The articulation was indistinct; he could go through the alphabet, but did not pronounce any consonant perfectly, those requiring the action of the lips and of the tip of the tongue being most imperfect. In speaking, the pronunciation was less distinct than in naming individual letters; and a *v* in his own name was always dropped. When he was asked to read, the articulation became worse and worse till, in two or three minutes, the words could not be distinguished. The intelligence was apparently unaffected; he read much, construed Greek well, wrote good letters; but his emotional condition was peculiar, as he was liable to an uncontrollable laugh, in which the expiratory phrase was a continuous squeak, unbroken by the successive explosions of ordinary laughter, quickly subsiding into a mere hiss, which was prolonged till it almost became alarming, when there would be a loud inspiration. The left ear was rather deaf; the optic discs were normal. Since his illness, the patient had become very stout.

The history of these symptoms was as follows. Syphilis was contracted four years before the patient came under observation. A year afterwards, left hemiplegia came on suddenly in the night; motor power was absolutely lost at first, but gradually returned, so that in six months he was able to walk. Twelve months after this attack, the speech gradually became indistinct, and in six months he could not speak at all. At this time, deglutition was very difficult; he swallowed solids better than fluids, these escaping by the mouth; the saliva also dribbled. The difficulty of swallowing was probably lingual, and not pharyngeal, consisting in a failure on the part of the tongue to bring the food within the grasp of the constrictors. Remains of this difficulty were, indeed, still to be observed. Afterwards, he gradually improved up to the point which he had reached when he came under observation. Under the influence of iodide of potassium and galvanisation of the affected limbs, further improvement took place, both in speaking and walking; but it appeared to me that irremediable damage had been inflicted.

The principal facts, in addition to the history, on which I based the conclusion that the disease was syphilitic, were the gradual paralysis of the tongue, lower lip, and larynx, its arrest, and the subsequent amendment. It is this grouping of the paralysis which localises the lesion in the medulla, the nerve-nuclei for the movements of all these parts being within reach of a single lesion in this situation. Disease of the motor tracts of both sides above the pons may affect speech greatly, and produce a relaxed condition of the labial muscles; but I have only met with the same peculiar modification of laryngeal phonation in one other case, and in this the evidence of disease in the medulla or lower part of the pons was complete, as there was cross paralysis; hemiplegia of one side, and facial paralysis of the other, with impairment of voice.

The hemiplegia was probably due to thrombosis of a branch of the right middle cerebral artery, and had nothing to do with the loss of speech which occurred twelve months later.

Syphilitic Affections of the Cerebellum.—The symptomatology of disease of the cerebellum is singularly obscure, corresponding, however, in this respect with the state of our knowledge of the physiology of this nervous centre. I still adhere to the theory of its action, which makes it the seat of the higher and more complex co-ordinations of movements. It is not the sole organ of muscular co-ordination, as was at first inferred from the experiments of Flourens. The spinal cord co-ordinates in a distinctly purposive manner movements which respond to tactile or other cutaneous impressions; but, when movements have to be guided by vision, some special and more complex apparatus is needed to bring muscular actions into relation with visual impressions, which are so far removed in character from tactile impressions (the immediate guides of muscular action), and are correlated with them only by inference and experience. This is essentially, though in a superficial and imperfect way, the view held by Mr. Herbert Spencer, and repeatedly expounded and illustrated by Dr. Hughlings Jackson, namely, that the cerebellum is to space-relations what the cerebrum is to time-relations. Dr. Ferrier's interesting experiments, which appear to make of the cerebellum simply a motor centre for the movements of the eyes, receive no support from the facts of pathology, and his results will probably find some interpretation more consistent with these.

As in the cerebral hemispheres, so in the cerebellum, there may be extensive disease without obvious symptoms, or the symptoms may be

such as will indicate only the existence of intracranial mischief, but afford no indication whatever of the seat of the lesion, namely, pain in the head, vomiting, and blindness; pain, however, being more constant as an accompaniment of cerebellar than of cerebral disease.

The special symptoms of disease of the cerebellum, when such exist, are a peculiar staggering gait and a vague purposeless character of the movements generally, together with a marked loss of vigour and energy. A want of control and co-ordination of muscular actions is evident, but it differs from that due to loss of spinal co-ordination, and it is not increased by closing the eyes. In the later stages, evidences of pressure upon the medulla, or of general intracranial pressure, due to effusion into the cerebral ventricles, may be superadded.

In the cases of disease of the cerebellum which I have watched to a fatal termination, and in which I have made *post mortem* examinations, the cause has not been syphilitic; and in the case I am about to relate, which I considered to be one of syphilitic affection of the cerebellum, no examination was made.

CASE III. Headache; Fits of Uncertain Character; Hemiopia; Later Slight Right Hemiplegia; Sudden Death.—Robert K., aged 26, shoemaker, first came under observation on December 7th, 1865. He had had syphilis seven years previously; six months before, and again recently, had had a fit, no account of which was obtained. Before the fit, he had had pain in the head, and at times saw double. The headache, which was frontal in situation, continued, was worse at night, but was especially exacerbated when the patient stooped or rose from the sitting posture. At times, he was giddy, but he did not fall. Ever since the fit, he had been "dark on the right side," *i.e.*, while he could see an object directly in front or to his left, he could not see anything to his right, and it was found, on testing the two eyes separately, that the inner side of the right retina and the outer side of the left were blind. He saw a little better with the right than with the left eye; the pupils were equal. The patient was a well-built and well-nourished young man, inclined to be stout; pale, but not anæmic-looking, with a depressed expression. Besides the pain, he complained of feeling weak, low-spirited, and deficient in energy. The bowels were open; the appetite failing. He was ordered a couple of aperient pills and four grains of iodide of potassium, with ammonia and gentian.

On the 14th, he was worse; had pain over the brow and in the mastoid regions. When he rose from the sitting posture, his strength seemed to leave him altogether. Vision was as before; the right pupil was a little larger than the left; sleep was imperfect; his appetite was good; his bowels were confined; the stools were of clayish colour; tongue white. The iodide was continued. He was ordered to take, twice a day, a quarter of a grain of pilula hydrargyri, two grains of compound rhubarb pill, and two grains of extract of taraxacum.

December 18th.—He was worse; and quinine was substituted for the iodide of potassium by the clinical assistant. He improved for a time after this change, and, on January 1st, had much less pain in the head, the state of the vision remaining the same, and the right pupil being still larger than the left. The headache, however, returned, and, on January 8th, liquor potassæ arsenitis, in doses of four minims, was given with calumba, in the place of the quinine; aloes and soap pill having also, on the 4th, been substituted for the pill previously taken. He took the arsenic for a month, improving on the whole; but, at the end of this time, he was made sick by it, and it was discontinued. After this, he continued to attend the hospital with the greatest regularity for more than a year, taking nearly all this time ammonia, with gentian and an aloetic pill. The vision was restored, and the headaches ceased, returning only at intervals. For a short time (about February 1867), he complained of tingling in the right face and pain in the right shin, the first of these symptoms acquiring significance from the subsequent history.

In this case, there was no doubt, from the first, as to the existence of some intracranial lesion. The fits alone would not have been conclusive; but the association of these with the peculiar affection of the sight left no room for uncertainty. The situation of the lesion could, moreover, be approximately fixed. The blindness in the left half of the two eyes, of the inner part of the right retina, and of the outer part of the left, showed that the left optic tract was involved at some point between the commissure and the corpora quadrigemina. The diagnosis I formed provisionally was, that there was a tumour in the cerebellum, giving rise to pressure on the optic tract, as in a case I had previously watched to a fatal termination; this conclusion being favoured also by the want of energy, which is a common result of disease of the cerebellum, and by the giddiness. It seemed reasonable to suppose that it might be syphilitic, and iodide of potassium was given at first; but, improvement following the substitution of quinine, it was discontinued before an

adequate trial had been made. After an interval of two years, during which the patient had been seen occasionally, he again came under observation on April 29th, 1869. He then complained that his head felt queer, and had done so for some time. At times he staggered, as if drunk, and a weakness of the right side had gradually come on. This he felt most after walking, and he noticed that he dragged the foot, wearing out the right boot at the toe very rapidly. The right arm was weak. There was evidence of slight paralysis of the muscles of the right side of the face, and articulation was indistinct. Sensation was not apparently impaired. The right pupil was a little larger than the left; both were small. The expression of his face was that of great depression of spirits; he sighed frequently and deeply. He was unable to stick to work, earning only three shillings to six shillings a week, instead of eighteen shillings to twenty shillings; he was indisposed to go out, and would sometimes remain in bed all day. He ate and slept well. The bowels were confined.

The treatment adopted was iodide of potassium, beginning with six grains gradually increased to sixteen, aperient pills, and a succession of blisters behind the left ear. He attended at the hospital for two months, and seemed to improve a little; he gained strength in the arm, dragged the leg less, spoke more distinctly, and did more work; but was never able to go on with it steadily through the day. He was lost sight of at the end of June, and, as was learned subsequently, became an out-patient at the Epileptic Hospital.

He died in January, 1870, and, from his father, the following particulars were obtained. For four months before his death, he scarcely did any work; would begin, but would cut the "uppers" all to pieces; he could not control his movements. He appeared "lost," was unable to give his attention to anything, and was extremely irritable. His appetite was enormous, and he became very fat, but was extremely weak. When he stooped, his face would become so red, that it seemed as if all the blood in his body rushed to his head. On the day before his death, he did one or two odd things, and went home to bed, where he was found next morning, at 10.30, lying in a torpid condition, but answering when spoken to. He refused to rise, and struck violently at a man who tried to drag him out of bed. Later in the day, he had a sort of fit, which consisted in violent movements of the legs only, and he became unconscious, lingering in this condition for four hours, and "dying from the feet upwards," *i.e.*, the feet became cold and the legs motionless. It may be conjectured that the immediate cause of death was hæmorrhage into the membranes at the base of the brain and round the medulla, though a sudden termination is not uncommon in disease of the cerebellum, independently of any such accident, in consequence of its proximity to the medulla. No *post mortem* examination was made.

In my next lecture, I shall consider the syphilitic affections of the brain: the most interesting and important part of my subject.

AMPUTATIONS IN PRIVATE PRACTICE AND IN HOSPITAL.

IN THE BRITISH MEDICAL JOURNAL for January 24th, I observe in Mr. Erichsen's lectures on Hospitalism, that he remarks, in speaking of smashes, such as are seen in railway accidents, occurring in the middle or upper third of the thigh, and in which amputation is performed, that "he is not acquainted with a single case in which such an operation has succeeded in general hospital practice". A short account of the following case, not in hospital practice, from memory, may be interesting. On November 27th, 1872, I was called to C. R., aged about 26, who was at work at one of the iron girders of a bridge on the Whitby and Lofthorne Railway, when it suddenly tilted, falling with the edge across the upper part of the left thigh, cutting to the bone, and causing a comminuted fracture. He was three miles from my house. On my arrival, I found him suffering much from shock. He had lost a large quantity of blood, which had been stopped by tying a handkerchief tightly round the thigh above the wound. He was removed to his home; and, having procured assistance, I amputated, making the anterior flap from the upper part of the wound, and a long posterior flap by transfixing. A fragment about three inches long was removed, which extruded close up to the trochanter. His pulse flagged very much during the operation, but he rallied considerably before I left him. I was sent for late in the evening, as the stump had been bleeding. I found there had been considerable loss, but it had ceased; and as there was no more during the time I remained (about two hours), I did not remove the dressings. Next day, he was in a very good state; and, with the exception of once or twice some tendency to erysipelas, he went on well. One or two small pieces of bone exfoliated about six months afterwards.

T. TINLEY, L.R.C.P.E., etc.