Of the mortality in the cases of primary amputation of the thigh, I need say little. Death, in each case, took place in three days at the farthest, and without any thorough rallying from the shock of the accident. The mode of amputation, or the treatment of the stump, can have little to do with the result.

The amputation of the arm and scapula was complicated with a laceration of the costal wall of the axilla, fracturing and exposing three of the ribs. The man died in twenty-four hours. The death in the arm amputation, as explained in the note, arose from causes unconnected with the operation. It is curious that, in this series, there are no shoulder-joint amputations. I published, in the *Liverpool Medical and Surgical Reports*, 1867, six cases which had occurred to me up to that time, and of which five recovered and one died.

When I was first appointed to the Hospital, twelve years ago, I found amputation below the knee a very serious operation. I became acquainted with hospital gangrene, sloughing of the flaps, and traumatic delirium. It is with great pleasure, then, that I look back on so many successful cases, in later years, of this operation, the mortality of which Mr. Erichsen, in his third edition, states as 40 per cent., from three years' statistics at the principal metropolitan and provincial hospitals.

I have been accustomed to follow the example of a very successful operator, Mr. Hussey of Oxford, in tying all large veins, as well as the arteries. In primary amputation, I have always seen more or less sharp febrile reaction, and I generally meet it with calomel and jalap, plenty of water and tea, and a diet restricted for two or three days to a little beef-tea and milk. As to the dressing of the stump, I usually touch its fresh surfaces with carbolic acid solution, bring them together as accurately as possible with silver sutures; then apply a layer of dry or wet lint covered over with one of lint soaked in carbolic oil; and then, as I was first taught to do by one of Mr. Callender's earlier papers, wrap the limb and stump in plenty of cotton-wool with a bandage loosely applied. I, of course, avoid all unnecessary disturbance of the stump. I do not avoid exposing it to the air during its dressing, and sponge it well with warm water. Sometimes it has not to be dressed from the first for many days. Arms and forearms often unite at once and require no dressing. Where matter forms in the cavity of the stump, and tension appears, I freely remove as many sutures as necessary, and break down all union between the flaps, if necessary, so as to ensure free and rapid exit to all discharges.

In conclusion, I may observe that I think the general results of amputations of late years have been much more favourable, and that the dangers of hospitalism have been much exaggerated. I cannot tell to what the improvement is to be attributed, unless to the superseding of ligatures by the use of catgut-threads cut short and of torsion, and to the greater use of antiseptics.

EPILEPTIFORM CONVULSIONS OF THE RIGHT SIDE, WITH APHASIA.

By J. LOCKHART CLARKE, M.D., F.R.S.,
Physician to the Hospital for Diseases of the Nervous System.

Mr. F., aged 74, a retired officer in the army, had jaundice two years ago, but was not confined to his house. During this period, he had an attack of epileptiform convulsions of the right side, followed by profound sleep for some hours. Six months later, he had a similar attack, the right side of the face being affected first by clonic spasm, and then drawn to the same side for several hours. After an interval of five months, he had another and similar attack; and one more, only a month later. After the last, he was profoundly unconscious for some hours, the arm and leg of the right side being somewhat flexed and stiff. The next day, he could use his right arm and leg, but they were both very weak. He was quite conscious, but had considerable difficulty in expressing his meaning, from inability to recollect the words he wished to use. To whatever question I asked him, he repeatedly answered, "my right arm, my right arm, my right arm is unable." The day after, he was somewhat improved, could walk, use his right arm, and say with difficulty nearly all that he wished, but he was quite unable to write even a short letter; for, whenever he attempted to do so, he substituted wrong words, or words that had no meaning, and, being sensible of his error, he at once gave up the attempt. During the next three days, there was a continuous jerking movement of the right arm, which was at length stopped by morphia.

Four months later, he had a very severe fit of convulsion of the right side. When I saw him, his head was twisted to the right, and both eyes deviated to that side. His lower jaw was in constant motion, and

his right arm was incessantly jerking. After a few minutes, both arms and both legs became convulsed. On no other occasion had the left arm and leg participated in the convulsive movements. The patient was quite unconscious, his bronchial tubes were choked with mucus, and after a few hours he expired.

In the intervals between the paroxysms, the treatment consisted chiefly of bromide of potassium, combined sometimes with digitalis.

A *post mortem* examination was made twenty-one hours and a half after death. The body was extremely emaciated. On opening the head, nothing remarkable was observed in the dura mater. On removing this membrane, a layer of fluid was found beneath the arachnoid, over the convolutions of both hemispheres. The quantity of fluid which escaped on removing the brain amounted to between two and three ounces. Most of the arteries were more or less atheromatous, and the gray substance of the convolutions and of other parts was exceedingly pale. The entire brain was small, and the whole of the left hemisphere was decidedly smaller than the right. The posterior lobe did not reach so far back on the left side as on the right, and was less pointed. The posterior half of the third frontal convolution, and the lower parts of the transverse frontal and transverse parietal convolutions, were very much wasted and depressed, so that the processes which run from them to the insula, and which normally are almost covered by them, were entirely exposed when the temporo-sphenoidal lobe was drawn aside. The heart was of average size. Its muscular tissue was healthy, but the wall of the left ventricle was much thickened. The aortic and pulmonary valves were quite healthy, as were also the tricuspid and mitral, with the exception of being a little atheromatous. The liver was small and firm to the touch, with a tendency to cirrhosis. There was a broad antero-posterior groove on the convex surface of its right lobe. The gall-bladder contained four gall-stones of the size of large marbles. Both kidneys were contracted; their cortical portions were diminished, and they were nodulated on the surface at different parts. On the surface of each, also, there was a cyst; that on the left kidney was about the size of a large marble. The spleen was small; its investing membrane had the appearance of having been inflamed, and its substance was softer than natural.

OVARIOTOMY SUCCESSFUL IN A GIRL EIGHT YEARS OLD.

By T. SPENCER WELLS, F.R.C.S.,
Surgeon to the Queen's Household and to the Samaritan Hospital.

OVARIOTOMY is now so generally and so often practised that a record of any single case is only interesting when there is something unusual in the history of the patient, or some peculiarity in the operation. The facts that an ovarian tumour of sufficient size to render surgical aid necessary has been observed in a girl only eight years old, and that ovariectomy has been successfully performed, appear to be of sufficient importance to call for the publication of a short account of the case.

On January 5th, 1874, a little girl, eight years of age, of German parentage, born in California, was brought to me from San Francisco, having been sent over by the advice of Dr. Levi C. Lane and Dr. Regensberger of that city. She was rather small for her age, and the central part of the abdomen was occupied by a tense movable cyst. I arranged a consultation for the 7th of January with Sir Wm. Jenner and Dr. Sutor; and we all agreed that there was a cyst in the abdomen, and that I should tap it with a fine trocar and aspirator to ascertain if it were hydatid or ovarian.

On the 8th of January, Mr. Thornton administered a little chloroform, and, with the kind assistance of Dr. Sutor, I removed twenty-six ounces of opaque slightly viscid fluid, which I had no doubt was from an ovarian cyst. A hard substance like half an orange, either the contracted cyst or a hard portion of the ovarian tumour, was felt to the left side, after the fluid had all escaped.

The child did not suffer at all after the tapping, but the fluid soon began to distend the cyst again.

On the 23rd January, Sir William Jenner saw her again, and wrote: "There is nothing in the chest to contraindicate operation. The result of tapping is very interesting. I fancy now I can feel connection with pelvic contents on moving the mass, as if there were a long pedicle. I have advised very decidedly the operation of removal. Children bear operations well."

On January 27th, assisted by Dr. Tracy of Melbourne, and by Mr. Thornton, chloro-methyl having been administered by Dr. Day, I made an incision, nearly four inches long, through the abdominal wall