by pinching its nerve. But there is more in the matter than this, I feel convinced, at least in many instances. In Mr. L. Holden's singular case of anomalous affection of the nervous system in a boy, aged 13, the tetanic seizures were brought on, not only by pressing his "bump", but any spot over the ten upper dorsal vertebrae; moreover, occasionally the "bump" lost, for a time, its peculiar power of producing the phenomena, though this seems to have persisted in the spinal integument. The same, or very similar, effects seem to have been produced by raising him into the upright posture. Again, the periodic quasi-maniacal daily paroxysms occurred without any provocative stimulation. The whole series of nerve-troubles seem to have depended mainly on the bump, since they disappeared as soon as the tumour was removed, and had not recurred four years later, when the last report was made. The only view I can take of such a case as this is, that the terminal extremities of the nerves in the bump, being organised in some peculiar way—say, having their axis-cylinders exposed—kept up a constant state of hyperexcitability or tension in the nervous centres, which was ever ready to display itself demonstratively on slight provocation, or even without any. This peculiar organ of excitement being removed, the nerve-centres reverted to their ordinary condition. A case which I have recorded in *Functional Nerve-Disorders*, p. 570, presented some resemblance to Mr. Holden's. The occurrence of physiological hyperexcitability in the male frog at the copulating season is well known, and is worth referring to in connection with such cases as the foregoing. The thumb of each hand becomes covered with large papillae, and, so long as these are present, the animal is enabled to grasp the abdomen of his mate distended with ova without any exertion, apparently, of volition, or any fatigue, even for weeks. If, however, the papillae be shaved off, the power of grasping in this remarkable way is lost. (Todd and Bowman's *Physiology*, vol. i, p. 315.) They regard the cord of the frog, or, rather, that segment of it with which the nerves of the anterior extremities are connected, as being, during this period, in a state "of exalted polarity".

[To be continued.]

A CONTRIBUTION TO THE PATHOLOGY OF ALBUMINUS EXPECTORATION AFTER THORACENTESIS.

By ALFRED B. DUFFIN, M.D., F.R.C.P.,
Physician to King's College Hospital.

DURING last summer, a lively discussion took place in Paris on the causes of albuminous expectoration after thoracentesis for pleuritic effusion. A very complete analysis was published in this JOURNAL during October last, reviewing not only M. Terrillon's thesis, which started the discussion, but also the various hypotheses that were advanced in explanation. They were four in number: 1. Perforation of the lung by the trocar (MM. Woillez and Marrotte); 2. Spontaneous rupture of air-vesicles after destruction of pulmonary epithelium (Férol); 3. Resorption of the residual liquid from the pleura after puncture; 4. Transudation of the sero-albuminous liquid through the alveolar walls by means of rapid pulmonary congestion and oedema (Héraud, Moutard-Martin, Béhier, Terrillon, etc.). The reviewer strongly expressed his adhesion to this last hypothesis, which I also believe to be the true one. Neither of the authors named has, however, pushed his theory to its consequences, so as to indicate the mechanism by which the characteristic expectoration may be presumed to be secreted.

Before contributing my quota to this task, I may say a word regarding a fifth hypothesis, which has been suggested by Professor George Johnson (BRITISH MEDICAL JOURNAL, Oct. 25, 1873). "While the lung is compressed by a copious liquid effusion into the pleura, it is probable—nay, it is certain—that, in consequence of the more or less complete blood-stasis within the compressed lung, coagula will form in the minute pulmonary vessels, and especially within the pulmonary veins." After thoracentesis, "it is obvious that the obstruction of the pulmonary veins by coagula would cause a backward passive congestion of the capillaries, and a consequent serous transudation into the air-cells and bronchi." The objections that tell against this view appear to me as follows. 1. Albuminous expectoration has generally been confined to those instances where the amount of pleuritic effusion has been greatest, and the lung consequently most crushed and anemic. 2. Thoracentesis has been performed four times on the same subject, and each time albuminous expectoration has followed (Boule's case, Béhier's lecture, *Union Médicale*, 1875). 3. In the four fatal cases* recorded, no mention is made of coagulation in the pulmonary vessels. The lung was in each

instance choked from top to bottom with the characteristic fluid, so that the coagula would require to be extremely numerous. 4. The phenomenon is itself a very transient one, seldom lasting beyond twenty-four hours, although in that time many pints of fluid may be expectorated. How are the coagula assumed to be subsequently disposed of?

Reverting to the hypothesis of rapid pulmonary congestion and oedema: the authors who uphold this have assumed that it is brought about by the intervention of the pneumogastric nerves, but they fail to inform us by what links of the chain the result is arrived at.

The effects of section of the vagi on the pulmonary circulation have afforded very variable results in the hands of different observers. To obtain any at all, both nerves must be divided. Legallois (*Expériences sur le Principe de la Vie*, 1812), whose experiments were chiefly performed on young rabbits, found section of the pneumogastric in the neck to be almost constantly followed by engorgement of the lung. De Blainville and Provençal (Claude Bernard, *Système Nerveux*, vol. ii, p. 353), on the other hand, missed all trace of pulmonary change. Claude Bernard (*ibid.*, p. 354) admits that death is not necessarily the result of asphyxia at all. He regards the pulmonary lesion, when present, as primarily of traumatic origin, and as induced by the difficulties that arise in the mechanical act of respiration. "The movements gain in amplitude what they lose in frequency; they tend to introduce the same amount of air into the lung." An animal breathing from a bell-jar takes in a notably larger amount of air at each breath after its vagi have been divided than before, and a traumatic emphysema rapidly results (*ibid.*, p. 367).

Dr. John Reid (*Experimental Investigations into the Functions of the Eighth Pair of Nerves*) admits the great frequency of pulmonary congestion and the effusion of frothy serum after section of the vagi; but he has also repeatedly known both to be absent. "The effusion of frothy serum into the bronchial tubes is not a necessary consequence of division of the vagi, even when the lungs were found loaded with blood, and when the respiration before death was very laboured." "I have also satisfied myself that this frothy fluid is not mucus, but is the frothy serum so frequently found in cases where the circulation through the lungs has been impeded for some time before death" (p. 202). He believes that all the morbid changes observed in the lungs may be traced to the diminished frequency of the respiratory muscular movements. Reid also quotes observations on the human subject of death with extremely slow respiration, where the vagi were apparently healthy, but great congestion and oedema of the lungs existed (*ibid.*, pp. 207-209).

It will thus be seen that Bernard and Reid are practically at one in ascribing pulmonary congestion after division of the vagi to interference with the mechanism of respiration; and I may add that, so far at least, Traube is of the same opinion. That we have not the whole truth even here, is rendered very probable by Valentin's observations (quoted by Schiff, *Nerven-Physiologie*, p. 411). He has shown that marmots, in whom the vagi have been cut during hibernation, get pulmonary congestion; although the mechanism of breathing is rarely materially disturbed, the animals do not swallow, and no secretions are formed in the mouth.

Such being the facts in reference to the pneumogastrics, I was led to inquire what evidence there might be of the existence of some vaso-motor tract influencing the pulmonary circulation, and capable of being interfered with by the pressure of an extensive pleuritic effusion.

It seems to me that we have in a limb compressed by Esmarch's bandage and rope a condition closely analogous to that of a lung compressed by an extensive pleurisy. When the apparatus is removed, it is often observed that a blush of redness diffuses itself over the part. Sometimes, but by no means invariably, a troublesome hæmorrhage will then ensue from the cut surfaces. The reason of these I take to be, that the pressure has not only emptied the vessels, but has also compressed the nerves of the limb, and among others the vaso-motor nerves. When the blood returns to the extremity, it finds the small vessels not only empty, but lax, owing to vaso-motor paralysis. A similar condition exists in a lung which has just been relieved from the pressure of a considerable effusion. Its vaso-motor nerves, I take it, are also compressed. As the lung again expands, the tension thrown upon its small vessels is not only sudden, but very great. When the nerves have had a few hours to recover, and blood to revive upon, their controlling action again asserts its influence, and the arterial tension is brought up to the normal. We can thus account for the rapid and violent onset, as well as for the very short duration, of this albuminous expectoration. It has generally been found proportionate to the rapidity and extent to which the lung again expands, and to the amount of pressure previously exercised. In eighteen out of twenty-one of M. Terrillon's cases, from 2000 to 5500 grammes of fluid were evacuated from the pleura, and in all instances rapidly.

Ranvier's researches on the nature of oedema (*Comptes Rendus*, lxi, p. 25) strongly suggest that the vaso-motor nerves of the lung play an

* Two quoted by Terrillon, one by Béhier (*Union Médicale*, June 24th, 1873), one by Dumontpallier (*ibid.*)

important part in this proceeding. He found that in dogs, in whom he had tied the inferior vena cava, œdema of the legs only ensued when the sciatic nerve was also cut. If the nerve were only divided in the one limb, the œdema was confined to that side. It is well worthy of notice, that the limb became œdematous from one to two hours after the section—a time closely corresponding with that of the first appearance of albuminous expectoration after thoracentesis. Proceeding further, if, instead of dividing the sciatic, he cut the three last lumbar and the sacral nerves in the spinal canal, then, in spite of complete paralysis of motion and sensation, *no œdema ensued*. Ranvier concludes that a paralysis of vaso-motor nerves must be added to the venous stasis to produce acute dropsy.

Of the vaso-motor nerves of the lung itself but little is known. In some experiments on the peripheral stumps of the spinal roots of the first thoracic ganglion, Schmiedeberg ("Ueber die Innervations-verhältnisse des Hundesherzen," Ludwig's *Arbeiten*, 1872) inquires whether, in the phenomena which result, we may not be dealing with a change in the calibre of the pulmonary vessels. Bernard, after removing the same ganglion (*op. cit.*, vol. ii, p. 519) found violent pleurisy and an extraordinary injection of the whole of the lung-substance. After removing all the four upper cervical ganglia (p. 537), he found an enormous liquid effusion into the pleura and pericardium, with hepatization of portions of the lung. The important observations of Schiff (*Nerven-Physiologie*, p. 409) induce him to ascribe the lung-œdema consequent on the section of the vagi to a *neuro-paralytic hyperæmia*. "The vagi usually contain a great part of the vaso-motor nerves of the lung. The more of these vaso-motor nerves course in the track of the vagus, the more general the hyperæmia, the younger the animal, the more rapidly does it set in, as is always the case with neuro-paralytic hyperæmias." He further accounts for the discrepancies among observers by suggesting that the vaso-motor nerves of the lung may be entirely absent from the vagus, and take some other track. With regard to the vaso-motor nerves of the liver, the contrasting experiments of Cyon and Aladoff (*Bulletin de l'Académie de St. Petersburg*, vol. viii, p. 91), and of Eckhard (*Beitrag*, vol. vii, part i, p. 19), indicate that irregularities in their course are frequent; and the observations alluded to in this paper point to the same inference with respect to the distribution of the pulmonary vaso-motor supply. In the majority of instances, they would seem to join the vagus high up, so as to be included in the section in the neck. In a few instances, they join lower down, and no lung-œdema ensues. For the production of albuminous expectoration after puncture, a previous crush of the lung by the pleuritic effusion sufficient to temporarily paralyse these nerves would appear to be the essential factor.

THE TEACHINGS OF THE GENERAL INFIRMARY AT LEEDS WITH REGARD TO THE OPERA- TION OF OVARIOTOMY.*

By C. G. WHEELHOUSE, F.R.C.S. Exam.,
Senior Surgeon to the Infirmary.

In the brief space of time allotted even to the major communications brought before this Society, it is impossible to do more than glance at some of the chief points involved in such a subject as that to which I ask your attention. I do not, therefore, intend to enter into discussion upon any of the points I bring before you, but merely to state what I, in the experience I have obtained in this institution, have learned concerning them; discussion will follow more appropriately afterwards.

In approaching the operation of ovariectomy, we are brought face to face with many difficulties; and, as the best method of dealing with these can only be solved by direct appeal to experience, it is with a view to bring out the accumulated experience of this institution that I have charged myself with the delivery of this Address. The points which I desire to bring briefly under your observation are the following.

1. The selection of cases for operation.
2. The influence upon the result of the condition of the peritoneum at the time.
3. The best method of dealing with adhesions.
4. The best method of dealing with the pedicle.

Upon each of these points, I am satisfied that the experience of this infirmary has taught me valuable lessons—lessons which I could have learned in no other way, and, in as few words as possible, I desire to make you sharers in the advantage I have thus reaped. First, then, as to the selection of cases. Bearing in mind that many of these tumours are of a mixed character, partly solid and partly fluid, we may yet, as a rule, classify them generally under these two heads. Accordingly, in

examining an ovarian case, I always direct my attention, in the first instance, to this point; for I know that, according as I have the one or the other kind of tumour to deal with, not only will widely different lines of treatment be required, but that as different an opinion also, as to the advisability of any operation being performed, will probably hang thereon.

Take, for instance, the case of a purely solid tumour first, and consider how great are the difficulties which, on the very threshold of our inquiry concerning it, may await our diagnosis. Consider how many things there are which *may* be, and every one of which *have* from time to time been, confounded with ovarian tumour: pregnancy, extra-uterine foetation, uterine fibroma, or the many forms of internal disease or displacements to which the uterus is liable; peri-uterine hæmatocele, fecal accumulation, mesenteric tumour, enlargement and disease of the kidney, enlargement and disease of the spleen—and you will not wonder when I tell you that I have learned, whenever a purely solid tumour is presented to me for examination, that I must never set the case down as one of ovarian disease, until, by an exhaustive process of elimination, I have assured myself that it can be nothing else.

The degree of fixity or mobility of such a tumour is a point to which I have learned to attach great importance, and also the presence or absence around it of ascitic fluid. I long ago learned to believe, and, up to the present time, I see no reason to doubt the correctness of the conclusion that, with a view to ovariectomy, no more favourable arrangement of circumstances can present itself than a solid ovarian tumour, freely movable, and surrounded by ascitic fluid; whilst, on the other hand, a solid ovarian tumour, fixed and immovable, I invariably approach with the most serious consciousness of impending difficulty. It is easy, I think, to understand why this should be so. If a solid ovarian tumour be tied in position by adhesions, the probability is, that such adhesions will be pelvic and posterior; they will be situated at the very points at which it will be most difficult to separate them, and, even supposing this successfully accomplished, the raw surfaces produced will occupy the very positions which are fraught with the greatest after-dangers to the patient. It will be into the pelvis, and to the back part of the body, that blood and effused fluids will gravitate; and from this part should they undergo decomposition, they will find the readiest and most direct entrance into the circulation, to the almost certain death of the patient.

I have said that a solid tumour, movable and surrounded by ascitic fluid, has, in my hands, proved the most favourable of all forms for operation; and that leads me to an observation which, I believe, all practical ovariectomists will endorse, though few attempts have been made directly to explain the reason why, after such operations, the patient generally progresses towards recovery in so remarkably quiet a manner. The question whether it is better to perform the operation whilst the patient is in robust health, or to wait until she is somewhat enfeebled by the disease, has been ably discussed both in this country and America; and, whilst all the earlier operators recommended that it should be done under the former condition, nearly all are now agreed that the greatest success has attended such as have been undertaken under the latter. Many reasons have been given for this, but I have nowhere seen it distinctly stated that the difference in the result is mainly dependent, as I believe it to be, upon an altered condition of the peritoneum.

Peaslee comes nearest to this point when he says, "At all events, a long distended state of the peritoneum very much diminishes its sensibility and the danger of traumatic peritonitis" (*Ovarian Tumours*); but neither he, nor any of the authorities whom I have consulted, lay any stress upon the altered structural conditions of the membrane itself. If you consider for a few moments what is the normal condition of the peritoneum, both in structure and as to function, you will have no difficulty, I think, in understanding the ground upon which I assert that I would rather operate in a case where I know that it has undergone structural change than in one in which it is presumably perfectly healthy.

Physiologically the largest lymph-sac in the body, it is, as you are aware, perpetually engaged in the process of absorption; and if, in its unaltered condition, septic fluids be presented to it, their immediate absorption is, of course, the result; and structurally the largest serous membrane in the body, it is, of all others, the most liable to become inflamed under conditions of provocation. Thus, in the performance of ovariectomy with a healthy peritoneum, we have the greatest liability to the two conditions which, of all others, are fraught with the greatest danger to the patient, viz., peritonitis, which alone destroys one-fourth of the cases which die, and septicæmia, which carries off at least as many. If, however, the membrane have undergone such changes, as the result of disease, as either weaken its power of absorption, or so change its structure that it is, in reality, no longer an active serous membrane, by so much, according to my experience, is the chance of the success of an operation improved; and this is a fact which, I think, my

* Read before the Leeds and West Riding Medico-Chirurgical Society.