invariably present ; and I have seen two cases where there has not been any pain at all, and another when the pains came on a week or two after the first attack of fever. The fever in this latter case was much more intense than I have seen it in any other case. In another case, swelling of the feet and ankles occurred from a week to a fortnight after the fever, and continued for three or four weeks, when it disappeared spontaneously. In this case, also, the fever was very severe. The pains often continue some time after an attack of dengue.

In dengue, as in many other diseases, it is possible to do too much. It is, I believe, better to treat the urgent symptoms in each case without adopting any routine plan of treatment. Purgatives at the commencement are not often desirable, for the patients are so exceedingly weak that they cannot well bear the effects of a brisk purge, and the patient seems worse after it than before : and, again, the pains are frequently so severe that the necessary movements during the action of a purgative are intensely agonising to the sufferer. When, however, the pain and debility are not great, a purgative is useful, as also is it when there is evidently great constipation, unless the other symptoms forbid it.

Following the advice of some Calcutta practitioners, I used the tincture of aconite in small doses, frequently repeated, with a little syrup, as a remedy against the severe pains, but I did not find it very serviceable. After reading Dr. Charles's clinical lecture on dengue, I tried the tincture of belladonna, and it proved very useful, in doses of from eight to twelve minims. The pain was quickly relieved, and the patient had sleep. The remedy need not be continued long after the pain is relieved, and the patient's restlessness calmed. It may be necessary, however, to repeat it if these symptoms should return. No other medicine is usually necessary, except perhaps a saline mixture, containing liq. ammon. acet., sp. ætheris nit., tinct. hyoscyami, and aqua camph.

The severe headache may be relieved by ice applied in a bladder to the head, and the strength should be kept up with nutritious soup or beef-tea, and milk, with occasionally a little brandy in it, etc. When the fever has abated and the terminal rash disappeared, a tonic is generally necessary, and a generous diet, with beer or wine, to bring up the patient's strength, which is often very much reduced. Patients often suffer from pains in the joints after an attack of dengue ; but they usually pass off soon, and seldom require special treatment.

THREE CASES OF SUDDEN DEATH.*

By J. MILNER BARRY, M.D., Tunbridge Wells.

THE first occurred in a man, aged 41, who during life had only complained of dyspeptic symptoms, including a peculiar choking sensation coming on at nights, and relieved by walking about. The heart and lung-sounds were apparently normal. Death occurred suddenly in the night. At the *post mortem* examination, the left pleura was found full of blood, which had escaped from a small sacculated aneurism, the size of half a walnut, of the descending portion of the thoracic aorta situated behind the heart. The case was characterised by a singular absence of the usual concomitant symptoms of aneurism. There was no dysphagia ; no erosion of the vertebrae. The choking sensation might be accounted for from the fact of the œsophagus being attached to the aneurism, lying directly on it. Dr. Barry mentioned that both Dr. Stokes and Dr. Walshe refer to the extreme difficulty of positive detection of small aneurismal sacs behind the heart.

The second case occurred in a young lady, aged 21, active, well-nourished, and with a good healthy colour. She was subject for nearly twelve months to nearly constant vomiting, and known to be the subject of heart-disease. On auscultation, loud blowing murmurs were heard over the pulmonary aortic and mitral valves, a thrill was perceptible over the situation of the pulmonary valves, and the area of præcordial dulness was increased. The vomiting continued unchanged, and, though apparently somewhat hysterical in character, seemed mainly attributable to irritation of the pneumogastric nerve from the diseased heart. She died suddenly in the night. At the *post mortem* examination, the pericardium was found generally adherent. The tricuspid and aortic valves were healthy. Two of the pulmonary valves were adherent by thin edges, and one of the mitral valves shortened and tied down. A short funnel-shaped canal passed through the septum ventriculorum from below the aortic valves to behind the tricuspid. Two rough ridges, one above and the other below the opening, probably impeded and roughened the current of blood, and gave rise to the peculiar "swashing" character of the murmur heard during life. The direction

of the canal was such as to cause the blood to pass more readily from the left ventricle into the right than *vice versa*. The entire absence of cyanosis was a noticeable feature in this case. No ulceration was found in the stomach ; but there appeared to be a sort of constriction midway in the greater curvature, where probably the spasmodic action was greatest, and this may have accounted for the peculiar character of the vomiting ; the food being suddenly and forcibly shot out, and in the convulsive effort made to reject it, a portion was probably dashed on towards the pyloric end, and then retained till the vomiting had subsided, thus, in a measure, accounting for the nutrition of the patient being maintained.

The third case occurred in a single lady, aged 41, five months previously. Dr. Barry had attended her with chronic peritonitis ; from this she quite recovered, and had never complained of any chest-symptoms. She died suddenly whilst dressing. At the *post mortem* examination, the pericardium was found distended with about eighteen ounces of serum and soft blood clots. The heart was loaded with fat and flabby, and showing fatty and granular degeneration, and almost total disappearance of muscular striæ under the microscope. The aorta was stiff, rugose, atheromatous, and dilated. A small aneurismal pouch, one inch above the aortic valves, as thin as a wafer, had given way, and allowed an escape of blood into the peritoneum. This patient's brother had died suddenly some months previously at the age of 52. The clinical lesson taught by the above and similar cases, as Dr. Barry insisted, was that, in cases of extremely sudden death our attention should be directed to the heart first, and not to the brain.

ON THE TREATMENT OF WOUNDS BY ALKALINE APPLICATIONS.

By R. T. MANSON, M.R.C.S.E., etc., Howden.

ALKALIES, as local applications, check alkaline secretions. Laudable pus has an alkaline reaction. The application of alkaline substances may, therefore, be supposed capable of beneficial action on pus-secreting surfaces. The official linimentum calcis, spread on fine cotton-wool and applied to the surface of a wound, seemed to afford the best method of testing the value of the theory. It has been tried by me in a great variety of surgical cases, from bruised fingers to amputations, and with uniform success. The advantages of its use are : lessening the secretion of pus from a wound ; easiness of application ; extreme adaptability to irregular surfaces ; prevention of the access of atmospheric air to the wound ; promotion of healing by first intention, or, when that does not occur, promotion of healthy granulation ; doing away with the use of nitrate of silver or sulphate of copper ; ease to the patient ; and saving of time to the surgeon. Used on lint, the linimentum calcis is not nearly so valuable as when used on fine cotton-wool. Lint does not get into the holes and corners of an irregular wound as cotton-wool does ; it is not removed so easily ; it becomes more easily dried and hardened, and possesses less of the nature of an artificial skin than cotton-wool. Linimentum calcis has long been used for burns and scalds. Some surgeons think it a nasty application ; but, when made with olive-oil instead of linseed-oil, as was formerly the case, the nastiness is abolished. The *modus applicandi* is as follows : On a piece of fine cotton-wool, rather larger than the surface of the wound to which it is to be applied, the linimentum calcis is poured and spread over the wool, which is then placed over the wound and covered with common wadding. Either bandages or strips of plaster may be used to keep the lint in position.

CASE I.—H. P., a pitman, while riding "in bye" to his work, incautiously raised his head, which came in contact with the roof of the pit. When I saw him, in addition to concussion of the brain, his scalp was torn, scored and grooved from forehead to occiput ; one ear was cut through, and he had a bruise on the face. I put the bits of skin in position as well as I could, and applied linimentum calcis on cotton-wool. The case did well, no other treatment being used.

CASE II.—J. S., also a pitman, was knocked down and a rail-plate forced into his mouth ; while he was so lying, the cage came down on his head. There were an extensive wound in the soft palate, fracture of the right superior maxillary bone ; the mouth was torn open ; he had scalp-wound, one ear cut, and a finger ripped up from end to end. The external wounds were all treated as in the last case, and the man made a good recovery.

CASE III.—R. T., a pitman, had compound dislocation of the ankle, for which I amputated the leg in the lower third. The flaps did not unite by the first intention, and some unhealthy action was set up. Poulices were applied, but changed for the wool and linimentum calcis, and the stump has done well.

* Read before the East Sussex District Meeting of the South Eastern Branch.