CLINICAL LECTURES

DELIVERED IN

THE OUT-PATIENT DEPARTMENT OF THE
HOSPITAL FOR SICK CHILDREN,
GREAT ORMOND STREET.

By W. B. CHEADLE, M.D., F.R.C.P.,

Assistant-Physician to the Hospital; Senior Assistant-Physician to St. Mary's Hospital; etc.

III.—MENINGITIS.

CASE.—*Constipation.*—*Vomiting.*—*Retraction of the Neck.*—*Insensibility of Pupils.*—*Hydrocephalic Cry.*—*Convulsions.*—*Squinting.*—*Insensibility.*—*Blindness.*—*Treatment by Iodide of Potassium.*—*Complete Recovery.*—Instances of recovery after symptoms of cerebral inflammation in children, of so serious a character as those which have occurred in the case of this little girl, Charlotte B., are extremely rare. In drawing your attention to her condition from time to time during the past six months, I have been careful to warn you that the improvement which appeared so marvellous might be unreal and delusive after all, and that some sudden access of convulsions, paralysis, and coma, might put an end to all the hopes excited by the seeming success of the treatment which had been adopted. But it has not been so; there has been no relapse. With the exception of some imperfection of vision still remaining, the child is now—more than six months since the first onset of the disease—to all appearance perfectly well. The improvement has been so uninterrupted and steadily progressive for so long a time, that I believe it is real and substantial, and that this will prove one of those rare instances of complete recovery from the meningitis of infancy.*

Before I say more upon this subject, however, let me briefly relate the history of this case for the benefit of those who have not been able to follow it closely throughout its course. Some of you will remember the condition of the child when she was first brought here on the 5th February. She was at that time seven months old, and at the breast, a plump and well nourished child. She lay taking no notice, with her head drawn rigidly back towards the spine. The pupils were somewhat dilated, oscillating slightly only under the stimulus of strong light. Although but half conscious to ordinary impressions, she screamed on being touched, and resisted all attempts to feel the pulse or listen to the heart. These symptoms had lasted several days, and had been preceded by constipation and severe vomiting. There had been no convulsions; there were no signs of teething; the abdomen was not sunken; the tongue was thickly coated, and the stools were reported to be very dark and offensive. The child could with difficulty be persuaded to take the breast. She was in a state closely corresponding to what is described as the second stage of acute hydrocephalus, when effusion has commenced as a result of the cerebral inflammation.

With a view of increasing the action of the bowels and relieving vomiting, rather than of obtaining any specific effect of mercury, a grain of grey powder was ordered three times a day. Four days afterwards, there seemed some small change for the better; the neck appeared less rigid, the bowels were more open, and the child took food with less difficulty; otherwise the condition remained unchanged. The grey powder was continued in doses of from half a grain to one grain three times a day for ten days, at the end of which time the condition of the patient was highly unsatisfactory. The head was still rigidly retracted, the pupils dilated and insensible, the vomiting constant. The bowels were indeed more regular, but against this gain was to be counted the supervention of another grave and significant symptom—the child, which lay perfectly quiet and half unconscious when still, uttered violent piercing screams when it was moved. The prognosis was extremely unfavourable. Two grains of iodide of potassium were now ordered to be given every six hours in two teaspoonfuls of caraway water; and this was continued for a week, the medicine being generally retained, although vomiting continued after food, and on being moved. No change was noted, however, until another week had elapsed, when the case seemed still more hopeless. The mother stated that the vomiting had become more incessant; for the last two days the medicine had been rejected as well as food, and squinting and convulsions had occurred the day before. The child was paler, and insensible to all external impressions. In the hope that the medicine might be retained if the quantity of fluid given at a time were reduced, a grain and a half of iodide of potassium in a

drachm of caraway water was ordered to be given every two hours. This treatment was, I believe, faithfully carried out by the mother, who is a very respectable, intelligent woman, willing to do anything which might save her child, and duly impressed by my urgent exhortations that the strict observance of my directions afforded the only possible chance of success.

When the child was again brought in the course of four days, there appeared for the first time some decided improvement; the vomiting had ceased, and the dilated pupils showed some slight contraction under strong light: the neck was still, however, rigidly drawn back. The improvement thus set in continued steadily, the medicine being still regularly administered every two hours. The child began to take food more readily; the stiff retracted neck gradually relaxed, and at the end of a fortnight she was able to sit up, became very restless and irritable, began to take notice of noises, and of her mother's coaxing and endearments. On careful examination by repeated experiments, it was ascertained that she was completely blind. The pupils remained only very slightly sensitive to light. The pulse, which was examined for the first time, was found to be about 100.

The iodide of potassium was continued in the same doses and with the same frequency for nearly three months, without any signs of iodism, except a certain amount of running at the eyes and nose. The patient progressed favourably, grew fat, began to utter intelligible sounds, and was in fact by this time perfectly well, with the exception of being absolutely blind. Gradually after this, however, sight began to return; the child would catch vaguely and unsuccessfully at any bright object waved before its eyes; and this improvement has gone on until she now sees with tolerable clearness, and can lay hold of things put before her with accuracy and certainty. As soon as it was discovered that the loss of perception through the eyes was something more than a passing manifestation of the general insensibility to external impressions—that since, as the sensorium regained its powers, the sense of sight did not return, while the senses of hearing and of touch returned, the fault must be in the eye itself or its special nerve, not in the brain—an examination was made with the ophthalmoscope. Mr. Walton, who was kind enough to conduct this for me, ascertained that the optic discs were exceedingly vascular and the veins large, as in optic neuritis in the stage of recovery. There was no sign of retinitis, no deposit in the choroid. The ophthalmic examination was not, however, satisfactorily effected until sight had in great measure been restored; several previous attempts having failed, owing to the extreme restlessness and irritability of the young patient when under manipulation.

You will perhaps have observed that I have restricted myself to the general term "meningitis" in speaking of this case; and I have done so purposely, because I am unable to offer any positive opinion that the cerebral inflammation which certainly existed was associated with tubercle. The terms tubercular meningitis and acute hydrocephalus are used somewhat loosely; and I wish to point out to you that not only are there several distinct affections exhibiting very similar symptoms, but in which there is no organic affection of the brain, which are constantly mistaken during life for cases of acute hydrocephalus, but also that the term is made to include several distinct morbid conditions of the brain and its membranes—that is, there are several diseases which simulate acute hydrocephalus, and acute hydrocephalus comprises distinct morbid conditions of the encephalon. By many, the designation of acute hydrocephalus or tubercular meningitis is applied to any case accompanied by cerebral symptoms in children, while by a few authorities these terms have been more carefully restricted to inflammation of the cerebral membranes, associated with the deposit of tubercle. The result of this discrepancy has been a wide difference of opinion as to the fatality of the disease. Some statistics, collected by Sir Thomas Watson from various sources, show an average mortality of one in four only. Dr. West, on the other hand, declares that he has never met with a single instance of recovery from advanced acute hydrocephalus (using the term clearly as synonymous with tubercular meningitis); that he has known but one in which the child got well after the second stage had commenced; and that he has seen but very few which ended favourably, even though they came under treatment in the very earliest stage. M. Guersant of Paris, whose experience was very large, regarded the prognosis of tubercular meningitis as equally serious, and affirms, in a passage quoted by Dr. West, that "tubercular meningitis may sometimes recover in the first stage, though the nature of such cases is always more or less doubtful. In the second stage, I have not seen one recover out of a hundred. As to patients in whom the disease has reached the third stage, I have never seen them improve even for a moment". Bouchut, who has studied the subject of tubercular meningitis with minute care, agrees that the majority of the recoveries which take place are from the first stage, rarely from the second, yet that some have been recorded in the third. Niemeyer mentions only

* This prediction has been verified. The child remained completely well when last seen (February 1872), twelve months from the commencement of the attack.

two cases of recovery from undoubted tubercular meningitis; and Trousseau declared he had seen but two instances in his extensive experience, in one of which the diagnosis was subsequently verified by *post mortem* examination, the child having died of some other disease a few months afterwards.

Now, the writers who take so desponding a view are speaking of the tubercular disease only, while the more sanguine are those who have been less precise in their definition, and have probably included under the term hydrocephalus various forms of cerebral disturbance which closely resemble tubercular meningitis in symptoms. And then most of the recoveries have been in the first stage; and, as Guersant remarks, the nature of such cases must always be to a certain extent doubtful. Sir Thomas Watson shrewdly guesses that, in many of the so-called recoveries from acute hydrocephalus, the disease has been what he terms simple encephalitis—not dependent on tubercle; and I think we might go further, and say that in many there has been no inflammation of the brain or its membranes at all. Trousseau, indeed, accounts for the apparent success of certain practitioners in dealing with this disease, by the supposition that they have mistaken for it typhoid with cerebral symptoms. The more strict observers are unquestionably right according to my experience: true tubercular meningitis is almost always fatal. I have seen one case only in which the disease had advanced so far that the diagnosis could not be mistaken, in which apparent recovery took place for a time; but the patient eventually fell a victim to the same disease, which shortly returned with fatal severity. And yet it seems absolutely certain that complete recovery has resulted in a few rare and exceptional cases. Whether the one before us must be ranked as one of true tubercular meningitis is an interesting problem, but one which we shall not find it easy to solve with certainty. That there was meningitis there can, I think, be no reasonable doubt; but, then, was this set up or accompanied by tuberculosis?

Let me explain to you first why I believe the case to have been one of acute hydrocephalus—not some form of disorder simulating it—and then we may attempt to determine with which form of that disease it may be classed.

There is such a vast difference in the danger involved in the several diseases simulating acute hydrocephalus, and that disease itself, that it becomes a matter of extreme importance to distinguish one from the other during life. The diagnosis is often difficult in the earlier stages; yet I think it can generally be made. With regard, however, to the recognition during life of the different forms of acute hydrocephalus itself, the difficulties are extreme; but they have been separated in the *post mortem* room, and the task will become easier now. The diseases which simulate acute hydrocephalus are three—viz., the simple convulsions of childhood, typhoid fever (the infantile remittent of the old authors), and what is usually called false hydrocephalus—that condition of insensibility accompanied by convulsion which occasionally supervenes upon protracted vomiting, diarrhoea, or starvation.

Now, if we compare the case which I have made the text of this lecture with each of these, and examine the evidence carefully, we shall be able to see at once that it presents essential points of difference from all of them. First, then, the case was not one of simple infantile convulsions, due to reflex irritation, without any primary lesion of the encephalon. The age of the patient, when the irritation of teething or intestinal disorder readily stirs up excited action of the cerebro-spinal centres, alone favours such a view: but there were no signs of teething, and, indeed, the child did not cut a tooth until five months afterwards. Then, again, in infantile convulsions, the attack comes on suddenly, and is not preceded by a long train of previous symptoms. The convulsion is the first, or one of the first, symptoms. This is followed by a period of insensibility of short duration only; unless, as sometimes happens, the convulsive seizures succeed each other at short intervals, when consciousness may not return for a considerable time. Yet this insensibility is not in itself of a persistent kind; its continuance is due to the constant renewal of the comatose state by successive attacks of convulsion. The stupor is due to the convulsion; it follows it and passes off shortly afterwards, unless reinduced by a recurrence of the exciting cause—viz., by a fresh convulsion. Since, then, in the simple convulsions of childhood the cerebral symptoms are not indicative of any primary lesion of the encephalon, but depend solely upon the condition of the brain, produced by the epileptiform attack—and in the case before us, as you will remember, no convulsion occurred until three weeks after the child was first observed in a semi-comatose condition—it is clear that the cerebral symptoms could not be due to this cause, but to some primary affection of the brain or its membranes. As additional proof, if any were needed, I may point out to you that persistent, rigid retraction of the neck, incessant vomiting, and intolerance of movement, are not met with in simple convulsions. It is equally clear that the disease was not

typhoid fever. The symptoms of this disease are often greatly modified in children, and at the outset closely simulate the invasive stage of acute hydrocephalus. There is often little or no purging at first, or indeed throughout; and the characteristic spots are not uncommonly entirely absent. The disease is usually less severe, and runs a shorter course. The absence of diarrhoea and of rose spots would not, then, be sufficient in themselves to prove this case was not one of abortive typhoid; but the dark offensive scanty stools, the moist, thickly coated tongue, the continued vomiting, the retraction of the neck, the blindness, and the duration of the attack, are in striking contrast to the light watery motions, the dried tongue, the transitory sickness, the muscular relaxation, and the definite shorter course of the enteric fever: and then typhoid is, as far as I know, never met with in the first year of infancy.

The remaining disorder, which might be supposed to have deceived us, is that state of convulsion and semi-consciousness which sometimes follows diarrhoea, or other exhaustive drain upon the system. It is due, no doubt, to anæmia of the brain and cord, and is closely allied to the convulsions and paralysis which result from excessive bleeding or extreme starvation. That such was not the condition in this case, is at once shown by the well-nourished state of the patient, and by the history. Neither anæmia nor wasting existed, nor had there been any antecedent flux or malnutrition to produce them, for the vomiting was not sufficiently constant and was of too short duration to have given rise to any serious results of this kind.

[To be continued.]

REMARKS ON THE MECHANICAL DILATATION OF THE CERVIX UTERI.

By LOMBE ATTHILL, M.D. Dubl.,

Fellow and Examiner in Midwifery in the King and Queen's College of Physicians of Ireland; Obstetric Physician to the Adelaide Hospital, Dublin.

THE practice of dilating the cervix uteri, in the treatment of certain affections of that organ, is now so common, that too great importance can hardly be attached to the consideration of the best method of accomplishing it. Dr. Matthews Duncan has, therefore, done good service in publishing his scientific researches thereon; but from the conclusions which he deduces from them I must most emphatically dissent on two grounds—namely, the results of his own experiments and of experience. I think I may assume that, the more nearly we can imitate the action of Nature in the production of any artificial process for the cure of disease, the more safe and satisfactory will be the result.

Dr. Matthews Duncan, in the able article which has just appeared from his pen on the subject under consideration, says that the uterus during pregnancy expands, by what he in a few lines before terms “a vital action,” “so as to remove the smallest active pressure on its contents”. To this “vital action” he attributes the effect produced by the sea-tangle tents. To quote his own words: “In ordinary cases the tangle does not expand by exerting force, but, a length of time—at least some hours—being allowed, through some vital change in the tissues touched. The superiority, then, as an expander in ordinary cases of a tangle over a metallic tent lies chiefly in this—that, while the metallic tent cannot enlarge, the tangle-tent does enlarge, and follows the expanding tissues and produces a further expansion in the same, but now also continuous, manner as the limited amount of expansion produced by a metallic tent is effected.” To allow the metallic tent to arouse this vital action, “hours at least” must, by Dr. Duncan's own showing, be allowed. But Dr. Duncan in a preceding paragraph says, in speaking of the bougie, “here the shortness of time used excludes vital relaxation or yielding, or nearly so”. Therefore Dr. Duncan, in recommending the use of the bougie in gynaecological practice, deliberately recommends that process which, by his own showing, directly violates the laws which Nature follows in dilating the cervix.

In this city, the practice of dilating the cervix uteri by means of tangle-tents has been adopted almost universally for several years past. In my own practice no unpleasant symptom has followed their use, nor have I yet met with a case which resisted their action. I do not mean to say that serious consequences may not follow their use, because, where predisposition to disease exists, the most trifling interference may be followed by such; but in a tolerably extensive practice such have not occurred to me, and I think I may safely infer that such cases are rare. In America, however, where dilatation of the cervix by graduated metallic bougies is extensively practised, serious and even fatal consequences are not uncommon; nor can that be a matter of wonder when it is borne in mind that, “the vital action” being disregarded, a rigid unyielding muscle is forcibly expanded in a few moments. That rupture of some of the muscular fibres takes place, we may assume as most probable; that the peritoneal covering of the uterus is injured in some cases,