seems certain. Thus Dr. Thomas of New York some time since published the particulars of a case in which, after the introduction of three of Peaslee's dilators, fatal peritonitis followed.

One other point in Dr. Duncan's paper demands notice. I have ever taught that no actual force should ever be used in attempting to introduce the uterine sound; and I believe the most serious consequences will ensue if, following the practice of Dr. Matthews Duncan, practitioners unskilled in the use of the instrument exert a force capable of bending it. I introduce the uterine sound daily, and have done so for years, but never yet perceived the most trifling bending. I would repeat here what I have already said in my published *Lectures* (p. 11): "If any obstruction occur, it should be overcome by gentle continuous pressure; force must not on any account be used, lest injury be done to the uterine walls. Better far that you should never introduce it, than that you should run the risk of injuring the uterus, and perhaps cause a fatal result by doing by force what should only be accomplished by tact."

ILLUSTRATIONS OF THE SURGICAL USES OF THE PNEUMATIC ASPIRATOR.*

By T. R. JESSOP, F.R.C.S.,

Surgeon to the Leeds General Infirmary, and Lecturer on Surgery at the Leeds School of Medicine.

AMONGST the many valuable additions which have been made within recent years to the means whereby the physician or surgeon is enabled to diagnose and treat disease—some of which, indeed, such as the stethoscope, the ophthalmoscope, the laryngoscope, and the thermometer, have become indispensable—I know of none which, in my own hands, has proved more useful—the loss of which I should feel a greater deprivation—than the instrument which I have selected as the subject of my communication; viz., Dr. Georges Dieulafoy's Pneumatic Aspirator. Invaluable, however, as it has proved in the hands of many, and widely as it is now known, at least amongst hospital surgeons, it is my daily experience to find that medical practitioners in general have not as yet been made fully alive to the great disadvantages which they labour under from the want of a thorough appreciation of the many and varied purposes which the aspirator has been found to fulfil.

It is with the twofold object of exciting the interest of such of my brethren of this Branch as have not yet had an opportunity of seeing for themselves, or of hearing from others, the very satisfactory results achieved by means of the aspirator, which are daily to be witnessed in the wards of our hospitals, and of eliciting a discussion on the part of those experienced in the use of the instrument, that I have determined to recount a few examples showing various conditions in which aspiration has been resorted to with more or less success. But before proceeding with the relation of my cases, I must occupy a few moments, first, upon the general subject of aspiration; and, secondly, upon a brief description of the instrument which we now employ.

Every medical man well knows how extremely difficult it is, under some circumstances, to decide upon the presence or absence of fluid; and, again, upon the nature of a fluid which is clearly proved to exist. How many anxious hours I myself have spent in vainly trying to solve these two questions! We have been in the habit of relying, in such cases, partly upon probabilities, partly upon that *tactus eruditus* which every surgeon strives to acquire, but which can only be acquired through long and patient practice; and occasionally upon evidence obtained from the introduction of small or, as they are termed, exploring trocars—instruments which are, to use Dr. Dieulafoy's words, at once too large and too small: too small to permit the passage through them of viscid or flaky fluids, and too large to allow their safe introduction into vascular or sensitive organs.

Having repeatedly observed how unimportant a lesion is produced by the insertion of the fine nozzle of the hypodermic syringe, and also the impunity attending the introduction of acupuncture needles, even deeply, into the substance of important and highly organised structures, Dr. Dieulafoy conceived the idea of creating a vacuum in connection with an extremely fine tubular needle, so as to exercise a powerful suction upon any fluid into which the needle should be introduced. This principle he embodied in his instrument, the pneumatic aspirator. And it became clear that not only could an amount of fluid be removed by this means sufficient for diagnostic purposes, but that, by a continuance of the process, the cavity might be entirely emptied of its contents, and so, in cases where the presence of the fluid constituted the sum of disease, a permanent and ready cure might be effected.

As is the case with all inventions, many improvements have been made

upon the original instrument. The one which I now show you is the latest issued by Messrs. Weiss and Son, of the Strand, London; and, in my judgment, it possesses decided advantages over all others which I have seen.* It consists of a brass syringe, capable of holding five ounces of fluid, fitted with an air-tight screw-piston, and having its nozzle guarded by a tap. Attaching the nozzle to the bottom of the syringe is a short tube of glass, which enables the surgeon to ascertain the quality of the fluid as it passes into the body of the instrument. To one side of the instrument, in its whole length, is applied a fine hollow glass tube, which communicates at its lower extremity with the interior of the syringe, and is shut off from the external atmosphere, at its upper end, by means of a tap. This glass tube is marked at regular intervals in half ounces, for the purpose of demonstrating the quantity of fluid contained in the syringe at any given time; and forms, besides, a means of exit, whereby the syringe can be emptied of its contents when, during an operation, it is desirable not to disengage the nozzle. Several tubular needles and trocars, some of which are of extremely fine calibre, made accurately to fit the nozzle of the syringe, either directly or through the medium of an elastic tube, complete the instrument.

In describing how to work the aspirator, I cannot do better than make use of Dr. Dieulafoy's own words, altered only so as to apply to the improved instrument. He says: "In order to produce a vacuum within the pump, it is necessary first to close the two taps, and then to screw up the piston. The vacuum is thus obtained as a preliminary measure, and the operator is in possession of a powerful aspirator, ready to be used when the proper time arrives." "Let it be supposed that we wish to examine an effusion into the cavity of the pleura. The tubular needle No. 1 or No. 2 must first be introduced into an intercostal space; and, when it has penetrated the tissues for about a third of an inch, it must be connected with the pump in which the vacuum has been established, either directly or through the medium of a caoutchouc tube. This done—and to this point I desire to call special attention—the tap must be opened between the needle and the vacuum, and the needle pushed gently forward. We may thus slowly traverse the tissues, so to speak, with the vacuum in hand, until we discover the effusion. The eye of the operator should be directed to the short glass tube between the needle and the pump; and, at the moment when the needle enters the liquid, the latter rushes forcibly into the instrument. The diagnosis is at once complete, the manœuvre is absolutely harmless, and the desired object is attained."

Such is the method of employing the aspirator which I have myself usually adopted; and the directions here given apply to all cases, medical and surgical, in which its use is indicated, equally with the example selected by Dr. Dieulafoy for his description. As a precaution against the introduction of septic materials, I always take care that the needle or trocar has been dipped overhead in carbolic oil. I have said that in the aspirator we possess a valuable aid both to diagnosis and to treatment. I will now go on to make a few selections from the numerous cases in which I have resorted to its employment.

And first, I will speak of its use for purposes of diagnosis. There are few groups of diseases which present greater difficulties, or which tax the skill of the surgeon in a higher degree, than *abdominal tumours*. Not many of us can look back upon our experiences without calling to mind numerous instances in which grave mistakes have been made, leading to pernicious treatment, and destroying the confidence of our patients both in ourselves and in our science. That in the aspirator we possess a powerful means of elucidating some at least of these cases, is sufficiently shown by the following examples.

CASE I.—On August 30th last, a woman, aged 46, was admitted under my care into the Leeds General Infirmary with an abdominal tumour of twelve months' growth, and equal in size to a seven months' pregnancy. The tumour was somewhat fixed towards the pelvis; several hard movable nodules could be distinguished through the abdominal walls; it was doubtful whether fluctuation existed or not. The pelvis was found, on vaginal examination, to be filled posteriorly by a firm mass, and the os uteri could not be felt, being apparently pushed up above the pubes, high out of reach of the finger. Menstruation had for some years been very irregular. My colleagues examined the case with me, and we all felt doubtful as to its nature. On September 21st, aspiration was made through the abdominal walls with the result of extracting four ounces of a thick glutinous brown-coloured fluid, such as is found only in ovarian cysts. The nature of the disease having been thus established beyond doubt, a fortnight later I performed upon her the operation of ovariectomy.

* I have received a note from Messrs. Weiss and Son within the last few days, informing me of a still further improvement which they have been able to effect in the mechanism of the instrument: by which they avoid all contact of the aspirated fluid with the syringe, and are at the same time better able to regulate the amount of vacuum created.

CASE II.—One day last week it was intimated to me that Dr. Allbutt wished me, as his surgical colleague, to examine a patient in his ward, a married woman, aged 34, with a view to decide upon the propriety of recommending the operation of ovariectomy. The enlargement was first suspected early this year, and was attributed for some months to natural causes. I found the abdomen distended to about the size of a full pregnancy, by a round, centrally placed, smooth tumour, in which a not very distinct fluctuation could be felt. On vaginal examination, the os uteri was felt high up posteriorly, small, and round; the cervix had disappeared; no movement of the os could be detected by displacing the tumour from side to side; and a Simpson's sound introduced into the uterus showed the cavity of that organ to measure an inch and three-quarters only in depth. The central position, the shape, and the general character of the tumour, the absence of cervix uteri, the duration of the symptoms, taken together, were sufficient to throw a little doubt upon the true nature of the case. On appealing to the aspirator, however, the ovarian character of the tumour and the correctness of Dr. Allbutt's diagnosis were at once determined, by the extraction of a syringe-full of thick stringy gelatinous fluid.

CASE III.—In July of this year, I was asked by Mr. Booth, of Mirfield, to see, with him, an elderly female, the right half of whose abdomen was filled with a prominent mass, the origin of which she attributed to her daughter having accidentally trodden upon her in getting out of bed a few months before. The increase in size had been gradual, and had been attended by considerable pain and some febrile disturbance. On percussion, the tumour was dull, and on handling, an indistinct fluctuation was made out. A few days afterwards, I withdrew by means of the aspirator several ounces of pus, and was thus enabled confidently to pronounce the case one of cellular abscess, and to confirm the patient's own opinion as to the origin of the disease.

CASE IV.—Not many months ago, a man, aged about 40, of weak intellect, and therefore unable to give any reliable history of himself, was sent from Tadcaster and admitted into the Leeds Infirmary under Mr. S. Hey, with complete retention of urine, and his bladder distended so as to reach above the umbilicus. Our resident medical officer, Mr. McGill, at once passed a No. 8 silver catheter, and drew off forty ounces of fetid urine—as much as would flow. And yet the distension was not wholly removed; above the pubes, a round fluctuating swelling remained, which, from the rectum, was found to communicate with a soft elastic bulging there distinctly felt. The aspirator was applied from the rectum, and by it twenty-five ounces of fluid were quickly pumped out, whereupon all tumour was found to have disappeared. The fluid was clear, had a specific gravity of 1004, and, besides large quantities of common salt, contained numerous hooklets and undamaged echinococci. The retention ceased, but the patient died a week afterwards from the inflammatory changes induced by the long continuance of the obstruction. At the *post mortem* examination, the hydatid disease was thought to have occupied the prostate itself, and similar disease was found both in the liver and omentum.

I will finish my illustrations of the diagnostic value of the aspirator by briefly relating a very important and not less interesting case which occurred recently in the hospital practice of my colleague, Mr. Wheelhouse, and which I had the opportunity of seeing in consultation with that gentleman.

CASE V.—J. M., aged 30, a striker by occupation, had been the subject of a right inguinal hernia for four months, and had never worn a truss. The bowel had frequently from time to time descended, and had always been returned without difficulty. About breakfast-time, on October 2nd, 1872, whilst he was at work, the hernia suddenly protruded with pain. Finding himself unable to reduce it as usual, he went home. Symptoms of strangulation soon supervened, and his sufferings became greatly increased. It was not, however, until the evening of the 4th that he sent for his medical attendant, who ordered his immediate removal to the infirmary. When seen on admission, he was vomiting frequently; his abdomen was highly distended, tympanitic, and tender on pressure; his countenance betrayed anxiety, and his pulse was small and thready. The right inguinal region was occupied by a swelling of the size of a hen's egg, not very tense, but upon which taxis produced not the least impression. Opium was administered, and taxis again failed. He was then placed under the influence of chloroform, and an attempt was once more made to reduce the rupture, but in vain. The finest of the needles of the aspirator was now introduced and attached to the syringe, whereupon two ounces of bloody serum were removed, and the swelling vanished altogether. The finger could now be passed without obstruction through the inguinal canal as far as the internal ring. Under these circumstances, it was decided to postpone any further proceedings until the following morning. Vomiting occurred at intervals during the night; but at the morning visit, the patient stated that the rupture had again descended, and that he had

been able to reduce it himself. It was not until the morning of the 6th that all vomiting ceased; but from the first there was a gradual and steady amelioration of his symptoms, and, finally, he has made a perfect recovery.

It is by no means an easy matter to state, with any degree of certainty, the actual order of events in this most fortunate case. It may be safely concluded, however, that the use of the aspirator precluded the necessity of following that good old rule—"when in doubt, operate."

I will now go on to mention a few instances in which aspiration has been resorted to as a means of treatment. Two of the earliest occasions upon which I made use of the instrument were in cases of advanced chronic hydrocephalus.

CASE VI.—In the first, a child, fifteen months old, I removed on May 18th, 1871, by means of the finest needle passed through the anterior fontanelle, six ounces of fluid; and on the 22nd, twelve ounces. The child died on the 23rd in convulsions.

CASE VII.—The second child was only three months old when first brought under notice. On July 22nd, 1871, I removed five ounces of fluid; on the 25th, three ounces; on August 1st, an ounce and a half; and subsequently, on seven occasions, at intervals of a few days, one ounce of fluid was removed. This case terminated fatally, also by convulsions, on September 13th.

A careful *post mortem* examination was made in each of the two instances, with the special view of discovering the immediate effects of the introduction of the needle, but no trace whatever could be detected either in the membranes or in the brain-substance.

Soon after I became familiar with the aspirator, it occurred to me that it was not unlikely to prove useful as an aid, under certain circumstances, to the reduction of strangulated hernia. I thought that, if the sac, for instance, contained a large amount of fluid, the withdrawal of the fluid might enable the surgeon to apply his reducing force more directly, and, therefore, more efficiently, upon the solid constituents of the hernia. And, again, if the strangled intestine were emptied of its fluid and gaseous contents, its reduction by taxis might be facilitated. In five examples of this affection, all presenting conditions apparently favourable to the proceeding, and in all of which taxis under chloroform first proved insufficient, I have practised the use of the aspirator, in the full hope of being able to avoid the major operation; but, hitherto, my efforts have not met with success. In all the five, herniotomy was finally resorted to; and it is interesting to observe that, in none was there any sign to show where the needle had penetrated. Cases have lately been referred to in the journals in which success has been achieved by operators abroad. I am still hoping that a further trial may lead to more satisfactory results in my own hands.

Some few months ago, several gentlemen showed great anxiety to obtain the credit for themselves, or for their friends, of having been the first to tap the intestines in cases of great flatulent distension. I remember the late Mr. Teale, in 1860, whilst introducing an acupuncture-needle twenty or thirty times through the abdominal walls in a case of excessive tympany following herniotomy, remarking to the students around him that, in his early days, he had been accustomed, under similar circumstances, to use the trocar after the manner of veterinary surgeons; but that, of late years, he had preferred the acupuncture-needle, as being at the same time less dangerous and equally efficient. Calling to mind this advice, I employed the aspirator on several occasions in May 1871, to the great relief of an elderly man, whose leg I had amputated by the supracondyloid method, and who in his last two or three weeks suffered in an unusual degree from excessive flatulent distension. On *post mortem* examination, a few red points, on both the parietal and visceral peritoneum, were thought to indicate the spots at which the needle had entered.

The instances are now numerous in which I have employed the aspirator, with decided advantage, in the emptying of chronic abscess. I will content myself with relating one example, by no means a solitary one of its kind.

CASE VIII.—In May 1871, I was consulted by Mr. T., aged 40, who for nearly six years had been suffering from excessive pain in the lumbar and lower dorsal spine, and at the top of whose right thigh a swelling had for nine months been gradually increasing. The case, as it presented itself to me, was, in short, one of marked angular curvature at the junction of the dorsal and lumbar vertebræ, accompanied with a psoas abscess, which, when he stood erect, was about as large as an orange. On June 10th, I removed seven ounces and a half of pus by aspiration, draining the abscess as completely as possible. In a few days fluctuation returned, and the swelling gradually increased. On June 28th, the aspirator was again applied, and three ounces and a half of pus removed. From this time the abscess may be said to have been cured; for although, on February 7th this year, I was induced once again to introduce the needle, from a feeling of suspicion that matter

had reformed, I was unable to procure a single drop. At the present time, no difference whatever can be detected between the two sides, and he is quite free from all pain. The curve, of course, still remains; and, as a matter of precaution, I am continuing to enjoin a large amount of rest in the recumbent position, and an entire freedom at other times from such movements as would be likely to affect injuriously a spine which has undoubtedly been extensively diseased.

There are few conditions which more frequently call for the attention of the surgeon than the various forms of joint-effusion; and, whilst some of these are of a transient character, or readily yield to appropriate treatment, there are others, the nature of which it is to be obstinately persistent; or, again, in which the effusion is but the precursor, if not indeed a chief factor, in the production of those processes which, if constructive in their tendency, end in ankylosis; but, if destructive, lead only to excision, amputation, or death.

It is within my own recollection—nay, I believe the practice still lingers in the dark corners of our profession—that surgeons were in the habit of spending month after month—aye, and in some instances, year after year—in leeching, blistering, burning, compressing, mercurialising, and drugging their patients, with the sole object of procuring absorption by a membrane, which, it may be, had ceased to be capable of absorbing, and which, at all events, obstinately refused to take the desired turn; or perhaps, goaded by the constant irritation from without, openly rebelled, as it were, in the form of suppuration and destruction.

Whilst no means were too severe to be applied to the exterior, it was a cardinal doctrine that to interfere with the interior of a joint—except in cases of acute suppuration—was an unsurgical and unwarrantable proceeding. The sacred character of synovial membranes, like many other phantoms, has vanished before advancing science. Dropsical joints are now treated by tapping, just as are the peritoneum, the pleura, or the pericardium. A couple of years ago, I published in our JOURNAL the history of a series of knee-joints which I had successfully treated, when in an early condition of inflammation, by *free incisions*, made under the protection of an antiseptic spray. And now that we possess the aspirator, we shall hardly be disposed, I imagine, to consume as many days as we formerly did months, in the endeavour to disperse a fluid whose presence is not seldom the sum total of disease.

The following are selected from a considerable number of cases which have been similarly treated.

CASE IX.—A gentleman, aged 21, in falling from his bicycle, struck his left knee a violent blow against the corner of a wall. Faint, and in great pain, he limped, as best he could, to his home, where I saw him on the following day, June 8th this year. The joint was distended from effusion within. He suffered but little, and could bear manipulation well. Rest in bed, an evaporating lotion, and an occasional purgative were prescribed. This treatment was duly persevered in for a month, without producing any perceptible change in the condition of the joint.

On July 5th, I applied the aspirator, making use of the smallest trocar, and drew off seven ounces of fluid, which consisted of little else than blood. It was the first time I had met with blood in a joint under similar circumstances, and I felt somewhat anxious as to the result. The knee, although emptied of fluid, looked decidedly thicker than the sound one, but was neither painful nor tender. The limb was placed upon a straight splint, and the evaporating lotion was continued. There was no return of fluid, and, indeed, nothing of an untoward nature at any time arose. On August 6th, he was allowed to get up, with the joint supported by a plaister of Paris bandage. In six weeks the bandage was removed; and at the end of September he could walk well—a little stiffness and a slight thickening of the joint alone remaining.

The following is a somewhat similar case.

CASE X.—F. M., aged 40, a dyer by occupation, was admitted into the Leeds Infirmary on September 16th, 1872. On August 31st, whilst attempting to climb over a high wall, he fell upon his left knee. Although in very great pain, he walked home—a distance of about two hundred yards—and on arriving there, he found that his knee was much swollen. Nevertheless, for three days he continued at his work, and only then desisted because of extreme suffering. After a few days' rest the pain ceased, but the swelling remained as before. Lotions and poultices were applied, and during two days the joint was painted with tincture of iodine. On admission, there was decided, though not excessive, swelling of the knee, and fluctuation was distinct. On September 18th, by means of the aspirator, I withdrew three ounces of a fluid deeply tinged with blood. The limb was placed upon a straight splint, until September 28th, when a plaister of Paris bandage was substituted, and he was made an out-patient. On October 14th, the bandage was removed at his urgent request, and, as recovery was apparently complete, he was permitted to return to his employment.

Satisfactory as were the results in the two cases just related, I think I am justified in attaching more importance to the two following; for, if my belief in the pyæmic origin of each be well founded, it would hardly be possible to exaggerate the value of a means whereby an issue so successful was accomplished.

CASE XI.—Joseph J., aged 21, a coal-miner, was admitted into the Leeds General Infirmary on September 19th, 1871, on account of acute periostitis with suppuration affecting the right humerus, of four months' duration. Incisions were made for the evacuation of pus. On October 14th, after several chills, not amounting to actual rigors, followed by feverish symptoms, and a rapid rise of temperature, he complained of great pain in the left hip-joint, which was intensified on the slightest movement. The application of the long splint afforded some relief; but the surgical fever continued without abatement, and the uniformly great difference between the morning and evening temperatures indicated the presence of acute suppuration. On October 27th, the left knee became suddenly swollen and acutely painful. On November 1st, this joint was aspirated, and two ounces of a clear fluid withdrawn—an operation, so far as that particular joint was concerned, which gave immediate and permanent relief. The general condition of the patient, however, was unimproved; and, as the affected hip continued still the source of sleepless nights and weary days, the aspirator was applied to this joint also on December 10th, with the result of extracting nine drachms of thick pus. The operation acted like a charm; his suffering was at once removed, and soon the balance of healthy action returned. On January 12th, 1872, he was discharged able to walk well, and with free movement in both hip and knee.

CASE XII.—The second case may be related in few words. A man, aged 52, was in the Leeds Infirmary suffering from old stricture of the urethra with many perineal fistulae. I had admitted him for the purpose of his undergoing the operation of perineal section. When he had been in the ward a few days he was seized with rigors, which were followed by inflammatory fever, and pain with some swelling of the left knee-joint. On the day after his seizure, fluctuation being decided, I removed by the aspirator an ounce and a half of turbid synovia. The pain at once ceased; and a week afterwards, he was sent home for awhile to recruit his health, lest an operation then performed should be followed by more serious consequences.

One case more, and I have done. It was the second of four I have had of its kind, and is selected now chiefly on account of its brevity.

CASE XIII.—On December 26th, 1870, a man, aged 38, was admitted under my care into the Leeds Infirmary with his bladder fully distended, and totally unable to pass a drop of urine. The retention had lasted for fifty-four hours, and several efforts had been fruitlessly made to enter his bladder. On my visit, he had already been placed by our resident medical officer in a hot bath, and opium had been freely given. A gentle attempt at catheterism sufficed to demonstrate a closure of the urethra a little beyond the bulb, and was followed only by the escape of a few drops of blood. Without further delay, I plunged the finest needle of the aspirator straight through the abdominal walls into the bladder, and, attaching the syringe, drew off thirty-eight ounces of urine. In about six hours, dribbling by the natural passage began, and, after a few days, I was able to commence the treatment of his stricture under circumstances not less favourable than if retention had never occurred.

EPITHELIOMA OF THE UPPER JAW.

IN the JOURNAL of Nov. 23, there is a short notice of a case of epithelioma of the upper jaw, exhibited by Mr. Lawson at the Clinical Society; and, in the general discussion which followed, I see it reported that Mr. Hulke "thought it doubtful if the epithelioma originated in the antrum, as the antrum did not exhibit any bulging out of its anterior wall, or other symptoms which would have been expected. Probably the disease originated in the alveolus, and invaded the antrum."

I have recently operated on a case of epithelioma involving both superior maxillary and palate bones, with a large part of the soft palate. In my case, the alveolar portion seemed, so far as I could discover, primarily and chiefly affected; but here also there was a large mass as big as a hen's egg filling the right antrum, without any bulging of its anterior walls; and I have no doubt that the mass grew from the alveolar portion into the antrum, in the way which Mr. Hulke thinks probable in Mr. Lawson's case. In my case, I did not discover the mass in the antrum until after I had removed it, together with the greater part of both superior maxillary and palate bones and soft palate.

As I hope presently to furnish the details of this interesting case, I refrain from further comment, and only call attention to it at this moment because of its similarity in some respects to Mr. Lawson's case.

NELSON C. DOBSON, F.R.C.S., Clifton.