of the City of Dublin, and possibly with the Dublin Ballast Board which has charge of the Port of Dublin. The question as to the constitution of an urban authority, and the area of its jurisdiction, should be determined by the Local Government Board (possibly by provisional order in large towns, and summarily in smaller places).

As to the expense of carrying out these proposals, we believe the total annual cost of the whole sanitary organisation, as proposed in the foregoing remarks, would amount to little over £100,000 a year. The commissioners at the head of the department should receive from £1,500 to £2,000 a year; the medical inspectors, about £1,000; the dispensary officers in rural and but sparsely populated districts, from £50 to £75 per annum, and in urban from £75 upwards to £200 or £300 for the smaller, and from £500 to £1,000 for the larger towns. We do not wish to suggest that our scheme is perfect; but no systematic organisation has up to the present been proposed, and we believe that, without a thorough, systematic, complete, and properly paid organisation to start with, no good result can be expected from further sanitary legislation for Ireland.

ON THE MODE OF DISINFECTING BY HEAT.*

By W. H. RANSOM, M.D., F.R.S., F.R.C.P.,
Physician to the General Hospital, Nottingham.

As it is my desire to treat this question as a practical one only, I assume your assent to the proposition that heat is a disinfecting agent of so much value as to justify public authorities in spending money to secure the necessary mechanical appliances. To this I venture to add, as an opinion which will, I think, receive your general assent, that heat is, for a certain group of things, the best and almost the only known trustworthy disinfecting agent. Nor can I here, without violence to the memory of an able man who has passed away—I mean Dr. Henry of Manchester—omit to point out that almost all our knowledge of the facts upon which we rest the proposition to which I have assumed your assent, is due to that gentleman. He alone, so far as I know, has had the courage to make the experiments so much needed; and he has apparently proved that a temperature of 200 deg. Fahr., continued for not less than one hour, destroys the activity of the virus of scarlet fever.

My attention was first drawn to this matter about eight years since, when, being asked by a public body to advise them as to the best form of hot closet for disinfecting purposes, I was unable to find any one in use which I could recommend. I therefore commenced some experiments with the view of suggesting a pattern to be tried; but the engineer who then kindly assisted me died, and the matter was put aside until 1870, when the occurrence of scarlet fever in my own family recalled my attention to this urgent public want.

Experiments made at home in 1870 with a hot closet, having a capacity of about twenty-five cubic feet, constructed upon the oven principle, heated by gas outside, and having its surfaces partly covered by a hot-air flue and partly backed with non-conducting material, led me to the conclusion that this form of hot closet could not be recommended. I was, however, enabled from these observations to form provisional opinions as to the temperature which might be safely used, the time that goods should be exposed, the need for ventilation in such a chamber, etc. I will not take up your time by stating in detail the results of these experiments, because, from the imperfection of principle in the closet, very few of the results were sufficiently exact. However, I found that clothes and bedding might be exposed to a heat of 250 to 260 deg. Fahr. for an hour and a half in such a chamber without much injury. As chambers made on this principle are much used, I will here briefly state the imperfections which were found out in working.

a. The chamber cannot be long maintained at a steady heat, however carefully you attend to the adjustment of the gas-tap; and, were it heated by coal, I cannot think it could be so well regulated as by gas.

b. The temperature varies enormously in different parts of the chamber at the same time.

c. Being imperfectly ventilated, its drying power is insufficient, and the heat therefore penetrates slowly into the interior of beds, etc.

I decided, therefore, to endeavour to construct a new one, which should fulfil the following conditions.

1. The temperature should at all times and in all parts of the chamber be within certain known limits; and it should be plain at a glance that they are not exceeded; the maximum being determined by the singeing-point of the materials of clothes and bedding, and the minimum lying as close as possible to the maximum, but always ten or twelve degrees above 212 deg. Fahr. Thus one might feel assured that clothing put into such a chamber really was exposed to a sufficient heat, and yet not injured by too much, which would seriously deter the public from using it.

2. The temperature should be easily maintained within the required limits, for periods of time sufficient to ensure adequate penetration of heat into the interior of bad conductors, such as beds, etc.

3. The chamber should have a current of heated air passing through it, so as to favour penetration of heat by drying.

4. It should be so moderate in cost and so simple in use, that small institutions might avail themselves of it, and a fairly instructed nurse manage it.

5. It should be self-regulating.

Accordingly, in the winter of 1871-2, a hot closet was erected, under my directions, at the General Hospital, Nottingham, which to a great extent did what was desired to be done. In this contrivance, the heat is obtained by burning gas with smokeless flames, and conducting the products of combustion, mixed with the heated air, by means of a short horizontal flue, into a cubical chamber through an aperture in its floor, and out of it by a smaller aperture in its roof. Fixed thermometers show the temperature of the entering and outgoing currents, which represent the maximum and minimum temperatures of the chamber. A self-acting mercurial regulator, adapted to this stove from Kemp's regulator, maintains the temperature of the entering current at any required degree. A thick jacket of felt and wood economises the heat, and makes it steadier. A valve at the outlet of the chamber, together with another at the inlet of the furnace, enables the maker to regulate once for all the speed of the current passing through the chamber—an item of great importance in the endeavour to keep the temperature of the outgoing current as high as possible. To this, if it be desired, may be added a conical short chimney at the outlet, in which three or four small fish-tail burners may be lighted. By this method, any active particles of poison which might, in the earlier moments of the operation, be carried up in the current, would be burned. Dr. Henry anticipated objections of this kind, and used in his apparatus a similar contrivance to that which is now employed in Fraser's disinfecting closet.

For about a year and a half this closet has been in use with but few interruptions, and, I think I may venture to say, has rendered important services to the hospital, especially during the recent prevalence of small-pox, when a number of cases were admitted to our fever-house. Whenever the fever-house is occupied, the hot closet is at work night and day, and ready at any moment for use. The sister in charge managed it without difficulty after a little instruction which I gave her. Although it has a cubic capacity of twenty-five feet only, and is chiefly used for the clothes of patients, it has sufficed to disinfect blankets and the contents of beds and pillows. Its average consumption of gas is nine cubic feet an hour, at a cost, assuming gas to be 4s. per 1,000 feet, of 6s. a week, counting seven days a week, and twenty-four hours a day.

Records of its work have been kept, showing the maximum and minimum temperature, the consumption of gas, and the pressure of gas at various hours. From these I am able to say that the heat can be maintained for weeks at a stretch without any greater diurnal variation than 5 deg. Fahr., which is usually due to the increase of pressure at the gas-works in the evening.

The average reading of the maximum thermometer is 250.5 deg. Fahr.; that of the minimum is 232 deg. Fahr., which I should like to see 10 deg. Fahr. higher. Nor do I doubt the possibility of making it so in any future closet which may be erected upon this plan.

Thus, it may be said, the five conditions stated as desirable have been fairly well fulfilled by this stove. When a large experience shall have taught us more, I do not doubt that much better results may be achieved by improvements in the details of construction of hot closets on the principle now described.

The chief advantages of this closet are, its moderate cost, its efficiency, cleanliness, constant readiness for use; its capability of being managed, without the cost of an engineer or stoker, by an intelligent nurse; and the facility which it offers for maintaining a steady known heat for the length of time necessary to ensure deep penetration into bad conductors.

The only disadvantages which I have noted are: a. The necessity of having occasionally (perhaps four times a year) the services of an intelligent workman to look at and adjust the heat-regulator if necessary;

* Read before the Public Medicine Section at the Annual Meeting of the British Medical Association in London, August 1873.

6. The action, which, I think, is occasionally though rarely seen, of the acid products of combustion upon some of the most delicate colours. This I will show you, and you will see how trivial it is. It would not be difficult to overcome this objection.

Before proceeding to draw up a card of instruction for the guidance of the sister in charge, I thought it prudent to repeat observations by the acid of this chamber on the singeing-point of various textile fabrics, and on the time required for penetration of heat into beds, etc. First, as to the singeing-point; here are tables recording these observations, not yet complete. I will sum up the results, so far as I have gone.

White wool, cotton, linen, silk, and paper, may be heated to 250 deg. Fahr. for three hours without apparent injury, although the wool will show a faint change in colour, especially when it is new. Perhaps this change is not more than the change which new white flannel undergoes when it is washed. The same may be said of dyed wools and printed cottons, and of most dyed silks; but there is one kind of white silk which easily turns brown by this heat, and pink silks of some kinds are also faded by it. The same or even slightly lower temperature will, if continued for a longer period—seven or eight hours—slightly change the colour of, but not injure, white wool, cotton, silk, paper, and grey linen. A heat of 295 deg. Fahr., continued for about three hours, more decidedly singes white wool, and less so grey and white cotton and white silk, white paper, and linen both grey and white; but does not, I think, injure their appearance materially. The same heat, continued for about five hours, singes and injures the appearance of white wool and cotton, grey linen, white silk and paper, some coloured fabrics of wool, or mixed wool and cotton, or mixed wool and silk.

These observations confirm the conclusion drawn from previous ones—that 250 deg. Fahr. was the best and most safe maximum heat to use. It is interesting to observe here that Dr. Henry (see foot-note in *Phil. Mag.*, vol xi, p. 22) had ascertained that cotton would resist this temperature. I was not aware, however, of this until the spring of this year.

It is noteworthy that the singeing of any fabric depends not alone upon the heat used, but also on the time during which it is exposed; and is influenced by the presence of other articles in the stove capable of giving up moisture. The maximum temperature which I find best for general purposes—viz., 250 deg. Fahr.—does not weaken the strength of cotton, silk, wool, or linen threads, even when continued for about eight hours; neither does it, so far as my experiments go, lessen the strength of textile fabrics composed of these yarns. This statement, however, refers only to articles which have been left aside long enough to have reabsorbed their normal hygroscopic moisture. A temperature of from 293 to 300 deg. Fahr., continued from three to five hours, weakens the strength of cotton-thread; probably also that of linen to a less degree—not silk, and not wool. But the fabrics composed of woollen yarn become dusty, and, I should think, would be weaker after washing. Neither they, nor yet linen fabrics, as in bed-ticking, are demonstrably weakened by this heat.

Turning now to the difficult question of securing the penetration of heat into the inside of beds, etc., I must again notice that Dr. Henry states that this is only a question of time; and so, in the main, I find it to be by observation. The accompanying table, abstracted from my notes, will show this. But the penetration of heat depends not only upon the time of exposure, but also upon the thickness of the bed, the amount of hygroscopic moisture it contains, the nature of the materials of which it is composed, and perhaps also their degree of compression, as well as on the temperature used. These conditions have to be borne in mind in working any disinfecting chamber.

No.	Name, thickness, and condition of sample.	Heat used. deg. F.	Time heated.	RESULTS.		
				Heat of inside. deg. F.	Loss of Weight.	Singed or not.
1	Horse-hair pillow, 5 in. thick; normal moisture	251-262 2 operations	8 hours	247	1-10th	Not
2	The same, nearly dry	247	2½ "	221	1-40th	Not
3	White blanket folded in 24 layers. 4½ in. thick; moist.	248	6½ "	214	1-12th	A little; not injured
4	Feather pillow, 5 in. thick; moist	241	7½ "	232	1-10th	Not
5	Flock pillow, 5 inches thick; moist	237-245 2 operations	23 "	252*	1 10th	Not
6	Horse-hair pillow, very moist, 5½ inches thick	293	3½ "	178	1-8th	Singed; injured
7	The same, much dryer	296	4¾ "	296	1-17th	Ditto
8	Flock pillow, 5½ inches thick; moist	293-300 2 operations	10½ "	280	1-10th	Ditto

* Perhaps an error in instrument. The inner and outer thermometers in this case may be taken as equal.

THE ACTION AND RELATIVE VALUE OF DISINFECTANTS.*

By J. A. WANKLYN, Esq.

If we heat the infectious material of, for instance, scarlet fever to a red heat, we destroy it. There can be as little doubt that, if we bring chlorine gas thoroughly into contact with infectious material, we destroy it likewise. If we boil it with oil of vitriol, or with permanganate of potash, we destroy it. Probably, too, if we soak it in concentrated carbolic acid, or treat it with excess of corrosive sublimate or arsenious acid, we render it inert. It is, indeed, highly probable that every kind of infectious material is capable of being rendered inert by thorough contact with any powerful chemical reagent.

But if the heat be only gentle, and if the chemical reagent be dilute, there is absolutely no reason for believing that, by the employment either of the one or of the other, we are so much as contributing towards the destruction of infection. There is a difference not only in degree, but even in kind, between the action of the same chemical when concentrated and when dilute. Concentrated sulphuric acid will convert cane-sugar into a lump of charcoal, but dilute sulphuric acid transforms it into dextrine and glucose, and, curiously enough, fits it for undergoing septic changes. So, again, very dilute bleaching powder has actually been found to favour the development of certain low forms of life; and Pettenkofer, as is well known, has found that germs whose development had been arrested by carbolic acid, start into life when the carbolic acid is still further diluted.

In the practical employment of disinfectants, the fact that dilution frustrates the action of a disinfectant has been very generally lost sight of. Attempts have often been made to disinfect the atmosphere. It is even said that, during the panic occasioned by the cattle-plague, the Commissioners endeavoured to disinfect the general atmosphere of the agricultural districts by turning cattle adrift with towels soaked in carbolic acid attached to their horns. I need not insist on the futility of such a proceeding; or, indeed, on the necessary futility of any effort to eliminate anything by chemical means from the general atmosphere covering our fields or occupying our streets. But it will probably not be quite needless to insist upon the impracticability of attacking the very limited atmosphere of a dwelling-house by chemical means. Certain very simple considerations will, however, suffice to throw the utmost doubt on the utility of endeavouring to purify air which has suffered contamination.

In a well known official memorandum, drawn up, I believe, by Professor Rolleston of Oxford, directions are given for the disinfection of a room with sulphurous acid. So much sulphur (the quantity proportionate to the size of the room) is to be burnt, and doors and windows are to be shut; and the memorandum winds up with the statement that, if a man be able to abide in the room for one instant whilst the disinfection is being carried on, then the disinfection is not to be depended upon. In other words, it is admitted that, not until you have put so much sulphurous acid into the air as to make it totally unfit to breathe, have you disinfected that air. The same certainly holds generally in regard to other agents; and, in short, we cannot hope to purify the air of a room by any chemical means without spoiling the air. It is, therefore, useless to try to disinfect the air. This is strikingly illustrated in reference to printed directions relative to the practice of disinfection. See, for instance, Dr. Wilson's little card, "Disinfectants and how to use them."

"Chlorine gas, poisonous and irritating to the lungs when in excess. For an occupied room, close fire-place, windows, etc., as directed under F. Pour over a quarter of a pound of black oxide of manganese in a dish placed high half a pint of muriatic acid (spirit of salt), and leave for six hours. It bleaches, and is apt to make white-limed walls sweat—useful for cabs."

Now if we take a room, say 13 feet by 13 feet and 13 feet high, or of a capacity of about 39 cubic metres (and that is not a very large room), and if we calculate what proportion by weight the chlorine liberated by the quarter of a pound of oxide of manganese will amount to, we get about 3 parts of chlorine in 1,000 parts of air. In point of fact, however, the proportion of chlorine in the atmosphere of such a room would never reach anything like 3 per 1,000, inasmuch as walls are not impervious, and during the six hours the air would have changed, and 3 parts per 10,000 would probably be nearer the true

* Read before the Public Medicine Section at the Annual Meeting of the British Medical Association in London, August 1873.