AN ADDRESS ON RECENT ADVANCES IN ANATOMY AND PHYSIOLOGY.

*Delivered in the Subsection of Anatomy and Physiology at the
Annual Meeting of the British Association, 1873.*

By WILLIAM RUTHERFORD, M.D.,

Professor of Physiology in King's College; President of the Subsection; etc.

GENTLEMEN,—In addressing you upon the subjects of Anatomy and Physiology, I would invite your attention to some of the features which characterise these departments of Biology at this present time, and to some recent advances in Physiology, the consideration of which you will find to be possessed of deep interest and importance.

Anatomy, dealing as it does merely with the structure of living things, is a far simpler subject than Physiology, whose province it is to ascertain and to explain their actions. It was not a difficult thing to handle such instruments as a knife and forceps, and with their aid to ascertain the coarser structure of the body. Accordingly, the naked eye anatomy of man has been fully investigated; and although the same cannot be said of that of many of the lower animals, it is nevertheless, as far as this kind of inquiry is concerned, a mere question of time as regards its completion. But minute or microscopic anatomy is in a different position. Requiring, as it does, the microscope for its pursuit, it could not make satisfactory progress until this instrument had been brought to some degree of perfection. Doubtless, much advantage is still to be derived from improvements in the construction of this instrument; but probably most of the future advances in our knowledge of the structure of the tissues and organs of the body may be expected to result from the application of new methods of preparing the tissues for examination with such microscopes as we now have at our disposal. This expectation naturally arises from what has been accomplished in this direction during the last fifteen years. For example, much valuable information has been gained regarding the structure of such soft tissues as the brain and spinal cord by hardening them with such an agent as chromic acid, in order that these tissues may be cut into thin slices for microscopical study. How greatly has the employment of such pigments as carmine and the aniline dyes facilitated the microscopical recognition of certain elements of the tissues. What a deal we have learned regarding the structure of the capillaries, and the origin of lymphatics, by the effect which nitrate of silver has of rendering distinctly visible the outlines of endothelial cells. What signal service chloride of gold has rendered in tracing the distribution of nerves by the property which it possesses of staining nerve-fibrils, and thereby greatly facilitating their recognition amidst the textures. Moreover, of what value osmic acid has been in enabling us to study the structure of the retina. In the hands of Lockhart Clarke, Beale, Recklinghausen, Cohnheim, Schultze, and others, these agents have furnished us with information of infinite value; and those who would advance microscopical anatomy may do so most rapidly by working in the direction indicated by those investigators. In human microscopical anatomy, indeed, there only remain for investigation things which are profoundly difficult, such as, for example, the structure of the brain, the peripheral terminations of nerves, the development of nerve-tissue, and other subjects equally recondite. But in the field of comparative anatomy there is far greater scope for the histological investigator. He has only to avail himself of those reagents and methods which have recently proved so useful in the microscopical anatomy of the vertebrates; he has only to apply these more fully than has yet been done to the invertebrates, and he will scarcely fail to make discoveries. For the lover of microscopical research there is, moreover, a wide field of inquiry in the study of comparative embryology; that is to say, in the study of the development of the lower animals. Since it has become clear that a knowledge of the precise relations of living things one to another can only be arrived at by watching the changes through which they pass in the course of their development, research has been vigorously turned in this direction; and, although an immense mass of facts has long since been accumulated regarding this question, Parker's brilliant researches on the development of the skull give an indication of the great things we may yet anticipate from this kind of research. Speaking of micro-

scopical study before this audience, I cannot but remember that in this country more than in any other we have a number of learned gentlemen who, as amateurs, eagerly pursue investigations in this department. I confess that I am always sorry to witness the enthusiastic perseverance with which they apply themselves to the prolonged study of markings upon diatoms, seeing that they might direct their efforts to subjects which would repay them for their labours far more gratefully. I would venture to suggest to such workers that it is now more than ever necessary to abandon all aims at haphazard discoveries, and to approach microscopy by the only legitimate method, of undergoing a thorough preliminary training in the various methods of microscopical investigation by competent teachers, of whom there are now many throughout the country.

With regard to Physiology, the present standpoint is not so high as in the case of Anatomy. Physiology, resting as it does upon a tripod consisting of anatomy, physics or mechanics, and chemistry, is many sided. The most minute anatomy, the most recondite physics, and the most complex chemistry, have all to be taken into account in the study of the physiology of living things; so that it is not surprising that it should, in its development, lag behind the comparatively elementary subject—anatomy. Until not very long ago, anatomy and physiology were in most of our medical schools taught by the same professor, who, although professing to teach both subjects, was generally more an anatomist than a physiologist. This arrangement gave to physiology a bias which was eminently anatomical, and this bias continued in many quarters, notwithstanding the separation of the physiological from the anatomical tuition. I am aware that there are still some distinguished anatomists who intermingle physiological with anatomical teaching. I am not questioning the usefulness of the practice when carried to a moderate extent. I wish merely to point out what appears to me to have been a result of the practice, and I believe that the result was to give to physiology an anatomical tendency. It was natural for the anatomist who dealt with visible structure to constantly refer to this in explaining physiological action or function. The physiologist with the anatomical tendency always tried to explain a difference in the action or function of a part by a difference in its structure; and when his microscope failed to show any structural difference between the cells which form saliva and those which produce pancreatic fluid, between the egg of a rabbit and that of a dog, he, baffled on the side of anatomy, was too ready to adopt the conclusion that, inasmuch as the microscope reveals no difference in the structure, there is really no difference between them, and that the only way in which the difference in action can be explained is by having recourse to the old hypothesis that the metamorphoses of matter, and the actions of force, are in the living world regulated by a metaphysical entity termed a vital principle, and that dissimilar actions by similarly constructed parts are only to be explained by referring them to the guiding operations of this principle. [After alluding further to the hypothesis of the vital principle and its supposed actions, and after stating that he did not follow the teaching of those who still adhere to this doctrine, the lecturer said that, viewed from the physical side, there appears to be no reason for supposing that two particles of protoplasm, which possess a similar microscopic structure, must act in the same way; for the physicist knows that molecular structure and action are beyond the ken of the microscopist, and that within apparently homogeneous jelly-like particles of protoplasm there may be differences of molecular constitution and arrangement which determine widely different properties.]

A great change is now taking place in physiological tuition in this country;—a superabundance of physiological anatomy, and an almost entire absence of experiment, are no longer the characteristic features of our tuition. The study of physics, too much neglected, is happily now being more and more regarded as important in the preliminary training of the physiologist, as the study of anatomy and of chemistry; and I trust that the day is not far distant when, in our medical schools, the thorough education of our students in mathematics and physics will be insisted upon as absolutely essential elements in their preliminary education. Until this is done, physiology will not advance in this country so rapidly as we could wish. I would not in this place have alluded to a question concerning medical education, but for the fact that the progress of physiology will always greatly depend upon the education of medical men, for only those who are conversant with physics and chemistry, and who, in addition, are acquainted with the phenomena of disease—that is to say, with abnormal physiological conditions—can handle physiology in all its branches. Physiology owes not a little to the study of pathology—that is, of abnormal physiological states. The study of a diseased condition has, on several occasions, given a clue to the discovery of the function of an organ. Nothing was known regarding the function of the spleen until the pathologist observed that an increase in the number of white

corpuscles in the blood is commonly associated with an enlargement of this organ. Hence arose the now accepted doctrine that the spleen is concerned in the growth of blood-corpuscles. The key to our knowledge of the functions of certain parts of the brain has also been supplied by a study of the diseased condition of that organ. The very singular fact that the right side of the body is governed by the left, and not by the right side of the brain, was ascertained by observing that palsy of the right side of the body is associated with certain diseased conditions of the left side of the brain. That the corpus striatum is concerned in motion, while the optic thalamus is concerned in sensation; that intellectual operations are manifested specially through the cerebral hemispheres, are conclusions which were indicated by the study of diseased conditions. Moreover, by the pursuit of the same line of inquiry the key has been given to the discovery of many other facts regarding the brain-functions. Some years ago, M. Broca made the remarkable observation that, when a certain portion in the front part of the left side of the brain becomes disorganised by disease, the person loses the power of expressing his thoughts by words, either spoken or written. He can comprehend what is said to him, his organs of articulate speech are not paralysed, and he retains his power of writing, for he can copy words when told so to do; but when he is asked to give expression to his thoughts by speaking or by writing, or even to tell his name, he is helpless. With a palsy of a portion of his brain, he has lost his power of finding words—he has lost his memory for words; and yet, although he loses his power of finding words, his intelligent perception of what passes around him and of what is said to him is not lost. It is true that this condition of aphasia, as it is termed, has been found to exist when various parts of the brain have been diseased; for example, it has been found to coexist with a diseased state of the posterior instead of the anterior part of the cerebrum. This fact renders it very difficult at present to assign a precise locality to the faculty of speech. It is not, however, my intention to discuss this question, because my object is merely to show how the study of disease has given a clue to the physiologist. Broca's observation led to the thought that, after all, the dreams of the phrenologists would be realised *in so far as* they supposed that the various mental operations are made manifest through certain definite territories of the brain.

It has until lately been supposed that the convolutions of the cerebrum are entirely concerned in purely intellectual operations, but this idea is now at an end. It is now evident, from recent researches, that in the cerebral convolutions—that is, in the part of the brain which was believed to minister to intellectual manifestations—there are nerve-centres for the production of voluntary muscular movements in various parts of the body. Dr. Hughlings Jackson, from the observations of various diseased conditions in which peculiar movements occur in distinct groups of muscles, came to the conclusion that in the cerebral convolutions are localised centres for the production of various muscular movements. It has always been taught that the convolutions of the brain, unlike nerves in general, cannot be stimulated by means of electricity. This, although true as regards the brains of pigeons, fowls, and perhaps other birds, has been shown by Fritsch and Hitzig to be untrue as regards mammals. These observers removed the upper portion of the skull in the dog, and stimulated small portions of the exposed surface of the cerebrum by means of weak galvanic currents; and they found that, when they stimulated certain definite portions of the surface of the convolutions in the anterior part of the cerebrum, movements were produced in certain definite groups of muscles on the opposite side of the body. By this new method of exploring the functions of the convolutions of the brain, these investigators showed that, in certain cerebral convolutions, there are centres for the nerves presiding over the muscles of the neck, the extensor and adductor muscles of the forearm, for the flexor and rotator muscles of the arm, the muscles of the foot, and those of the face. They, moreover, removed the portion of the convolution on the left side of the cerebrum, which they had ascertained to be the centre for the movements of the right forelimb, and they found that, after the injury thus inflicted, the animal had only an imperfect control over the movements of the part of the limb in question. Within the last few months these observations have been greatly extended by the elaborate experiments of my former pupil and now able colleague in King's College, Professor Ferrier.

Adopting the method of Fritsch and Hitzig—but instead of using galvanic he has employed Faradic electricity, with which, strange to say, the investigators just mentioned obtained no very definite results—he has explored the brain in the fish, frog, dog, cat, rabbit, and guinea-pig, and lately in the monkey. The results of this investigation are of great importance. He has explored the convolutions of the cerebrum far more fully than the German experimenters, and has investigated the cerebellum, corpora quadrigemina, and several other portions of the brain not touched upon by them. There is perhaps no part of the

brain whose function has been more obscure than that of the cerebellum. Dr. Ferrier has discovered that this ganglion is a great centre for the movements of the muscles of the eyeballs. He has also very carefully mapped out in the dog, cat, etc., the various centres in the convolutions of the cerebrum, which are concerned in the production of movements in the muscles of the eyelids, face, mouth, tongue, ear, neck, fore and hind feet, and tail. He confirms the doctrine that the corpus striatum is concerned in motion, while the optic thalamus is probably concerned in sensation, as are also the hippocampus major and its neighbouring convolutions. He has also found that in the case of the higher brain of the monkey there is what is not found in the dog or cat—to wit, a portion in the front part of the brain, whose stimulation produces no muscular movement. What may be the function of this part, whether or not it ministers to certain intellectual operations, remains to be seen. These researches of Jackson, Fritsch and Hitzig, and Ferrier, mark the commencement of a new era in our knowledge of brain-function. Of all the studies in comparative physiology there will be none more interesting, and few so important, as those in which the various centres will be mapped out in the brains throughout the vertebrate series. A new, *but this time a true*, system of phrenology will be founded upon them; by this, however, *I do not mean* that it will be possible to tell a man's faculties from the configuration of his skull, but merely this, that the various mental faculties will be assigned to definite territories of the brain, as Gall and Spurzheim long ago maintained, although their geography of the brain was absurdly erroneous, and their notions regarding the indications afforded by the configuration of the skull ridiculous.

I have alluded to the subject, not only because it affords an illustration of the service which a study of diseased conditions has rendered to physiology, but also because these investigations constitute the most important work which has been accomplished in physiology for a very considerable time past. We may, I think, term this the Renaissance period of English physiology. It seems strange that the country of Harvey, John Hunter, Charles Bell, Marshall Hall, and John Reid, should not always have been in the front rank as regards physiology. The neglect of physics must be admitted as a cause of this; it is also to be attributed to the, until a few years ago, almost entire absence of experimental teaching; but it would be unjust not to attribute it in great measure to the limited appliances possessed by our physiologists. It is to be remembered that physiology could not be successfully cultivated without proper laboratories, with a supply of expensive apparatus. Without endowments from public or private resources, how can such institutions be properly fitted up and maintained by men who can, for the most part, only turn to physiological research in moments snatched from the busy toil of a profession so laborious as that of medicine? In defiance of these difficulties, we are now striving to hold our place in the physiological world. A new system of physiological tuition is rapidly extending over the country. In the London schools, in Edinburgh, Cambridge, Manchester, and elsewhere, earnest efforts are being made to give a thoroughly practical aspect to the tuition of our science, and notwithstanding the imperfect results which must necessarily ensue in the absence of suitable endowment, we can nevertheless point to the fact that the effect of these efforts has been to awaken a love for physiological research in the mind of many a student, and the results of this awakening are already apparent in the archives of Royal Societies, in the *Journal of Anatomy and Physiology*, and elsewhere. But physiological research is most expensive and laborious, and it is, moreover, unremunerative. The labours of the physiologist are entirely philanthropic; all his researches do nothing but contribute to the increase of human happiness by the prevention of disease, and the amelioration of suffering; and I would venture to suggest to those who are possessed of wealth and of a desire to apply it for the benefit of society, that, in view of the wholly unselfish and philanthropic character of physiological labours, they could not do better than follow the admirable example set by Miss Brackenbury in endowing a physiological laboratory in Manchester. In London and elsewhere the laboratories exist, but they need endowment: indeed, there is no physiological laboratory in this country which is adequately endowed, consequently research is greatly crippled. The endowment of a dozen such laboratories throughout the country would immensely aid in the development of physiological research amongst us.

We anticipate great benefit to the community not only from an advance of physiology, but from a diffusion of a knowledge of its leading facts amongst the people. This is now being carried out in our schools on a scale which is annually increasing. Thanks to the efforts of Huxley, the principles of physiology are now presented in a singularly palatable form to the minds of the young. The instruction communicated does not consist of technical terms and numbers, but in the elucidation of the principal events which happen within our bodies, together

with an explanation of the treatment which they must receive in order to be maintained in health. Considering how much may be accomplished by these bodies of ours if they be properly attended to and rightly used, it seems to be a most desirable thing that the possessor of the body should know something about its mechanism; not only because such knowledge affords him much material for suggestive thought—not only because it is excellent mental training to endeavour to understand the why and the wherefore of the bodily actions, but also because he may greatly profit from a knowledge of the conditions of health. A thorough adoption of hygienic measures—in other words, of measures which are necessary to preserve individuals in the highest state of health—cannot be hoped for until a knowledge of fundamental physiological principles finds its way into every family. This country has taken the lead in the attempt to diffuse a sound knowledge of physiological facts and principles among the people, and we may fairly anticipate that this will contribute not a little to enable her to maintain her high rank amongst nations; for every step which is calculated to improve the physiological state of the individual must inevitably contribute to make the nation successful in the general struggle for existence.

A CASE OF MELANOTIC TUMOUR DEVELOPED NEAR A CONGENITAL MOLE.*

By FRANCIS MASON, F.R.C.S.,
Senior Assistant-Surgeon to St. Thomas's Hospital.

THE patient, from whom this specimen was removed, was a man, aged 64, who was the subject from birth of a black mole, about the size of a pea, situated on the right cheek, an inch from the angle of the mouth. He remained quite well until a year before his admission into St. Thomas's Hospital, in October last, when he noticed what he described as a "pimple," situated on, or at least very close to, the mole already spoken of. "The pimple" steadily increased in size, and made rapid progress during the last two months. He had never suffered any pain in the tumour, and, excepting for its unsightly appearance and the local inconvenience, he would not have sought advice. For the following notes of the case, I am indebted to the dresser, Mr. E. Scott.

On admission, a fungating mass, about the size of a small orange, was observed on the right cheek (see fig. 1). The skin at the base of the

Fig. 1.

tumour was thin, and had a purplish appearance. The growth was soft to the touch, and was freely movable, apparently not involving the mucous membrane of the mouth, nor occupying any portion of that cavity. The open surface was scabbed over, and, on removing the scabs, a dark livid mass was exposed, from which a thin, sanious, inoffensive fluid exuded. On raising the tumour, the congenital mole was seen on its lower surface, apparently isolated from the rest of the growth. There had never been any severe hæmorrhage from the part. Several glands in the submaxillary region were enlarged, and evidently were deeply adherent. There was no family history of tumour, nor of cancer in any form, and there were no other similar growths visible about the patient.

On October 23rd, I removed the tumour by an incision around the base, dissecting it carefully away; and, after leaving a perfectly healthy raw surface, the edges of the wound were brought together with hare-lip pins. An attempt was made to take away the enlarged glands; but, in enucleating them with the fingers, the capsules burst, allowing a dark brown semi-solid fluid to escape, and, as they were firmly attached deeply, their complete extirpation was scarcely possible, having regard to the important neighbouring structures which would thereby be necessarily involved. As it was, the facial artery was opened and had to be ligatured.

The subsequent history of the case is told in a few words. There was some oozing in the neck soon after the operation, but pressure alone sufficed to arrest it. The hare-lip pins were removed from the face on the third day, and, on the fifth day, the wound had healed by first intention (see fig. 2, from a photograph taken ten days after the operation).

Fig. 2.

The wound in the neck, too, which had been dressed with a weak solution of permanganate of potash, was almost closed, there being little, if any, discharge from it. He left the hospital in about three weeks, quite well.

I may add that I have recently received a letter from the patient's wife, who gives a completely satisfactory account of her husband. He continues his work as a labourer, and there is no evidence of return of the tumour either in the cheek, or, more strangely, in the neck. A section of the tumour presents, to the naked eye, numerous characteristic points. It has generally a nodulated form, and is studded with black spots, the colour varying from light brown to a decided black. Large dark, as well as perfectly white, patches are also observed, with various intermediate shades of pigmental infiltration. The growth is firm in some places, and readily breaks down under pressure in others.

A microscopic examination ($\times 250$) shows the tumour to be made up of a number of delicate cells, chiefly of a spindle shape, containing a varying amount of dark granules, some cells being quite colourless, others having a jet black hue. Here and there, scattered throughout the tumour, but more especially observed at the circumference, there are masses of round cells, containing more or less of the same brownish material. On examination under a higher power ($\times 400$), some cells, containing one or two distinct dark-coloured nuclei, are seen. (Fig. 3.)

Fig. 3.

The chief points of interest in this case seem to be the following. 1. The fact, already noticed by pathologists, that melanotic tumours have a remarkable tendency to be developed either in, or in the immediate neighbourhood of, a congenital mole. I have reason to think that in this case, for it was carefully examined, the tumour was connected with the mole. 2. The curious circumstance, that the congenital mole should remain quiescent, and give no inconvenience whatever for over three-score years; and, lastly, the remarkably healthy condition of the patient, both locally and constitutionally at the present time; for, as some of the tumour was undoubtedly left in the neck, it is surprising that there should be, after ten months, no sign of the recurrence of the growth in this region. How long this satisfactory state of the part is to continue, of course remains to be proved. But the case is one which I venture to think is not altogether devoid of interest, nor wholly unworthy of being brought under the notice of the members of the British Medical Association.

* Presented to the Surgical Section at the Annual Meeting of the British Medical Association in London, August 1873.