absorbed by the blood, it does not actually coagulate the albuminous matters; but it probably renders them less fluid, and reduces their permeating property. Alcohol interferes with the disintegration of blood-corpuscles; and in cases where this is going on very rapidly, and where fluid is passing through the walls of the vessels in considerable quantity, in consequence of the walls themselves being stretched and too readily permeable to fluids, alcohol is likely to be of service; but where these changes are occurring very rapidly, and the patient's strength fast ebbing, it may save life. We may, therefore, explain the beneficial action of alcohol, without assuming that it is a food, or contributes directly to the process of nutrition." And again, he says: "This view of the action of alcohol accords with many broad facts familiar to all. It accounts for the shrivelling of the hepatic cells, the shrinking of the secreting structure, and the increased hardness and condensation of the entire liver, which results from the continual bathing of the gland-structure by blood loaded with alcohol. It accords with the gradual shrinking and condensation of tissues which occur in persons who have long been accustomed to excess. The tendency to increased formation of adipose-tissue, which occurs in persons who live generously, and seems to be augmented by alcohol, may be explained upon the same view; and the stunting in growth which follows its exhibition to young animals is readily accounted for."

These extracts will show that Dr. Beale has, to a very considerable extent, anticipated me by ten years in the theory of the action of alcohol which I advanced in the JOURNAL a short time ago. For a fuller exposition of his views, I must refer my readers to the little pamphlet* itself; since no number of detached extracts can do complete justice to an author's opinions.

I am happy to be able to add that, in a very kind private communication, Dr. Beale disclaims the idea of accusing me of intentionally appropriating his ideas without due acknowledgement; and that he cordially congratulates me on coming to the same general conclusion from a different point of view. The hypothesis was originally suggested to me by Mr. Herbert Spencer's remarks upon narcotics, reference to which I have already given. I must leave others to judge whether what I have said upon the subject has given to the hypothesis additional significance.

PULMONARY APOPLEXIES (HÆMORRHAGIC INFARCTIONS) IN CASES OF CEREBRAL APOPLEXY.

By J. HUGHLINGS JACKSON, M.D., F.R.C.P.,

Physician to the London Hospital.

IN his highly interesting Harveian Oration (BRITISH MEDICAL JOURNAL, July 5th, p. 5), Professor Rolleston refers to Brown-Séquard's important experiments, which show that injuries of certain encephalic centres are followed by pulmonary hæmorrhages. I wish to point out that some of the "experiments of disease" on the higher nervous centres of man produce the same result. I have alluded to this briefly in my article on Apoplexy in Reynolds's *System*, 2nd edition, vol. ii.

I shall in this article use the term pulmonary "apoplexy". It is true that to use this term for local internal hæmorrhages is to write inaccurately; but I never, or scarcely ever, hear used any other term than pulmonary apoplexy to describe what is better called pulmonary hæmorrhagic infarction. In the expression "cerebral apoplexy", the term apoplexy is used strictly; but it is well in this paper to qualify it by the word cerebral. In the two expressions, pulmonary apoplexy and cerebral apoplexy, the word apoplexy has quite different meanings.

After cerebral apoplexy—the effects of any comatous lesions being included in this term—numerous conditions of lung are found. As regards differences in mere quantity of blood in these organs, there are two causative factors—1, the robustness of the patient when taken ill; 2, the rapidity of his death. As a rule, the more robust the patient, and the more rapidly he dies, the more blood in the lungs; but in one case of death in an hour from large cerebral hæmorrhage, I found that the patient's lungs weighed together only twenty-two ounces. They were very like—quite like, I think—the lungs of patients who have died in cholera collapse. This case puzzles me very much, as I have found the lungs much congested in patients who have only lived apoplectic half an hour or less (as after meningeal hæmorrhage). But, as a rule, the lungs are the more congested the more rapid the death, provided death

does not occur *very* rapidly, does not occur within, let us say—and I only state my impression—two hours.

I have above spoken as if there were only degrees of congestion of the lungs in cerebral apoplexy; but in a considerable number of cases we find other appearances *post mortem* which certainly cannot be passed by as "congestion". It is of these that I wish to speak. I omit further reference to well known conditions of posterior congestion, and need not detain the reader by speaking of œdema and acute emphysema.

We see in some cases, on slicing the lung, not only a general reddening of the dependent parts, but *patches* also. These patches are not confined to lobes, although they affect the lower lobes most. They vary in tint from a black to a granite colour; and in some lungs we find both colours and intermediate degrees of colouration. The sections are margined, slightly raised; the masses can sometimes be felt from the surface, but very indistinctly, as nodules. The black nodules vary in area of section from a half-crown to a split pea. The granite-coloured ones vary similarly. These appearances—the differences of stages being well marked—were very striking in the case of a largely built patient aged 50, whom I saw a few months ago with Dr. Stephen Mackenzie. In this case, a few patches were broken down into a pulp.

I think it is plain that the above description of the dark patches is that of "apoplexies"—hæmorrhagic infarctions of the lung. The large black patches are indeed, I consider, unmistakably "apoplectic". The patches (nodules) in heart-disease are, however, for obvious reasons, harder. In a case of coma from injury to the head in a very robust man, I found lungs which were very like, if not quite like, the "damson-lungs" of heart-disease; and I found the same in a case of apoplexy from opium-poisoning. The suicide was a healthy man eighteen years of age.

As to the pathology of the granite-coloured patches, there will be a difference of opinion. As we find them along with black patches in some cases, and as there are all degrees of intermediateness, I suppose them to be decolorised pulmonary "apoplexies". This is what Brown-Séquard has said. But I often hear them called pneumonic. There is, I suppose, the hypostatic pneumonia of Piorry. But, whatever the pathology of the patches may be, I think that the term pneumonia is an unfortunate one. There is not clinical pneumonia. Pneumonia should be kept for an acute affection—of one lung in most cases—which has a fairly definite course, beginning by a severe rigor, and attaining its climax in about a week. When the patches I have mentioned are so numerous as to be confluent, we may find the dependent parts of the lungs bulky and solid. But even then both lungs are nearly always affected—one more than the other, it is true, in many cases; and they are not affected in one or more lobes only, but in all lobes and in the dependent parts of these. When, however, there are a few scattered granite-coloured masses, the condition will occasionally be called lobular pneumonia. The geographical part of the expression is, I think, correct; it is a *lobular* affection, but it has not the clinical associations of lobular (catarrhal) pneumonia. It is lobular; but there is no evidence that it is inflammatory—that is, inflammatory *primarily*. Sometimes the pleura is inflamed over a patch seated near the surface; but the same can occur in the pulmonary "apoplexies" of heart-disease, which are certainly not inflammatory primarily. It occurs also in the hæmorrhagic infarctions of pyæmia. The latter, by the way, present stages of decolorisation analogous to those of which I speak in severe cases of coma. It is true, they go on to suppuration; and so, although very rarely, may the apoplexies of heart-disease.

The general opinion of medical men in this country, so far as my knowledge goes, is, that pulmonary "apoplexy" in heart-disease is owing to impeded *exit* of blood. It is not, however, easy to understand how this process can lead to *localised* hæmorrhages, even if it be easy to understand that it can lead to general congestion, splenisation, etc. The more likely explanation seems to be, that the "apoplexies" result from pulmonary embolism. Virchow's experiments, indeed, demonstrate that embolism does produce pulmonary apoplexies. The embolic material in cases of heart-disease is believed to be blood coagulated during partial stasis on the chordæ tendineæ; and I suppose the same explanation holds in the pulmonary apoplexies of cerebral apoplexy. Every one knows that splenic and renal blocks are *red* when recent; and the accepted explanation of "red softening" of the brain is, that it is dependent on *obstruction* of small arteries. It is true that there are difficulties in understanding how increased quantity of blood, and even extravasation, result from plugging of arteries; but of the facts of the matter there is no doubt. It is warrantable, then, to infer that the pulmonary apoplexies of heart-disease or of cerebral apoplexy arise in a like manner from blocking of arteries (pulmonary). It is not necessary to deny that impeded exit has something to do with the production of pulmonary "apoplexies".

* *On Deficiency of Vital Power in Disease, and on Support; with Observations upon the Action of Alcohol in Serious Cases of Acute Disease.* By Lionel S. Beale, M.B., F.R.S. London: T. Richards, 37, Great Queen Street. 1863.

If the above reasoning be correct, the pulmonary apoplexies from lesions of the encephalon arise, not directly from the nervous lesion, but indirectly and by a quasi-mechanical process; there being primarily produced imperfection in the respiratory and circulatory movements. The great emphysema which in some cases we find anteriorly is complementary to the congestion, oedema, and solidifications found posteriorly. Thus it admits of a mechanical explanation too. The distinction is important. If my explanation be correct, the hæmorrhages, congestion, emphysema, etc., do not result by vaso-motor or other nervous actions directed straightway upon the lungs themselves.

Several of the pulmonary states mentioned occur, of course, under other conditions than those of cerebral apoplexy. But I suggest cases of cerebral apoplexy for study, because there is often a *sudden* and *local* lesion. In traumatic cases in healthy and occasionally robust persons, the process is near akin to the experiments which the physiologist makes. I have not been able to conclude as to the influence of exact position of encephalic lesion; I do not think it makes a great difference. Indeed, I have seen, as before remarked, lungs like those of heart-disease in a young suicide who died apoplectic from opium. In cases of brain-disease, the two elements of size of lesion and suddenness of lesion are more important than position of lesion. Without hastily concluding, I would suggest that the matter of greatest importance is to note the *mode of death* in numerous cases of apoplexy, and especially the alterations of respiration and circulation in cases of lesions, large and small, differently placed. But it will be essential also to note the health of the patient before his illness, the suddenness of the attack, and the rapidity of death. These, however, are very difficult matters, and I shall consider the methods and objects of such observations another time.

THERAPEUTIC MEMORANDA.

CROTON-CHLORAL HYDRAT.

THE profession and the public are chiefly indebted to Dr. Oscar Liebreich for the introduction of chloral hydrate; and this obligation is further increased by the addition of croton-chloral hydrate, which will doubtless prove an equally valuable therapeutic agent. It is of the greatest service in cases of nerve-pain. Every sufferer from neuralgia is anxious to obtain speedy relief from pain; this may be obtained by taking croton-chloral hydrate, and then the antecedent causes of the neuralgia may afterwards be inquired into and treated accordingly. The following cases are interesting, as showing the immediate relief from pain that this drug affords.

A. suffered from facial neuralgia of a most severe character; it affected her hearing and eyesight. She could not rest or take food. She took one grain of croton-chloral hydrate every hour. In three hours, she was considerably better. After taking three more doses, she was entirely free from pain.

B. suffered much from facial neuralgia dependent on decayed teeth, and had not been able to take food or sleep for three days. She was ordered croton-chloral hydrate in grain-doses every hour, and obtained great relief after two doses. Six doses removed the pain completely. She slept that night.

C. This patient suffered from concussion of the spine caused by a railway accident some years ago. She has had every variety of treatment for the pain she suffers in the spine and the nerves proceeding therefrom. She took potassium bromide gr. 20, and croton-chloral hydrate gr. 1, three times a day, with marked relief and no bad symptoms.

E. This is a young dyspeptic and neuralgic patient, and suffers greatly from dysmenorrhœa. She took two-grain doses when the paroxysms of pain came on, with marked relief.

F. has been under treatment for various neuralgiæ for some years. She has had, at one time or another, almost every external and internal therapeutic agent in the *Pharmacopœia*—strychnia, iron, quinine, ammonium, chloride, aconite, belladonna, iodine, bromine, blisters, hypodermic injections, galvanism, together with baths and other hygienic appliances, including change of air. In this case, two grain-doses of croton-chloral hydrate every hour afforded more speedy relief from pain than any of the above remedies. After taking eight grains, she was almost free from pain.

In thirteen patients who have taken croton-chloral hydrate, not a single bad symptom has been observed. In grain-doses, it relieves pain quickly; causes natural sleep; no subsequent headache or furred tongue. In several cases, it acted as a gentle laxative.

BENSON BAKER.

REPORTS

MEDICAL AND SURGICAL PRACTICE IN THE HOSPITALS OF GREAT BRITAIN.

ST. MARY'S HOSPITAL.

INCISED WOUND OF WRIST, INVOLVING THE CARPAL JOINT: RECOVERY, WITH SLIGHT IMPAIRMENT OF MOVEMENT.

(Under the care of Mr. EDMUND OWEN.)

A LITTLE girl, just under two years of age, was admitted under the care of Mr. Edmund Owen on October 21st, 1872, with a severe wound about the left wrist, which had been caused, about an hour previously, by the blade of a chaff-cutting machine. The patient had suffered from considerable hæmorrhage on her way to the hospital. On examination, it was ascertained that the flexor carpi ulnaris, the ulnar nerve and vessels, and some of the more internal of the flexor tendons of the fingers, as well as the ulna itself, had been divided by an incision which

Diagram of Wrist, with dotted course showing line of division of Ulna (u) and Radius (r).

passed obliquely downwards and outwards from an inch above the head of the ulna into the radio-carpal articulation. The blade had also cut through the lower end of the radius in such a manner that the lesser sigmoid cavity was separated from the shaft, the inferior radio-ulnar articulation being uninjured, whilst the interior of the wrist-joint was freely exposed.

Mr. Owen, having consulted with Mr. Knott, the resident registrar, came to the conclusion that the proper course of treatment would consist in endeavouring to save the hand, notwithstanding the severity of the injury, remarking that the processes of repair were extremely active in healthy young patients, and that even if Nature should fail to effect a cure in this case, the surgeon could subsequently come to her assistance. The ulnar artery was then tied at the wound with a carbolised catgut ligature, of which the ends were cut short; the edges of the wound were brought together with sutures of the same material; and, dressings soaked in weak carbolic acid lotion having been applied, the arm was lightly bandaged on a splint. When the dressings were removed upon the third day, the appearance presented by the wound was very satisfactory. Pulsation, too, could be distinguished along the ulnar side of the little finger, showing that at this early period collateral circulation had been established between the radial and ulnar arteries. All went on well for three weeks, when erysipelas and abscess attacked the back of the hand. Incisions and poultices were employed, and steel-wine and port prescribed. Under this treatment the child gradually improved, and was discharged about three weeks subsequent to the formation of the abscess, with the wound entirely healed, and the limb still supported upon a splint.

When the child was last seen, at the beginning of May, there remained some stiffness at the wrist-joint, and an extensive cicatrix indicating the position of the incision—the child using the hand, the mother said, almost as freely as before the accident.

WOLVERHAMPTON AND STAFFORDSHIRE GENERAL HOSPITAL.

A THIRD YEAR'S CASES OF LITHOTOMY.

(Under the care of Mr. VINCENT JACKSON and Mr. NEWNHAM.)

OF the examples of stone in the bladder admitted into the hospital during the year 1872, the following eight cases were successfully (excepting one) operated upon by lithotomy. The operation is always