even none. But this I can affirm : I cannot call to mind a case in which a stricture that gave trouble was not relieved by rest, recumbency, and the other means which have been mentioned. If this be so, should it not form an essential and primary part of the treatment of stricture? Should not the most be got out of these measures before we resort to instruments?

If it be urged that this would be waste of time, which may be saved by employing instruments at once, it may be replied that chiefly out of such mischievous practice have arisen such convictions as those to which I referred at the commencement of this paper. But then it will be argued that after all you come at last to instruments to accomplish the cure. Not always—very often not. I do not pretend that, under the sole influence of these measures, which are simply hygienic, all traces of stricture usually disappear ; but the patient is very often, and sometimes speedily too, restored to such comfort, he is so relieved from every trouble, the stream of urine flows so naturally, that he is well satisfied with the result, and declines any further interference.

But by these means you have not thoroughly cured his stricture : you have given him only temporary relief, and his troubles may at any time return. That will in many cases depend on himself, and the care he takes of his health. No doubt in most cases of severe stricture these simple means are only palliative. Frequently, indeed, they will succeed so well that the patient himself is no longer conscious of anything amiss, yet the passage of an instrument will still detect an abnormal condition. But supposing you fall back on instruments, what then? How much more than this can you do? Dare you, in the majority of cases of severe stricture, promise by their aid a complete and permanent cure? Can you do more than restore a patient to a condition of comparative comfort, yet still with liability, under unfavourable circumstances, to relapse?

The sum of my argument, then, amounts to this : while I am not suggesting that instruments are never to be employed in the treatment of stricture, I venture to affirm that they have been, and still are, grossly abused ; that an untold amount of mischief is perpetrated by their abuse ; that the routine employment of instruments in the treatment of stricture is in the highest degree unscientific and improper ; that, in a word, in the treatment of stricture, instruments of any kind should never be employed in any way except as a last resource.

REMARKS

ON

CASE OF INTESTINAL OBSTRUCTION FOLLOWING OVARIOTOMY.

By THOMAS KEITH, F.R.C.S., Edinburgh.

THE subject of the following history was my 160th case of ovariectomy. She was a young Irish lady in her 20th year, of a highly nervous temperament, thin, and fragile looking. For some months, she had suffered severely, and had been much pulled down by the disease. She had been tapped in February by Dr. Wallace of Parsonstown ; and, when he sent her to me in the end of March last, she was already as large as before.

Ovariectomy was performed on the 1st of April, 1873. Dr. Zell of New York and other friends were present. There was nothing unusual in the operation. Omental and general parietal adhesions were separated, several catgut ligatures applied, and, as a good deal of blood had been lost during the breaking up of the semi-solid portion of the tumour, I did not follow my usual practice, in young subjects, of letting ten or twelve ounces of blood escape from the pedicle before securing it. In this case, both ovaries were diseased. The pedicle of the right was of fair length ; that of the left was extremely short. Both were divided by the cautery, and, after waiting till all oozing had ceased, the wound was closed by silk sutures in the usual way. Ether was given during the hour and a half that the operation lasted. The weight of the broken down cyst-walls and contents was upwards of 40 lbs.

She did not do well from the beginning. From the afternoon of the operation, there was an amount of restlessness, such as I have never seen after ovariectomy. The pulse rose the first evening to 140, and continued so for the first five days ; the temperature ranged from 101 degs. to 102 degs. ; the limbs were rarely for a moment still ; the nights were restless as the days, and were spent tumbling and tossing about. There were thirst and dry tongue ; yet elimination went on well. There was an unusual amount of clear urine, and free perspiration from time to time. Flatus first passed twenty-six hours after the operation, and

continued to pass fairly well, yet the abdomen became distended, and, on the fourth day, there was much vomiting. On the fifth day, fluid was for the first time detected in Douglas's space, on examination by the rectum—for no vaginal examination was, in this case, at any time practicable. The fluid collected rapidly, and, on the seventh day, there was distinct fluctuation above the pubes ; on the ninth, this reached to within an inch of the umbilicus. The wound, though much stretched, was healed, and the general condition improved. In the evening, as tension, which had hitherto been slight, had suddenly become great, and the patient was very uneasy, a small trocar was pushed by the rectum into Douglas's space, and, after six ounces of thick dark blood had flowed, the cannula became choked by clot, and was removed. This so far relieved the tension, that the patient had some hours' sleep. Next morning—the tenth day—a long syphon trocar was used, and thirty-six ounces of thick syrupy blood, not the least foetid, were removed by the rectum as before ; then about three ounces of clot were sucked out by a powerful syringe, care being taken to admit no air. This gave much relief ; the pulse fell to 100 ; the abdomen became soft ; a large deep hollow being left above the pubes, very distinctly arched over by a roof of adherent intestine.

It was now that the anxieties of the case began ; for it soon became evident that, as the roof of the hæmatocele fell inwards, a trap had been formed somewhere from folds of intestine adhering at angles ; and, though the solid contents of the bowel moved downwards, or were moved by medicines, yet no air was allowed to pass. The fault seemed to be not far from the lower end of the colon.

The eleventh day after the operation was spent quietly, though scarcely any flatus escaped ; and, by the twelfth, this had ceased altogether to pass outwards. Towards evening, the abdomen became distended, and windy pains were troublesome. On the thirteenth day, the bowels were freely moved by oil and an enema, without any relief to the flatulence. Douglas's space was again punctured, and ten ounces of bloody purulent matter, with broken down pieces of clot, were removed. The abdomen was raised by coils of distended intestine. On the fourteenth, the pulse was again up to 120 ; the distension was general and excessive ; the rectum was dilated and empty ; all attempts to pass a tube into the colon failed, as they had done before, and changes of position gave no relief. Enemata brought away some solid fæces, but never any gas followed. By evening pain was severe, and opiates were, as on previous days, necessary to moderate the increasing peristalsis.

On the morning of the fifteenth day, a little fluid was again felt in Douglas's space, and five ounces of thick pus were removed by the syringe. The cavity was then washed out with warm water, as the matter was a little foetid. This was the last of the hæmatocele. The distress and pain were this day great, and she was restless exceedingly. The distension was now enormous ; the girth at the umbilicus was 52 inches, greater than before the removal of the tumour ; the ensiform cartilage was turned upwards, and the stretched, though firm, cicatrix was almost at a right angle to the pubes. Prominent amongst the ever-restless coils of intestine, the colon—now as big as one's arm—from the thinness of the parietes, was easily traced upwards and across the epigastrium, but lost below and to the left of the umbilicus. By evening, the patient was much exhausted ; the pulse was 140 ; and it was clear that the heart could not much longer bear the tension upon it. Late at night, my brother and Dr. Sidey gave me their help, and we agreed to make one more attempt to get a tube into the colon, and, should that fail, to open the abdomen up. At her request—for she disliked the smell of ether—chloroform was given. The hand was put into the rectum, and the fingers were with ease passed above the brim of the pelvis ; but, farther than two inches beyond, no tube could be made to go. This took some time, and on turning the patient round to proceed with the proposed operation, respiration was found to have ceased, and it seemed for a time that we had to do with a death from chloroform. She slowly recovered, however ; but the feebleness was so great, that such a severe proceeding as opening up the abdomen was not now to be thought of. The finest needle of an aspirator was then pushed into the transverse colon ; it was put in several times, but seemed to be at once choked. I then punctured with a small trocar, and a very large quantity of air escaped through the cannula. Some nepenthe was injected under the skin, and she was left at midnight in a deep sleep ; the pulse was then 140, though steadier since the relief given to the heart by the puncture.

Next day—the sixteenth after operation—there was less pain, and the abdomen was not so full. Some flatus passed in the forenoon, and several times during the day ; pulse, 136 ; temperature, 101.8.

On the nineteenth day, much flatus passed, and the bowels were moved six times. She was nervous and feeble all day, with a tendency to faintness, and free stimulation was necessary. Towards evening, flatus again ceased to pass downwards, and the spasms became severe ;

pulse, 140; temperature, 102 degs. On the eighteenth, the abdomen was again distended, and, though discharges of fluid feces went on during the day, not a particle of air was suffered to escape. The next five days were spent without change; the tension again became frightful, and the sufferings during these weary days were very great. Sometimes the peristalsis was allowed to have its full swing; generally it was kept within bounds by opium or belladonna, and a light poultice over the abdomen gave much relief. On the night of the twenty-fourth, chloroform was carefully given, and the colon was again punctured. The cannula was passed well downwards, and then held against the abdominal wall to prevent its slipping, as it had done before. It was retained for some time; much gas escaped, and, as the peristalsis came and went, there were great discharges. The tension was immensely relieved; opium was again injected, and, as before, she slept all through the night.

During the next day, and the day after, she was feeble, scarcely able to speak. The pulse had fallen to 120; the bowels had been moved freely by oil, and, soon after, flatus passed in great quantity. Preceding this improvement, there was a profuse diuresis, sixty ounces of urine having been passed within twelve hours. On the twenty-ninth day, the belly was quite flat; there was no return of distension, and, though reduced nearly to a skeleton, she rapidly gained flesh. Seven weeks after the operation, she returned to Ireland, and remains well.

There can be little doubt that the puncture of the intestine saved this patient. It did so by gaining time; for, as already stated, it was pretty certain that a portion of the adherent bowel, which formed the roof of the hæmatocele, had become trapped as it fell downwards after evacuation of the fluid. It was hoped, if the patient could only live long enough, that more or less absorption of the connecting lymph would soon take place, and lead to some change of position in the arrangement of the part at fault, that would allow of air to pass along with the solid contents. Fortunately, this happened. The profuse diuresis, which occurred just before the relief, I have seen take place once before in a case of obstruction, which terminated favourably.

I do not guess at the source of hæmorrhage in this case. Some might, perhaps, blame the cautery, or the slipping of some of the catgut ligatures; but there have been similar cases of hæmatocele after ovariectomy in which the clamp was used, and in which no adhesions existed.

I have never seen any harm in puncturing hæmatocèles or collections of fluid in the pelvis by the rectum. If, on the occurrence of tension, I wish to empty Douglas's space, I prefer to do so by the rectum rather than by the vagina. With a proper instrument, this is a safe procedure. What I have long used for this purpose is the ordinary trocar and cannula, of the size and length of a No. 6 catheter. The shoulder of the cannula is moved an inch forwards from the end, so as to allow of a long India-rubber tube being slipped on as the trocar is withdrawn. With the slightest attention, no air can thus enter. The operation can be repeated as often as necessary, and, if the fluid is too thick, it can be sucked out with a syringe. Of course, I would not drain by the rectum; but drainage by the vagina is a very imperfect way of keeping Douglas's space free. With a patient on her back, and a dilated recto-vaginal pouch, a tube passing through the vagina will not keep Douglas's space empty, but will leave ample room for the stagnation of fluids, which can hardly help becoming putrid. It seems to me, therefore, that Dr. Marion Sims's scheme for the drainage of the abdomen by the vagina, in all cases of ovariectomy, will not satisfactorily accomplish his object. Even if it did, it would almost always be unnecessary, and it might, by carrying putridity inwards, not seldom lead to blood-poisoning of the patients.

THERAPEUTIC MEMORANDA.

ONYCHIA MALIGNA.

I WAS much interested in perusing Mr. MacCormac's remarks on this distressing malady, and having met with many cases of this and the allied disease "ingrowing nail," I was delighted to find what I considered, and experience has proved, a perfect cure. Rest and attention to the state of general health having preceded, the fungous growth is then burnt with strong nitric acid, washed with water, and poulticed. The relief is certain, and the repetition of the application seldom necessary. If there should be any trouble with the nail, the tender flesh may be protected by the insertion of a thin piece of compressed sponge, kept in its place by strips of plaster applied longitudinally to avoid compression.

BENJAMIN BLOWER, Liverpool.

CLINICAL LECTURE ON CONVERGENT SQUINT.

By R. LIEBREICH,

Ophthalmic Surgeon and Lecturer, St. Thomas's Hospital.

I SHOW you to-day three patients with strabismus convergens, who represent three forms of strabismus, different with regard to treatment. In the first case, the squint shows itself only while the look is attentively fixed upon an object, whether near or remote. If, however, the patient look absently before him, contenting himself with an undefined image of the object, his eyes remain perfectly straight. This state is characteristic of the first stage of the ordinary strabismus convergens based on hypermetropia. In order to see distinctly with eyes the refraction of which is not sufficiently strong, it is necessary to make a great effort of accommodation. There is a connection between the accommodation of each eye—that is, of the power of the eye to adapt itself to different distances—and between the muscles which make the two eyes converge to a single point. On account of this connection, a hypermetropic eye can only make the effort of accommodation necessary to correct its fault of refraction, by making at the same time a very great effort of the two recti interni—that is, by producing too high a degree of convergency, which, in fact, constitutes the strabismus. So the periodical squint becomes a means by which hypermetropical children succeed in seeing distinctly. It is true that, by doing so, they give up the binocular vision; but they see better with one eye sharply fixed than with both eyes incompletely adapted to the distance of the object. If, in these cases, the optical fault of the eyes be corrected by adding glasses which make the vision equal to that of a normal eye, the squint will disappear; for then every distance requires only the normal quantity of effort of accommodation, as well as the normal degree of convergency. The first indication of treatment is to find out the degree of hypermetropia and of the astigmatismus which may be combined with it, and to advise the use of spectacles which correct these optical faults. This is not always easy. The principal difficulties we meet with are the following.

1. The youth of the patient. There are cases where the affection begins as early as the second year, and even earlier, at a time when the use of spectacles is out of the question. In such cases, one must be content to draw the attention of the parents to the matter, in order as far as possible to prevent the child from fixing his eyes on small near objects; and, later, reading must not be attempted without the use of the right spectacles.

2. A difficulty based on spasm of accommodation. In order to ascertain the refraction of an eye—that is, to find out which is the farthest point to which it can accommodate itself—it is necessary to suspend the accommodation. It is, however, not always possible to obtain this in eyes which, on account of hypermetropia, are in a constant state of tension of the accommodation. On being examined, they either appear not at all hypermetropical, or at least not so much as they really are. Their hypermetropia is then called latent. In order to make it manifest, different ways may be adopted: either the accommodation may be entirely suspended by the instilling of atropine, and then the refraction is easily ascertained; or, choosing a slower way, we may diminish by degrees the effort of accommodation by the use of convex spectacles. These spectacles are to be chosen first approximately; then gradually stronger ones must be used, until they are in accordance with the degree of hypermetropia.

3. A third difficulty will often be found in the strong aversion of parents to seeing their children wear spectacles. They often urge to have an immediate operation performed, in order that the deformity of squinting may be removed at once. Some surgeons give way to such wishes; but let me earnestly caution you against this. The result of an operation in such a case is, that the convergent squint is, in the most favourable case, removed; that is, during the fixation, under the conditions which produced the convergence before, the axes of the eye are now more or less well adapted. On the contrary, while the look is not definitely fixed on an object—that is, under the conditions which, before the operation, made the axes of the eyes stand more or less parallel