increase of its influence for good. This had been effected by the diffusion of a purer ethical code impressed on the minds and spirits of men, rather than by written laws enforced by tribunals. Out of this there had sprung a charitable feeling, which made effort to aid the least fortunate among them, and to succour those who had been left unprovided for by them by means of a benevolent fund.

Secondly: Much had been done already towards consolidating the vital interests of the profession. Thus it was greatly due to the voice of the Association that a Medical Council had been established to register all legally qualified practitioners, and to form a standard of therapeutics embodied in a national Pharmacopœia, that had supplanted the too numerous productions of colleges, and to aim at uniformity of education and tests for the entire kingdom. Although these last desiderata had not yet been accomplished, there was ground for hoping that when the mind of the profession was sufficiently educated to be of one accord on the essentials, these great points of internal reform would certainly be achieved.

Thirdly: Its external utility was amply displayed in the successful operations of its "State Medicine Committee", to which it was in a great measure due that they had any Public Health Act, supervising the eradication of the causes of diseases in cities and towns, by sanitary measures; and, although much still remained to be done, yet it might be confidently predicted that the voice of their profession, heard in the repeated warnings of this Committee and its district Branches, would make itself not merely listened to, but attended to. It had already triumphed frequently in the cause of the public, over the narrow obstructiveness of corporations and vestries. Again, by vigilance of the Parliamentary Bills Committee, erroneous measures had been corrected before obtaining the force of enactments. Undoubtedly, through its influence, much had been gained by those of them who undertook the care of the sick poor, and by those whose office it was to treat sick and wounded sailors and soldiers in the service of the State. For the last divisions, he might state that they were fully sensible of, and gratefully acknowledged the friendly hand that had been extended on more than one occasion, assisting them to obtain a due consideration for their labour and calling.

Fourthly: The Association might glory in its having conferred lasting benefits on the sick poor by persistent efforts to obtain mitigations of the too levelling and centralising policy of the Poor Law Board, in whose economical reforms acts were done oppressively to the sick and those disabled by age. The details of the Poor Law were much more mercifully administered now than they were a few years since, thanks to their profession in a great degree.

Lastly: In a scientific direction, the Association had done much by its *Transactions*, and of later years by its *JOURNAL*, which, under its present editorship, was second to none, and had become the fair exponent of the views and studies of its individual members. And, as much by its annual meetings, for the general discussion of the objects and management of the Society, as well as for the advancement of science, it brought home to them concisely and lucidly a summary, and the results of all the recent investigations devotedly pursued by the most illustrious men of the day. And now, at length, a point had been reached when a more direct participation would be taken in such investigations. To this end, a recommendation would be made at the Annual Meeting, 1874, to grant £200 in aid of researches in anatomy and physiology, the action of remedies, clinical medicine, and surgery. All these things were demonstrations of innate strength, marked by slow and steady growth, and as successive triumphs worthy of being recorded, and likely to arouse enthusiasm among them, akin to that the soldier felt when he enlisted in a regiment with untarnished laurels, to follow its glorious colours into fields where fresh honours were to be participated in. Each of them should strive to augment and maintain what had been already secured by the elder Branches of their Association, and to prove themselves not unworthy of being enrolled in the same list.

Proceeding next to local arrangements, he spoke of the results obtained by the Contagious Diseases Act, which the Admiralty had been prominent in instituting and maintaining in spite of an unprecedented clamour in the name of a pseudo-morality and of public decency, which, as he contended, had been greatly advanced by its operation; and he called on his audience to proclaim their knowledge, so that the arm of the Government might be strengthened, and the nation enlightened, so as to give a wider adoption of a great sanitary principle. In another direction, he believed that equal good would be derived in the direct action of Government as a corrective to the slowness of municipal authorities in stamping out the causes of some zymotic diseases. Among these, he instanced the fact that the causes of enteric or sewer fever were never absent in certain districts of these towns, especially in the hastily constructed houses allied to the barracks at Eastney, which were

of themselves, perhaps, the finest in England, from among the men quartered, in which there were more cases of the disease sent to Haslar Hospital than from all the more numerous bodies of men under control of the Admiralty quartered here. He attributed this to the very imperfect sewerage and water-supply of the neighbourhood of the barracks in which the families of the soldiers resided for the convenience of being near the barracks. In addition to his personal observations, he quoted those of the medical officers immediately concerned, in proof of the facts. While admitting that much was being done in securing better conditions in the newer portions, he contended that much remained to be done to correct past neglect, and expressed a hope that the Government would use its weight in stimulating municipal authorities to correct all proved evils arising from defective sewerage. The necessity for activity in this direction had become the more needful, inasmuch as otherwise the health of the Portsmouth district was second to that of none in the kingdom; and, with this corrected, it would stand superior to all. To this end, he advocated the strengthening of the hands of the health-officer, who had done very much, and was alive to all these imperfections. Whatever influence established or increased the public safety, the united strength of civilian and Governmental practitioners in working to the advantage of the community at large, would be seen and felt, and the hand should not be stayed so long as there was any remediable cause of which disease was an inevitable product. Reminding his audience that to aim at the prevention of disease was more conservative of life than the most successful treatment of it could ever become, he remarked that this fact had been strongly impressed on military medicine from the days of the great conqueror Cyrus, whose father Cambyses, according to Xenophon, taught him that the prevention of diseases in the camp was one of the most honourable functions of a general, while the office of treating diseases stood relatively to it, as the office of repairing the cast-off garments of the army. In this respect the medical duties had undergone great changes in modern times, although the executive had not yet fully surrendered to the medical element its authority in such matters, in apprehension that military strategy might be embarrassed or retarded by any secondary interposition. Whatever might have been the recent experience and practice of other conflicting nations, they might trust that at least the salutary lesson had been learned from the management, and in the results of their own little war in a deadly climate, that medical foresight and executive reliance on it would deprive war, under its worst aspects, of many of its horrors, while they tended to ensure rapid and complete success now, when brevity of a campaign had become an absolute necessity. Sooner or later the lessons now accumulating in social and military science must bring about a just appreciation of the fact that all questions concerning public health, that is of communities, were most safely dealt with by the science of medicine and its educated professors. To this desirable end, nothing could be more conducive than a deep sense of its responsibilities actuating the profession, and that sense was becoming more fostered now than at any previous period, mainly through the co-operation of all classes in one fraternity—the British Medical Association. They were striving in the right path by uniting to form a new Branch of it, to be guided by its principles in pursuit of its prominent objects, while hoping for the same results within their own local sphere. Let them, then, work together, zealously holding this sole end steadily in view.

THE GAINS DERIVED FROM THE USE OF THE LARYNGOSCOPE.

By PUGIN THORNTON, Esq.,

Assistant-Surgeon to the Hospital for Diseases of the Throat, and Surgeon to St. Marylebone General Dispensary.

THE profession will, I am sure, look forward with interest to the evidence resulting from Dr. George Johnson's experiments and pathological observations. In my desire not to encroach on the space of the *JOURNAL*, I fear that the observations I ventured to put forward in explanation of his interesting case were not quite clear. I will, therefore, briefly explain.

1. By idiopathic disease of the muscle, I mean disease of the muscular structure, independent of the nerve-supply. The laryngeal muscles are so superficially situated, that they are very readily affected by inflammation of the mucous membrane; and it not unfrequently happens that, when the inflammation subsides, the structure of the muscles remains impaired. This subject has been elucidated by Dr. Morell Mackenzie in his work on *Hoarseness and Loss of Voice*; and, under paralysis of the adductors of the vocal cords (page 5), he specially calls attention to the views of Professor Gerhardt, who enumerates two

kinds of rheumatic paralysis of the muscles of the vocal cords. In addition to this cause of paralysis, syphilitic ulceration and gummatous tumours occasionally leave atrophy of the muscles.

2. As regards the state of the crico-arytenoid articulation, it would be interesting to know if this joint was opened at the *post mortem* examination of Dr. Johnson's case.

3. It would be desirable to ascertain, if possible, what was the real condition of the muscles in this case. Were they submitted to a microscopical examination? I ask this question, because the information contained in the *Transactions* of the Pathological Society from Dr. Curnow's examination was briefly summed up as follows. "He reports that 'the laryngeal muscles on the left side are decidedly atrophied; those on the right side are somewhat larger; but he is inclined to consider them atrophied also'."

ABSTRACT OF LECTURES ON THE SURGICAL TREATMENT OF ANEURISM.

Delivered at the Royal College of Surgeons of England.

By T. HOLMES, F.R.C.S., Professor of Pathology and Surgery.

LECTURE IV.

IN commencing the subject of popliteal aneurism, the lecturer stated that he could only hope to touch somewhat lightly on the numerous points which that subject presents for discussion. He hoped to establish, at any rate, the encouraging fact, that the treatment of this disease in our present hospitals in this kingdom is exceedingly successful, both by the Hunterian method and otherwise.

In the first place, the different forms of popliteal aneurism were spoken of as follows. 1. Traumatic and other rapidly growing aneurisms, in which the sac is very thin or imperfect. These are not readily made amenable to any of the milder methods of treatment. 2. Aneurisms which grow into and irritate the femur, and in which the irritation is liable to be reflected on to the sac of the aneurism, provoking its speedy rupture. Many of these aneurisms have been found to be only curable by amputation, of which some instances were alluded to. 3. The shape of the tumour influences its curability, those which are of the bud-like form being more amenable to milder measures than those which approach the cylindrical. 4. The progress of the symptoms has much weight in the choice of treatment, those in which the disease is acute being of evil omen for the success of compression. 5. The presence of concomitant disease, though it does not always forbid the success of ligature or amputation, certainly indisposes the surgeon to any cutting operation. The first step, then, in deciding on the method of treatment to be adopted in any case of popliteal aneurism is to ascertain, as well as can be done, these facts, and to give them their due weight on one side or the other. It is thus chiefly that our choice is regulated as to the selection of the ligature or of compression. The next question is, supposing compression to be first adopted, have we such definite evidence of its great success, as compared with the Hunterian ligature, as to make it our duty to persevere with it at all hazards, even when it seems at first to be doing no good? There seems at present no proof of this. The results of treatment all over the country by compression have not as yet been shown to be much, if any better, than those by ligature. Whether they could be made so is a different question. The effect of previous compression on the success of subsequent ligature does not seem to be favourable, but the reverse; and the success of the Hunterian operation in late years appears to be much greater than is shown by the statistics of Norris and others.

After these preliminary observations and a slight sketch of Hunter's operations (the preparations from two of which are in the College Museum, both successful), the lecturer went on to discuss the results of the ligature of the artery on the method of Hunter, *i.e.*, when undertaken as the sole means of treatment with no previous compression. The number of such cases contained in the "Table of Hospital Cases", on which these remarks are founded, was seventy-seven in popliteal and ten in femoral aneurisms. Two of the latter died; the rest succeeded. Of the seventy-seven popliteal cases, there were eleven deaths, and, in three other cases, the ligature failed, but the patients recovered—in one case by the old operation, and in the other two by amputation. In several of the fatal cases, death occurred really from other diseases, and in two from direct injury to the vein in the operation. The rarity of cases of secondary hæmorrhage, gangrene, or supuration of the sac was very remarkable in contrast with what Norris's table shows. In fact, the whole result shows that, in the hands of British hospital surgeons, the Hunterian operation has proved about twice as successful of late years as was the case formerly, accepting

Norris's table as a perfectly authentic and sufficient record; and the great success which has attended the same operation in the hands of individual operators, such as Syme and Mott, also tends to show that its dangers have been overestimated. In all probability, this improvement is due to improved surgical practice in this particular. The old operators (Hunter, Roux, etc.) were by no means careful to avoid injuring the vein; they exposed the artery much more freely than is now done; they could not dissect so quietly and systematically without chloroform as can now be done; and, as a general statement, it may be said that surgeons now are more practised anatomists and dissectors than they used to be. It results, then, as a practical inference, that, if for any reason compression seem contraindicated, or if its trial be unsatisfactory, the surgeon may resort to the ligature with confidence, as being a most successful operation.

Next, the general results of compression were stated from the same table. Compression had been tried in sixty-six cases with success, and in fifty-eight without success; in other words, it had failed almost as often as it had succeeded. It must be remembered, however, that these results might probably be much improved. It seems doubtful (to say the least) whether the compression-treatment is not often carried out rather negligently in our hospitals, being entrusted to students, nurses, and others, who have not the necessary knowledge or the necessary interest in the case. The success of compression depends on minute and long-continued attention to a number of details by a perfectly competent surgeon, and, in the absence of such minute attention, we do not obtain the results which the method is capable of yielding. Yet, even if compression were to continue to fail as often as it succeeds, this ought not to deter us from adopting the treatment, unless there be some plain proof that, when it fails, it decidedly spoils the prospect of successful treatment otherwise, of which as yet we have no proof. The average results of the whole number of cases treated by pressure are no worse than those of ligature, though they are not decidedly better, considering that compression is not generally used (as the ligature is) in the worst class of cases. At the same time, it is not to be denied that compression has dangers of its own. Excluding cases such as are recorded in old books, where the injudicious use of pressure has led to deep sloughing and arterial hæmorrhage—cases which should never occur now—there are a certain number where it has been followed by gangrene, and some in which the tissues of the vessels compressed seem to have been injured. This latter point was illustrated by Verdier's case, in which the condition of the external iliac artery was shown after four years' use of a compressing apparatus, and by Mr. O. Pemberton's case of the formation of an arterio-venous communication after compression of the femoral; and, though neither case was absolutely decisive, yet, taken in conjunction with other recorded instances, they showed, at any rate, the possibility of damage being done by the compression to the vessels and to the cellular tissue which surrounds them.

The further consideration of the subject was deferred to the next lecture.

CLINICAL MEMORANDA.

CASE OF MAMMARY ABSCESS IN AN INFANT THREE WEEKS OLD, CAUSED BY RETENTION OF MILK.

ON June 17th last, Mrs. R. was safely delivered of a healthy female child, well developed. The child continued well till July 3rd, when I was called to see her. I found her suffering from colic, which was relieved by medicine. Next day, I also observed a swelling of the right mamma, which was hard and inflamed, and tending to point on the inner side. The following day, I opened it, when about an ounce of thick greenish pus escaped. On examination, I found a milky secretion escaping from the left breast, some of which I collected on a slide, and, for the purpose of comparison, put a drop of the mother's milk also on another slide. On examining them by the microscope, I found them to be alike in composition, only that the mother's milk contained about five times as many corpuscles as the child's, while the child's milk contained colostric corpuscles, which I did not see in the mother's. I do not find a case of this kind mentioned by any of the authorities within my reach, and several medical friends, to whom I have spoken of it, have never heard of a similar case. There can be little doubt that the abscess was caused by retention of milk in the right breast, which is a common cause of it in adults, as the left breast was sucked out regularly by my direction, and did not inflame. There is still an oozing of pus from the right breast from a second abscess which formed on the outer aspect of the mamma, and discharged its pus through the opening on the left side.

WILLIAM STEPHEN, M.D., M.A., Fettercairn.