mother; during our journey, he entered most fully into the history of his mother's illness, detailing all her symptoms, but did not let drop one word about habits of drinking. On my arrival at the patient's house, the nature of the case was but too evident; the lady was dying of drink; but I got no hint of the real cause, until I made inquiry from my colleague after the examination was ended. In another case, in which the anxiety of the parents about their daughter was excessive, and was also justified by her death, I paid my first visit without receiving information on this subject from anyone with whom I came into communication; it was only when I pointedly suggested a possible cause for uncontrollable vomiting at my second visit that a free admission was made by my colleague, who had been restrained from offering any communication to me by the absolute horror on the part of the parents of any reference being made to the fact of their daughter having aided herself to endure a husband's neglect by the sustaining power of stimulants.

ALCOHOLISM IN PRIVATE PRACTICE.

By H. SUTHERLAND, M.D.,

Lecturer on Insanity at the Westminster Hospital.

PREVENTION is better than cure in all diseases, but especially in those produced by an excessive indulgence in alcohol. Intemperance, just short of being absolutely drunk, is now far more common amongst the upper classes than it was ten years ago; and it is more than ever needful for those who see the ill effects of an inordinate indulgence in stimulants to remind their patients that the good things of this world are intended to be used, and not abused.

The whole question, however, is beset with many difficulties. Some men can take a very large quantity of alcohol, and are all the better for it; others, with weaker heads, try to follow their example, and suffer in various ways in consequence. The methods of treatment or of prevention are in many instances faulty and prejudiced. On the one hand, we read of sensational temperance meetings, where the use of alcohol is altogether condemned. On the other side, we have the fashionable doctor prescribing champagne at lunch for his lady-patients. Many people err in the use of alcohol from mere ignorance, and are really grateful for a few practical hints as to what they may and what they may not take during the day.

The following observations, deduced from numerous inquiries, may possibly be of service to those who have doubts on this subject. They are intended to point out some of the premonitory symptoms of alcoholism, selected from typical cases, and the most rational means of avoiding excessive indulgence in stimulants.

CASE I. *Extreme Nervousness from Alcohol.*—Mr. W. H., a stock-broker, complained of a painful state of nervousness. He was afraid to walk out of sight of his house in the morning. He drove to the City in a brougham with his wife, and did not dare to go alone. In the morning he was always in a state of collapse. He transacted his business, but was so nervous at the office, that he was obliged to lean on the arm of one of the clerks in going down stairs. This condition necessitated two or three glasses of neat brandy, but he never felt that he was "himself" until after lunch. At that meal, he drank a pint bottle of stout and two glasses of sherry. He had a glass of sherry in the afternoon; three or four more at dinner; a pint of claret and about two glasses of whiskey and water at night. He was ordered to leave off the brandy and whiskey altogether, to drink water at dinner, and to take wine only at the end of the meal. After a week of this treatment, he was much improved.

CASE II. *Transitory Mania from Alcohol.*—An officer in the army, aged 24, came up to town for a spree. He very seldom took much. One night, he was dining at one of the military clubs, and drank a bottle of champagne, besides sherry, hock, and claret. After dinner, he had two glasses of neat brandy. He soon became much excited, and executed a "view-holloa" in the club dining-room, much to the astonishment of other members. He left the club with his friend, and took two more glasses of brandy at a public-house. After this, he became more excited, struck a stranger on the back with his stick in Regent Street, tore down a large advertisement, and was ultimately taken to his lodgings with much difficulty. Here he attempted suicide with a razor, and, in flourishing it about, the blade fell back on his knuckles and cut him severely. Next, he boxed at his looking-glass, and struck it with such force as to cut a piece of skin as large as a

sixpence off the back of his hand; not breaking the glass, however. After that, he tried to get out of the window; but assistance was procured, and he was strapped down in bed, and prevented from doing any further mischief. Next day, he left town apparently well, no treatment by drugs having been resorted to.

In both these cases, the patients were advised to reduce their quantity of alcohol, and were told that no healthy man ought to exceed four glasses of wine a day. They both had sufficient common sense to know where to stop; and there is no reason why they should not recover from their incipient symptoms and become temperate and healthy men.

The prognosis in the following cases, however, was far less favourable, as the habit was of long standing, and was indulged in to the extent of getting drunk frequently and without regard to consequences.

CASE III. *Dipsomania.*—On March 21st, a clergyman called on me, and gave me the following particulars about the state of mind of his daughter. Miss P. E., aged 28, had run away for the third time from her home, and was found in London, alone, in a lodging-house. There was no hereditary history of insanity, phthisis, or nervous disorders. The father was very hysterical, and burst into tears several times while he related his story; but this was sufficiently accounted for by the conduct of his daughter. She had always been healthy as a child until she was sixteen years old, when she had scarlatina. After this illness, she went out of the house before she had properly recovered, and caught a chill. This was succeeded by an attack of meningitis. For five years after that, she suffered from great debility, accompanied with various mental symptoms, such as slight loss of memory, weakness of judgment, irritability, wilfulness, cruelty to her sister, lying and spreading false reports about her relatives. These symptoms gradually became worse; and, as there was considerable loss of tone and want of sleep, the medical treatment included stimulants and also opiates. The patient then got into the habit of treating herself, and would take large quantities of chlorodyne, sal volatile, paregoric, or laudanum, during the day, if she fancied she required it. For the last four years, she had been addicted to taking opiates to a great extent. She employed a number of agents in the parish to procure opium for her secretly, and rewarded them by giving them her old or new clothes recklessly. Latterly, a taste for alcoholic stimulants was also developed, which she gratified by frequent visits to public-houses, and was found on several occasions staggering about the streets of her father's parish in a drunken condition. During this period, she was placed under medical treatment, and the opium and stimulants were cut off entirely. This sudden change in her mode of life had immediately disastrous effects. She became very restless, weeping all day, begging for wine and laudanum, and at last was so debilitated that she was confined to bed for four weeks. The treatment succeeded temporarily, however, and she abstained entirely for three months. At the end of that time, she took to her old habits again, and became more intemperate than before. She took a violent dislike to her relatives; stole a twenty-pound note from her father; pawned a watch, which originally cost £20, for 15s.; and succeeded in inducing her friends to give her dresses costing altogether about £50, saying she was going to India, which she never had any intention of doing. The father procured the necessary order and certificates, and with no little trouble she was traced to a lodging-house near a convent, where she had been residing, and from which she had been expelled for misconduct. After some manoeuvring, I and my friend Mr. Bond succeeded in obtaining an interview. Four days previously, she had been out at night in search of a situation as governess, and had got drunk and fallen down an area. She was picked up by a policeman, taken to the station-house, and locked up in the cells for the night. Her hand was severely injured by the fall, and she had bruised her eye, nose, and chin. Her dress was covered in every part with dried blood; and, beyond a night-dress, she had nothing with her but the clothes she was wearing. After a prolonged interview, in which she told several lies, which she afterwards confessed, it was found that there were no evidences whatever of insanity about her. The father was, therefore, informed that there the doctor's duties ended, as the case was not one which could be legally treated in an asylum. A most distressing scene followed. The father begged his daughter to return home; and this she obstinately refused to do, although there appeared to be no chance of her getting a shelter anywhere else. In the course of our interview, this lady confessed to taking an amount of alcohol equal to about eight glasses of wine a day, and, in addition, six drachms of laudanum every night. It was, therefore, necessary to order four glasses of wine daily, and two drachms of the tincture of opium every night, although there was apparently not the slightest hope that she would limit herself to this quantity if she had an opportunity of further gratifying her extraordinary propensities.

CASE IV. *Dipsomania*.—Mr. A. D., aged 60, a rich merchant from Lancashire, sent for medical advice on the morning of Sunday, March 22nd. He was one of a large family, none of whom had been addicted to habits of intemperance; nor was there any hereditary taint of insanity, phthisis, or nervous disorder. He had always been a strong healthy man. He had been remarkably prosperous in business, and was noted for possessing an unusual amount of shrewdness and common sense. For the last forty years, however, he had lived very freely, and had been frequently intoxicated. Latterly, this habit had become confirmed, and he had drunk to such an extent as to become a perfect nuisance to his sons, who were all sober and steady, and who were carrying on his business for him. About four days ago, he came up to town to visit his branch establishment in the City. He confessed that on one day he had drunk a bottle of champagne in the afternoon at the office, two bottles of sherry at dinner, besides beer and whiskey during the day and also when he returned home at night. On Sunday morning, he awoke in a great state of tremor, having slept very badly, and felt so unwell that he became alarmed, and sent for a doctor.

He was a remarkably fine robust old man; and, although his memory was said to be affected by his long continued potations, it was impossible to discover any proofs of intellectual aberration. On the contrary, he asked some very pointed questions about the proposed treatment of his case. The skin was warm and perspiring. In front, the lungs were fairly healthy; but behind there were loud sonorous *râles* all over the chest, and dulness with tubular breathing at the base of the right lung. It was afterwards ascertained that he had caught a severe chill when coming home in a cab in a semi-intoxicated condition. The heart's action was feeble, but regular; and there were no murmurs. Pulse 104, sharp and rather small. The liver was somewhat enlarged, the hepatic dulness extending to two and a half inches below the ribs. There were also pain and tenderness in this region, both in front and behind. The tongue was very large, soft, flabby, and tremulous, and covered with irregular white patches. On making him raise the hands with the fingers extended, it was observed that they were also very tremulous. The question of treatment resolved itself into present and future. The lung-symptoms, being the most urgent, were treated first; and a night-draught containing hydrate of chloral and bromide of potassium was ordered. The patient was also advised not to take more nor less than eight glasses of wine a day, which quantity was to be gradually reduced. The improvement was very rapid, although it was probably due more to the fact that the old gentleman was kept in bed and away from unlimited alcohol, than to any medical remedies which were prescribed.

The lesson to be learnt from such cases is, that our code of laws is decidedly in a defective condition, when such patients are allowed to be at large, and to injure both themselves and their relatives by their habits of intemperance. In the first case, there was a young lady bringing disgrace and ruin upon her parents and brothers by getting into debt and drinking to excess on every possible occasion. As she was of age, her father could not legally detain her; and, being sane, she was unfit for an asylum. In the second case, an old man was imperilling the prospects of his relations, and putting them into a continual state of alarm, by getting frequently intoxicated in public. He being the head of the family, they were afraid to offend him; and, as he was of sound mind and obstinate, they were compelled to keep him at home.

We have every reason to believe that there are many persons amongst the upper classes who are suffering from minor ailments, and are sowing the seeds of greater ones, by their ignorance of the use and abuse of alcohol. What is required for such persons is an establishment in which they could seek refuge until they had conquered their dangerous propensities, and where they would enjoy greater liberty and privileges than they could in a lunatic asylum. The patients might be of two classes, and in the same or in different institutions. One class should include such cases as those detailed above, who are absolutely ruining the prospects of their families by their bad habits, and who should be confined, whether they like it or not. The other class should consist of those who, feeling the necessity for surveillance, would voluntarily place themselves under the discipline and *régime* of asylum-life.

The treatment of alcoholism in its earliest stages need seldom include the administration of drugs. The patient's cure is in his own hands, and he is usually in good bodily health. Some people ridicule the idea of there being any necessity at all for limiting their daily amount of alcohol: others are really grateful for a few practical hints on the subject. Living, as we now do, in an age of public dinners and general festivity, it is especially necessary to lay down a few simple rules for our guidance. Some of these, which we frequently prescribe for our patients, are here given in detail, for the benefit of those who suspect

they are taking habitually a larger allowance of alcohol than is compatible with health.

Alcoholic stimulants should never be taken in the morning before lunch. If one be obliged to do so habitually, it is certain that alcohol is gradually getting a hold on the individual, which he will not easily throw off. If, in exceptional cases, a person be called upon to undergo unusual fatigue in the morning, and feel the necessity for a stimulant, a glass of beer may be taken. No one in good health should ever take wine or spirits in the morning.

If anyone fear that he is taking too much alcohol, the easiest way to limit the quantity is to drink the same allowance of the same wine every day. This, for various reasons, should be brown sherry, and should be taken as follows: either two glasses after lunch and two after dinner, or one and a half after lunch, the same after dinner, and one with a biscuit in the afternoon. During lunch and dinner, water must be drunk, and wine taken only at the end of the meal.

No healthy man should ever take more than four glasses of wine daily as his usual allowance, and should reduce this to three if possible. Those who are accustomed to wine should not take less than this every day.

Those who wish to become teetotallers should leave off alcohol very gradually, and the period during which this reformation is brought about should extend over not less than two years.

No healthy man should ever touch spirits, whether raw or diluted, except under specially exceptional circumstances, as, for instance, after great exposure to wet or cold. Liqueurs at dinner are also to be avoided.

Roughly speaking, one tumblerful of beer contains about the same amount of alcohol as one wineglassful of wine or half a wineglassful of brandy.

Ladies who are in good health should take one glass of sherry or claret at lunch; one and a half ordinarily at dinner; one glass of champagne and one of claret at dinner-parties; and two glasses of champagne with seltzer water at balls.

Ladies in delicate health, who are going to ride in the park, if they have breakfasted early and do not lunch until late, may take a glass of dry sherry and a biscuit before starting for their ride.

Men, at dinner, may take a glass of sherry with their soup, if they have not had one in the afternoon; no hock; a glass and a half of champagne and one of sherry or claret after dinner. If they will keep to this, they may dine out as often as they like, and it will not hurt them.

At public dinners, the same rules should be observed; but, as there are occasions on which it is almost impossible to avoid drinking much, it is then recommended that they should either have a tumblerful of water by their side, and drink a mouthful of it after every sip of wine, or that they should drink champagne and seltzer water, half and half, all through dinner.

At balls, the safest drink as refreshment is claret-cup. Bad claret is the least offensive bad wine. A biscuit should be taken with it. It is astonishing what reparative powers there are in a biscuit. This hint was obtained from a volunteer who had tried it successfully during the autumn manoeuvres.

At supper, under all circumstances, champagne should be diluted with seltzer water. If the champagne be questionable, claret-cup should be taken, or beer, if it can be got.

Wine should never be taken undiluted, especially on an empty stomach, when one is hot and dusty. This rule particularly applies to travelling.

Brandy and soda-water should be avoided as rank poison. Diluting the brandy does not destroy its pernicious effects; and the soda-water, by its bulk, acts most injuriously by separating the food from the walls of the stomach, thus preventing the gastric juice from having free access to it and assisting in the process of digestion.

If these rules have been neglected, and a severe bilious headache be the result, nothing short of an emetic, which may consist of a table-spoonful of mustard in a tumbler of hot water, will be effectual.

If a person suspect that he has taken bad wine, and fear the result, it is a good plan to take about thirty grains of carbonate of soda, *on the same night*, before going to bed.

Mild bilious attacks may be treated by soda, sal volatile, ether, and peppermint-water, taken before breakfast and during the day between meals.

The best kind of supper to have at home is a biscuit and a glass of water.

Finally, wine is a good friend, but a bad enemy. It is the abuse, and not the use of alcohol, which is injurious; and, in using it, let us not forget the motto of Abernethy, who, when asked what virtue was most conducive to a healthy life, replied in one word—"Moderation".