cinical work, when tried solely on its own ground, that of histology, the data from which distinctions are drawn appear to be too trifling or insecure. Malignant sarcomata, for example, are classed with comparatively innocent growths, and separated from carcinoma, on the ground that, while both are composed of cells, the cells of carcinoma are of an epithelial type, while those of sarcoma spring from connective tissue. I do not say that this is the sole ground of distinction. There is the very subordinate difference, probably dependent on this, of arrangement of those elements. But this, I take it, is the radical distinction; yet of what after all consists the line of demarcation between connective tissue and epithelium? Is it strong enough—that is, physiologically distinct enough—to be the durable basis of such a classification?

But, at all events, if hereafter a more stable basis for the arrangement of tumours shall be devised than any which is founded upon the character of their elements, in the meanwhile I would suggest that, in structures of a doubtful kind, in tumours of no definite description, in growths of a mixed nature, and as an aid to, or in criticism of, conclusions drawn from current views, more attention than heretofore might with advantage be given to a study of the relation of a low grade of development to dissemination and recurrence.

In summary then:—The tissues which are simplest in structure and mode of development are most apt to form the substance of tumours, as, for example, connective tissue. The most elaborate tissue in these respects—striated muscle—is very rarely, if ever, found as a distinct morbid growth, and it is comparatively rare to find even unstriated muscle as a separate mass in the substance of tumours. In the elementary structures of tumours, there is an inverse ratio between the tendency to progressive metamorphosis and the tendency to reproduction. The less the structures of which a tumour is composed tend to change from their primary or embryonic form, the more abundantly will they multiply; so that those tumours whose structures retain most nearly their primary form are the most malignant. And, as the structures of a tumour are capable of transformation, so do they lose their power of repetition; so that those tumours which consist most completely of fully formed tissue are the most innocent.

LABORATORY NOTE: ON THE TENSILE STRENGTH OF THE FRESH ADULT FŒTUS.*

By J. MATTHEWS DUNCAN, M.D.,
President of the Obstetrical Society, Edinburgh.

THERE are almost innumerable laboratory experiments demanded in the present state of obstetrical science; for, while many important points in the mechanism of natural and of unnatural delivery may be admirably illustrated or explained by them, others can in no other way be made out or settled. To give a list of those which at once occur to the mind would be tedious and, to me at least, very tantalising, for I have been able to make very few; and this deficiency arises, in great part, from the want of material. Yet those acquainted with the recent progress of obstetrics can easily recall to mind some of the most valuable of our recent steps in advance, which have been attained through following that experimental method, the dawn of whose usefulness is as yet scarcely recognised by obstetricians.

The experiments which I am now about to describe were made with a view to arrive at correct ideas as to the force available in delivering a child footling; for example, in the operation of podalic extraction now extensively recommended in the conduct of certain cases of contracted pelvis. I knew that even among our best recent authors there has prevailed gross error on this point, the available force being absurdly overrated and the force that could be exerted in the alternative forceps operation almost as absurdly underrated. It would be easy, but it is unnecessary, to cite strong asseverations to this erroneous effect from the most recent authors, widely reputed as our best. Now the point is not of second, but of first-rate importance in the question; indeed, a fundamental matter. It is, therefore, well worthy of careful discussion, and fortunately it is easily settled in a quite conclusive manner.

I took a fresh newly-born child, and, after passing its body through an aperture so cut in hard wood as to represent the brim of a contracted pelvis, or after fixing its head below the biparietal diameter between two parallel bars, I applied above an ankle an apparatus consisting essentially of a hook, whereby to suspend the weights, which were to

be gradually increased till they produced disseverment of the body of the child.

As is well known, the disseverment in actual practice takes place, as it did in all my experiments, in the neck. The disseverment was, therefore, always decapitation.

The force effecting decapitation is in these experiments equal to the combined weight of the child beneath the line of disseverment, of the apparatus attached above the ankle, and of the special weights applied. But great nicety was not attempted, the increments of weight attached being made by several pounds at a time.

It was desirable to limit the time during which in each trial the weight was allowed to operate, and the term of a half-minute was adopted, the weight not being allowed to act longer than that time in one trial. A limit of this kind was necessary to bring the experiments into conditions somewhat analogous to the pains or force of labour and the efforts of the accoucheur in a difficult case in actual practice.

The decapitating force thus discovered is the measure of the tensile strength of the fœtus; and, of course, in podalic extraction, whether by the feet or by the Prague seizure, no greater force effective in delivering the head can possibly be applied, for the materials give way when the weight reaches the decapitating amount.

In actual practice, a greater dragging force may be exerted by the accoucheur and applied to the child. But such force, so far as it exceeds the decapitating amount, does not directly help the delivery of the head. It is either usefully spent in overcoming friction of some part below the line of disseverment, or uselessly or even banefully spent, because misdirected. In actual practice, however, when the head is fixed on the brim, the decapitating force must be very nearly an exact highest limit of the force that can be exerted by the accoucheur.

These remarks are made without regard to the life of the child, and with a view merely to the strength of the materials. The experiments to be narrated will show that the life of the child is compromised before the limit of the tensile strength of the fœtus is reached. This is easily explained by the destruction of the spinal column, which occurs under a force considerably less than the decapitating force; and this circumstance must be kept in mind when podalic extraction is compared with forceps-extraction; for, in making the comparison, it is supposed that the life of the child is not compromised. Neither operation would be selected as the best, if the securing a living child were hopeless.

Experiment 1. A fresh adult female fœtus, weighing 5 lbs. 6 oz., and measuring eighteen inches in length. The spinal column gave way under 90 lbs. Decapitation took place under a weight of 118 lbs. The disseverment occurred between the fifth and sixth cervical vertebra.

Experiment 2. A fresh adult female fœtus, weighing 7 lbs. 7 oz., and measuring twenty inches in length. The spinal column gave way under a weight of 120 lbs. Decapitation took place under 141 lbs. The disseverment occurred between the sixth and seventh cervical vertebra.

Experiment 3. A fresh adult female fœtus, weighing 8 lbs. 15 oz., and measuring twenty-three inches in length. The spinal column gave way under a weight of 122 lbs. Decapitation took place under 136 lbs. The disseverment was between the fourth and fifth cervical vertebra.

Experiment 4. A fresh adult male fœtus, weighing 5 lbs. 12 oz., and measuring twenty-one inches in length. The spinal column gave way under a weight of 91 lbs. Decapitation occurred under a weight of the same amount. The disseverment was between the fourth and fifth cervical vertebra.

Experiment 5. This experiment is not to be included with the former, as it was performed not on a newly-born child, but on one a fortnight old. It was fresh, and weighed 7 lbs. 5 oz., and measured twenty-two inches in length. The spinal column gave way under a weight of 147 lbs. Decapitation took place under a weight of 163 lbs. The disseverment was between the third and fourth cervical vertebra.

The dissevering or decapitating force in these experiments was 118, 141, 136, and 91 lbs. respectively, or, on an average, about 120 lbs. This is probably far from being what most obstetricians would regard as a great force; and it may be said with truth that, if we are to avoid premature decapitation, the force used should not reach this amount; should not, indeed, exceed a hundredweight. Such a force, if used only for a short time, if it cause no great contusion, and especially if it do not break the continuity of the mucous surface, may be safely used in repeated efforts in delivering. On this point, there is much inexactness from our ignorance as to how much and how long the soft parts will endure compression without great resulting evil to the mother.

This podalic dragging force of a hundredweight is considerably greater than any ever known to be exerted in propulsive efforts in natural parturition, and so also is the force of 100 lbs., which is about the limit of what is available without destroying the life of the child. Estimates of the force of labour-pains have, as is well known, been published, which make it much above a hundredweight; but such esti-

* Read before the Obstetric Medicine Section at the Annual Meeting of the British Medical Association in Norwich, August 1874.

mates have been shown to be unreliable in themselves and in the methods by which they have been reached. It is a part of the theory of podalic extraction in contracted pelvis, that the force exerted there by the accoucheur exceeds that of the best labour-pains, and it is interesting to observe that the experiments confirm at least this part of the theory. At the same time, it is necessary to realise that the excess of force in favour of podalic extraction is truly very limited.

I am sure that most accoucheurs, not being familiar with dynamometrical experiments, will regard each his own available muscular power as being far more than he can dare to use in podalic extraction; that is, far more than a hundredweight. No doubt he is right, in so far as that he can easily lift a hundredweight, perhaps even two; but that is a very different and a very easy matter compared with applying a dragging force of a hundredweight in a special direction. Probably no ordinary man can stoop over a bed and pull in a direction across the front of his body, or in what would now be called a coronal direction, with a force of a hundredweight, and this is what many attempt and erroneously fancy they can do. Numerous failures arise from practitioners erroneously estimating the amount of force they are exerting, and still more from erroneously estimating the amount of extractive or dragging force they are exerting on the fetus, since it is common to confuse the amount of force exerted with the amount made efficient as a dragging force; and it is the direct dragging force that may be allowed to rise to about a hundredweight. If a proper judgment be formed of the great difficulty of using a dragging force of a hundredweight, except when the accoucheur is favourably placed, it will be easily understood how important it is to have the patient in a convenient position; for the practitioner must wield his great force skilfully, and gently, and easily, if he is to do his work in the best way.

In the foregoing experiments, it will be observed that the cervical part of the vertebral column invariably gave way, and the dis severed vertebrae became widely separated from one another before the whole neck yielded and decapitation was effected. While the vertebral column yielded when the weight or force was about 105 lbs., a further addition of about 15 lbs., forming a total of 120 lbs., was required to produce decapitation. From this, it is evident that at first almost the entire dragging force is exerted through the vertebral column; that the tensile strength of all the tissues of the neck is not simultaneously and equally tried, and that the vertebral column is weaker than the combined soft parts. The vertebral column yields with a snap or jerk, and immediately a marked elongation of the foetal body occurs. Of both of these events, the accoucheur can scarcely fail to be aware, even if he did not expect them, and this information is important practical direction to him, if he wish, as he almost certainly does, to avoid decapitation. The yielding of the column of bones does not restrict the force he can exert; for he can generally, not always, apply more, to the extent of about 15 lbs., without producing decapitation. But the events referred to form a distinct warning to him as to how near he has reached the amount of force available by dragging the foetal body. He cannot, when they have occurred, pull with greatly increased force, even for half a minute at a time, without incurring the imminent risk of decapitation.

Another point is clearly shown by these experiments; viz., the sufficiency of a single limb for dragging, so far as mere strength of materials is concerned; and this determination has many evident practical bearings. In the trials narrated, we uniformly attached the apparatus and its connected weights to one leg just above the ankle. The parts were protected by a piece of leather wrapped around them from the cutting sort of edge presented by the cord used to tie on the apparatus. One leg always proved stronger than the whole neck. It is interesting to remark, with reference to the leg, that, just as in the neck, the bony column was weaker than the other combined soft parts; for, during the dragging, various cracks or snaps in the limb were heard, and were subsequently found to have been produced by separation of femoral or other epiphyses. One limb always proved more than sufficient for the neck. Whatever advantages may or may not arise from using the Prague seizure, one thing is sure, that, with a view to merely having sufficient strength of materials, it is not required; one leg is, in this point of view, sufficient. It is plain that this is not exactly proved by the experiments, for in them the weight acting on the neck exceeded that acting on the ankle by the greater part of the weight of the body; but the ankle will not have this advantage, except when the body is dragged downwards vertically, or nearly so, which is not done in practice.

I have spoken incidentally of the comparison between the power that may be exerted by dragging the child in podalic extraction without compromising its life, and the power that may be exerted by dragging the child by the forceps. While the limit of the former power is easily

ascertained and has been just stated as about 100 lbs., it is no part of my present object to determine the higher limit of the latter, and it is a far more difficult matter. But there are various ways of showing that the latter is considerably higher than the former, which, as I have already said, is not the universal or even the general belief. I have made many experiments in connection with this subject, but I shall not here enter on them, satisfying myself with giving some easily appreciated evidence.

An accoucheur can decapitate a child by the efforts of podalic extraction; in other words, his available power overcomes the whole tensile strength of the fetus. Few men pulling in an advantageous manner find much difficulty in applying a dragging force of 120 lbs.; but it is quite possible to apply a forceps to a foetal head, at the brim of an actual pelvis or of one artificially constructed for experimental purposes, so as to defy the utmost efforts of the most powerful accoucheur. These iron hands can be made virtually of any strength, and the grasped spheroidal foetal head is firm enough to resist such quasi-cephalotriptic compressing or otherwise destroying force, as they exert in maintaining their hold against a dragging force greater than can be used in podalic extraction, because greater than is sufficient to rend in twain a foetal body.

A CRITICAL EXAMINATION OF DR. MUNRO'S VIEWS ON MALARIAL FEVERS AND CHOLERA.

By W. C. MACLEAN, C.B., M.D.,

Surgeon-General; Professor of Military Medicine in the Army Medical School at Netley.

THE fourteenth volume of the *Army Medical Department Reports*, recently published, contains a paper on "Malarious Fevers and Cholera", by Deputy Surgeon-General Munro, C.B., head of the Medical Branch in the office of the Director-General, on which I feel constrained to offer some observations. It would not, perhaps, be difficult to assume the office of a critic, either anonymously or under a *nom de plume*, and thus escape thrusting myself in *propria persona* before the profession. In the present case, such a course, it strikes me, would be neither fair to Dr. Munro nor otherwise suitable. Dr. Munro does not write anonymously; he puts forward his deliberate opinions in an official publication with all the weight he derives from his official position and his deservedly high reputation in the Medical Department of the Army. Moreover, the issues he raises in his paper are not matters of merely speculative interest; they involve questions of the gravest practical importance in medicine and sanitation. Opinions coming from such a source are entitled to respectful consideration and fair and responsible criticism. To what Dr. Munro has written I have given careful consideration, and I propose, in the following observations, to discuss the issues he has raised, I hope with the courtesy and fairness the author has a right to expect, and with the fulness the importance of the subject demands.

In all controversies the first thing to be done is to understand clearly, and to state exactly, the matter in dispute. In this case there is no difficulty, as Dr. Munro has plainly put the difficult thesis he has set himself to maintain with sufficient distinctness. It is—1. That there is no such poison as malaria; 2. That so-called malarial fevers, cholera, yellow fever, and heat-apoplexy, are merely different manifestations of one and the same disease, the difference being one not of cause, but of degree; 3. That the cause of the above diseases is "certain electrical conditions" (the nature of the said conditions is not explained), which are the outcome of "heat acting on moisture".

It will be seen at once that Dr. Munro admits the action of two agents which have always been thought active in evolving malaria from the soil; but, instead of agreeing with those who believe in the existence of any poison or "telluric influence" to which the convenient name of malaria has been given, he attributes the disease-producing agent to the above named electrical conditions acting with paralysing effect on the sympathetic nervous system.*

* It is not denied that the sympathetic system of nerves is implicated in grave malarial fevers, and in specific yellow fever in particular. Many of the most notable objective symptoms in the latter point to this implication. Bellot the younger even insists that this can be demonstrated pathologically *post mortem*. The other nervous centres are implicated too. A pigmented condition of the brain is one of the most familiar of the morbid appearances in malarial fever. It is for Dr. Munro to connect the morbid appearances with the "electrical conditions" of which he writes so positively.