Another difference is found in the fact that scarlet fever is not often associated with other diseases, while diphtheritic affections are not uncommon in the course of other pyrexial disorders. It is recorded in the Report of the Diphtheria Subcommittee of the Epidemiological Society, that, in the epidemic of 1861, this disease occurred fifty-seven times alone, thirty-four times in association with scarlatina, nine times with small-pox, seven with measles, six with fever, and three times each with ordinary sore-throat, croup, and catarrh. But in the nature and symptoms of the two diseases are points of unlikeness too great to allow us to rank them as varieties of the same species. Thus, the rash of diphtheria is often absent, is very variable as to the time of its appearance; it occurs seldom at the outset of the malady, and I have seen it as late as the third week of its course. The period at which albuminuria sets in is also different, often appearing in severe cases of diphtheria within the first two or three days, whilst in scarlatina it seldom sets in until degeneration of the kidney is commencing. There is, indeed, an entire absence of definiteness about the duration of the symptoms of diphtheria, which of itself marks it off from the regular sequence of events in scarlet fever. I have known it last for six weeks, and even two months, without any evident affection of the cervical glands, but with constant formation of exudation on the fauces or on other mucous surfaces.

The mode in which diphtheria localises itself on the mucous membranes, or on the broken surfaces of the skin, would probably be sufficient to separate it from any other disease. I do not know that scarlet fever ever attacks the larynx; but I have seen the diphtheritic membrane appear in different cases upon wounds of the skin, upon blistered surfaces, and upon nearly all the mucous membranes of the body—the lips, cheeks, nose, ears, pharynx, larynx, trachea, and bronchi, and on the anus and vulva; and different epidemics have differed strangely in this respect; in one, the throat would always be the point selected for attack; and in another, the other mucous membranes. It is somewhat remarkable in the latter case that, although the disease was mild and there were no deaths, the subsequent complications were much more frequent and troublesome.

The last point of difference that I shall mention is the fact that, while scarlet fever seldom reappears again in the same individual, diphtheria seems by its first attack to confer no immunity from subsequent seizures; on the contrary, the susceptibility of the throat seems to be rather increased.

Upon taking a survey of all these relations existing between scarlet fever and diphtheria, it is evident that they are distinct diseases, and yet that there is some very close and definite connection between them. Wherein the bond consists, it is not easy to point out. From the manner in which the diphtheritic poison seeks out the highest points of the best-drained localities, it has appeared to me most probable that the virus of this disease must be of a more rarefied and subtle nature than the germs that produce scarlet fever.

It is possible that the nature of some other ferments may throw some light upon this subject. I would specially call attention to the discovery by M. Berthelot (*Comptes Rendus*, vol. 1, p. 980) of an unorganised glucosic ferment in an infusion of yeast resembling diastase; also to the fact that one fermentation by organised beings frequently prepares the way for another. It may be suggested, that in one of these ways the singular relations existing between scarlatina and diphtheria might be accounted for. I do not venture to affirm that there is a strict analogy between these fermentative actions and the zymosis of scarlatina and diphtheria; but, as there exist in nature ferments requiring the presence of living germs, and others acting independently of them, so there are disease ferments with organic germs reproducing themselves, and others which have no definite term of life, and which are not self-reproductive.

It is probable that, in most cases of fermentation, the living being connected with each kind of this action is of a specific nature, and peculiar to the medium or to the food upon which it lives. In some instances, a ferment either during or at the end of its operation will produce the conditions favourable to the growth of another kind of germ, as seen in the crowd of vibrios appearing at the end of a mycoidermic fermentation; so the pyrexial state may produce the condition most congenial to the diphtheritic ferment, whether that may prove to be an organised germ or an unorganised catalytic ferment. As my excuse for offering these suggestions, I will quote, in conclusion, the words of the great English philosopher Robert Boyle. "He that thoroughly understands the nature of ferments and fermentation shall probably be much better able than he that ignores them to give a fair account of several diseases (as well fevers as others) which will perhaps be never thoroughly understood without an insight into the doctrine of fermentation." (*Works*, ed. 1744, vol. i, p. 476.)

MALARIA.

By T. INMAN, M.D.Lond.,

Late Lecturer on Medicine at the Liverpool Royal Infirmary School of Medicine; Consulting Physician of the Liverpool Royal Infirmary.

As I anticipate that my friend Dr. Oldham will, in due time, make his remarks upon what has appeared on this subject in the JOURNAL, I do not intend farther to prolong the controversy, especially as I find it difficult to do so without being satirical upon the logic of those who have supported the old idea of malaria.

For example, to my mind it appears strange that an ague-fit coming on after a gravel-walk has been raked, or a pit dug, must necessarily rise from a disturbance of the earth. I can recall plenty of cases in which ditches full of vegetable debris, stagnant ponds, and soil of all sorts, have been dug out and dug over. In one, this process was marked by an attack of sciatica in the master; in another, by gout; in another, by chicken-pox; in another, by pneumonia; in another, by a drunken bout; in another, by catarrh; in another, by paralysis; in another, by chorea; and in another, by gonorrhoea. Would not every one "write me down an ass" if I associated any one of these complaints with an upturning of sods, delving ditches, taking up potatoes, storing carrots, turnips, or mangold, or with other farm-labour? Then, if an attack of ague should come on, are we to be asked to jump to the conclusion that the complaint was caused by raking a gravel-walk, or digging a grave in the nearest cemetery? Again, all the correspondents have avoided the very important point, that emanations, whether from marsh or river, trench or pit, are the most abundant and incessant during the heat of the day, a period when nobody pretends—except in such cases when the body is suddenly chilled by a hailstorm, or by wading through cold rivers, or by a cold wind from the mountains, or simply by a drying zephyr when the body is covered with thin garments soaked with perspiration, rain, etc.—that agues are caught. The well known Maremma and Pontine Marshes and the Campagna around Rome are innocuous during the day, when the sun is hot and every emanation strong.

When night falls, and emanation ceases, then comes the danger of ague, etc. Clearly, then, the cause must be sought in a something which there is in common between nights in general and those conditions that will give ague during the day. This something, most unquestionably, is not malaria, or an emanation from any soil.

I commend the inquiry to those whose experience is commensurate with their acumen. I am quite aware how easy it is to try to put down a thinker by an array of so-called facts; and I have known a "stay-at-home" astronomer convict of inaccurate observations the whole of the officers of an exploring expedition, who insisted that their chronometers went wrong for three days, under the influence of a land-breeze from Australia. The head man was not so silly as to stick up for the malari-ous effect of proximity to land, but re-examined his calculations, and discovered the error.

Some of the writers in the JOURNAL seem wholly to have misunderstood the bad effect of sleeping on the deck of a ship being dependent upon (1) hard work and exhaustion under a hot sun; (2) sleeping where the body is certain to be chilled by radiation or a cold land-wind; and they make merry over the fact that many are taken ill between decks. Unquestionably they are, when all the air-holes are kept open, and admit a chilly land-breeze. Men on board ships in the deadliest African rivers escape fevers—as far as they are caught on ship-board—by closing the windows at sundown. At sea, the air in the tropics is equable day and night, when not chilled by a cold land-breeze. In like manner, one may cross the Pontine Marshes by night, by excluding from the carriage in which one rides the bitter cold; not that which we call bitter in England, but very severe to those accustomed to a day temperature of 140 degs. in the sun and 88 to 92 degs. in the shade, and who are too lightly clad to bear a sudden fall to 60 or 50 degs. during the hours of darkness. No amount of experience in the field can alter the physical fact that in many places in hot climates the nights are colder than the days, especially when the sky is clear. Even frost occurs occasionally by night on the Sahara; and the Arab always keeps his most valued horses under cover after sundown, and in good stables too.

Nor can any amount of experience, from the Indus to the Pole, convince us that very formidable diseases—*e. g.*, pneumonia and bronchitis—are not frequently produced by a severe chill after fatigue and exposure to heat. I should pay no compliment to the writers if I were to count *seriatim* all those complaints—even including purulent periostitis and necrosis—which have fairly been traced to violent chill after exposure to heat, without a microscope being able to detect in the air or water

to which the sufferers were exposed either fungoid spores or marsh malaria.

Once more I would add, that I have had the privilege of conversing familiarly with medical and other residents of Sierra Leone, Cape Coast Castle, Rangoon, Birmah, Fernando Po, the Gold Coast, the aguish districts of South America and the United States, the worst part of the West Indies, the fen districts of Cambridgeshire, and the worst parts of Italy. Some of my informants have had ague repeatedly; others have seen every one around them, including negroes, down with fever (I do not refer, of course, to yellow or other contagious fever), whilst they have escaped wholly during a residence of five years on the same spot.

The experience thus obtained is not to be despised, even although I have no personal knowledge of other malarious districts than those of Cambridgeshire and Italy.

THOUGHTS ON THE MANAGEMENT OF INCURABLE CHRONIC HEART-DISEASE.

By C. M. DURRANT, M.D., F.R.C.P.,
Physician to the East Suffolk and Ipswich Hospital.

I HAVE been induced to jot down a few passing reflections on the subject of this paper, from the circumstance that, while the treatment by drugs has undergone an almost exhaustive investigation by writers on heart-disease, its general management by moral and hygienic suggestions has been proportionately overlooked, and its paramount importance scarcely recognised.

The first question, then, that occurs to me—and a very important one it is—Do we, as practitioners, sufficiently inform our patient suffering from the consequences of an incurably diseased heart of the nature of his malady, with a view to his adopting such measures in the conduct of himself as may tend to ward off a suddenly fatal termination? The result of the daily inquests that are held will, I think, give a negative answer to this question.

A man uses sudden, and perhaps violent, exertion; falls; dies; an inquest is held. His usual medical attendant states that he has known (for some years possibly) that the individual has been the subject of an organic disease of the heart, and a verdict of death from natural causes is returned. The question arises, Had the patient been made aware of the precise nature of his complaint, and of its fatally hazardous character under the influence of sudden exertion? Too frequently, I believe, the individual undertakes the hurried movements which result in death in entire ignorance of his precarious condition; whereas, had he understood it, he might possibly, by his own prudence and care, have postponed the fatal termination. I am sure that we must all have been familiar with cases of sudden death from heart-disease, in which the patient, from a kind, but mistaken, feeling on the part both of physician and friends, has been left in absolute ignorance of his danger, or with only a vague conviction of something being wrong, yet without the remotest idea of what he should himself do to prevent a suddenly fatal catastrophe. The fear lest the shock, on being informed of his dangerous condition, produce the very result that we wish to avoid is, I believe, an altogether mistaken one. I am satisfied that no one with heart-disease will die one moment sooner for having been *properly* made aware of its existence; and I feel that we are all in duty bound, as honest responsible practitioners, and as Christians, to adopt the golden rule of doing as we should wish, under like circumstances, to be done by. If the patient be gently and kindly told by his medical attendant that, although his heart is seriously affected, still, if he carry out implicitly in his own person the rules laid down for his guidance, he may hope with confidence to have his life prolonged, I do not believe that anything but good can accrue from adopting this course. I am convinced that these suggestions have hitherto been very culpably neglected or insufficiently enforced, and that, if they were generally adopted, many cases of heart-disease now terminating with such painful suddenness might have life prolonged, and thus afford to the patient and to his family a lengthened interchange of social happiness. If, then, we admit the propriety of enlightening our patient as to his condition of danger, we have next to consider the especial influences which he will be called upon, for his own safety, to resist.

The first that will present itself to our minds will be the careful avoidance of sudden and hurried motion. Nothing is so likely to cause sudden death in advanced heart-disease as this. Hurrying to catch a train, attempting to walk rapidly up hill, especially in the face of a strong wind; dragging or lifting weights, and many other movements which will occur to the medical attendant to interdict, must be sedulously avoided by the patient. In cases of regurgitant aortic disease and great thinning of the ventricular walls, as well as in fatty de-

generation of their structure, this injunction will be doubly necessary. I believe that prolonged travelling by train may be highly prejudicial, and tend to a fatal termination by exhausting nerve-force. I have seen two cases of sudden death follow upon a railway journey, which I could only explain in this way. Both were subjects of advanced valvular disease.

Another very necessary precaution, and one not to be neglected by the patient without the greatest hazard, is the avoidance of partaking at any one time of a large distending meal, even though the food taken be of an easily assimilable character. This is a fruitful, but not a sufficiently recognised, cause of sudden death in heart-disease. This senile syncope, as it has been called, may be generally traced to the filling the enfeebled stomach beyond its power of disposing of its contents. Undigested food, acting as a foreign body, irritates the stomachal nerves; and, thus irritation being reflected upon the cardiac nerves, they in their turn fail to influence sufficiently the coronary vessels, and hence the cardiac syncope, and, if not relieved, fatal termination. In all cases of organic heart-disease, the meals, let it be remembered, should be small in quantity, of easy assimilation, and of frequent repetition; so that the heart may not be pressed upon by a distended stomach, on the one hand, or its due supply of nerve-force exhausted by prolonged abstinence, on the other.

Cold liquids, if taken in excess, and especially if quickly followed by exercise, and more particularly if this exercise be up an ascent, are very likely, by exciting nerve-irritation, to produce cardiac syncope and sudden death. In these days of almost universal tea-drinking, and at all hours, it requires some courage to inveigh against the popular custom; neither is it my intention to do so, beyond a few observations bearing upon the especial subject under our consideration. I think that the discrepancy in the minds of medical men in regard to the effects of tea, and the diametrically opposed opinions which are so freely enunciated, are much to be deprecated. Tea acts chiefly as a nerve-stimulant, and, when taken in moderation as to quantity, and not too strong, it can in no wise be injurious in heart-disease. When taken under these circumstances, it exhilarates and restores, and may often well supply the place of alcohol, with its subsequently depressing and intoxicating effects. With some persons, it should be remarked, tea will be found to act as a powerful nerve-irritant, amounting almost to a poison, and producing cardiac feebleness, flatulence, irregularity of the heart's action, and the many nervous disturbances which we must all have witnessed, even if we have not in our own persons experienced its discomforts. In such idiosyncracies, the use of tea and coffee must be strictly prohibited.

Another question arises, as to how far a patient labouring under serious heart-disease should debar himself from sexual intercourse. I believe, as a rule, that it should be absolutely forbidden, especially in the more serious conditions of heart above referred to.

A very troublesome symptom, difficult to combat, and attended with extreme hazard, is the great restlessness of some patients, the subjects of advanced valvular disease. If the patient have been in the habit of frequent travelling when well, we shall find the greatest difficulty in keeping him at home. Now, sudden death away from his own family circle, as I have more than once seen, is by no means unlikely to occur. The disturbed circulation, affecting the brain, and producing great irritability, renders these cases peculiarly trying and difficult to manage. Firmness, kindness, and the most truthful admission of the real condition of the sufferer, while the attendant dangers are distinctly set before him, will alone avail to influence him, and to keep him at home.

Passive carriage-motion is good; but, as a rule, it will be better, I believe, to debar the patient from horse-exercise. Early hours must be enjoined; and, if the patient be wakeful at night, nourishment in the shape of beef-tea with brandy must be taken, and especially in the early morning, when the nerve-force is at its lowest ebb. Milk, with rum or brandy, may also be taken shortly before rising. In all cases of advanced heart-disease, the patient, as a rule, should invariably carry a small flask of brandy in his pocket.

In reference to medicinal treatment, it is not my intention to dilate further than to state that, in mitral valve-disease, aperients freely given, with a view to unload the venous capillaries, will afford the most relief. In aortic regurgitation, I have often seen much advantage derived from the infusion of senega, with henbane and spirit of nitrous ether, as recommended by the late Dr. Barlow.

In senile syncope, the result of an overdistended stomach, no time must be lost in emptying this organ by an emetic, to be followed as quickly as circumstances will admit by the administration of full doses of brandy with hot water. In those cases of advanced heart-disease in which dropsy is superadded, and in which diuretics often signally fail, temporary relief may sometimes be obtained from full doses of quinine.