in several cases of comminuted fracture of the body of the bone, drilling the bone through the mouth or through a wound which already existed. Mr. Bickersteth of Liverpool (*Medico-Chirurgical Transactions*, vol. xlvii) has suggested and practised a method of securing the fractured ends of this and other bones by means of a drill, and has given a most interesting account of several cases in which it was successfully adopted. The employment of special external incisions, as in the case related, has rarely if ever been practised in connection with recent fractures of the lower jaw. The objection to them is the resulting external cicatrix or cicatrices which, although not of so much consequence in the male, as the whiskers will usually completely cover them, must cause some deformity in the female. For this reason, I would only advocate an external incision or incisions in cases which cannot be successfully treated by other means. In the removal of the tongue, or portions of it, and in some cases of tumour of the jaw, I always now bring the ends of the divided jaw together by means of the wire suture; and my experience of this method is, that it is safe, efficient, and easy of application. Since writing this, Mr. Mason of St. Thomas's Hospital has been kind enough to direct my attention to the ingenious method of employing the wire suture in bringing the end of bones together, practised by him (*Medico-Chirurgical Transactions*, vol. liv). His method allows the wire to be easily removed at any time, and is, therefore, to be preferred to that which has usually been employed, should the case admit of the needle being passed in an oblique direction through both ends of the bone.

ANOTHER CASE OF POISONING BY HOMŒOPATHIC SOLUTION OF CAMPHOR.

By GEORGE JOHNSON, M.D., F.R.S.,
Physician to King's College Hospital.

In a paper of mine published in the last volume of the Clinical Society's *Transactions*, there will be found notes of five cases in which symptoms more or less distressing and alarming were caused by homœopathic concentrated solution of camphor. I am indebted for the particulars of the following case to a former pupil, who gives me permission to publish them, but begs me to withhold his name, for the reason that it is a rule of the public service in which he is engaged that no medical officer shall publish his cases without special permission, which is not always granted when asked for. I give the case as reported in a letter addressed to me by the medical officer who saw the patient.

"I was called in to see a lady about thirty-five years of age, and found her with a very pale face and weak pulse. She said she had been suffering from a bad cold and feverishness; that she had got up late, and had her dinner at about two o'clock. About an hour after dinner, her sister, who is an experienced nurse, dropped seven drops of homœopathic solution of camphor on a lump of sugar, and gave it to her. Immediately after taking it, she was attacked with a very faint feeling, which compelled her to lie down flat on the hearth-rug; and she nearly lost consciousness. This lasted about five minutes. When I arrived, her face was very pale, and her pulse was weak. I ordered her to bed, and gave her some aromatic spirit of ammonia and lavender. She was very drowsy after she got to bed, but had no more vertigo or faintness; and in a day or two she got rid of her cold. I am led to believe that this was a case of camphor-poisoning similar to those which you have published, for the following reasons: the attack occurred immediately after taking a dose of camphor; the patient says that she had never before had a similar feeling of faintness; and the vertigo or faint was followed by drowsiness."

There can, I think, be no doubt that the symptoms were a direct result of the camphor. The dose was smaller than in any other case which I have met with in which poisoning has resulted from the incautious use of this dangerous compound. The homœopathic concentrated solution of camphor is a saturated solution of camphor in alcohol, the proportion being an ounce of camphor to an ounce and a quarter of spirit. Seven drops would be an uncertain quantity; but seven measured minims would contain rather more than five and a half grains of camphor. In one of the cases which I have recorded in the *Transactions* of the Clinical Society (vol. vii), a dose of twenty-five drops, taken by a young lady for a cold, caused violent convulsions and nervous sequelæ which continued for several months. In another case, eight doses of three drops each, taken by a clergyman within forty minutes, caused violent headache, sickness, and inability to work for two months. In a case communicated to the *BRITISH MEDICAL JOURNAL* by Dr. Clifford Allbutt (Dec. 6th, 1873), two doses of at the most ten drops each rendered a robust mountain-guide so giddy that he was unable to proceed on his journey, and he had to be deposited in a

cave for safety. Lastly, in two other cases, a teaspoonful dose caused convulsions and coma.

It is time that the public should be warned of the danger which they incur by the use of this poisonous homœopathic concentrated solution of camphor, which is more than seven times the strength of ordinary spirit of camphor. It is notorious that the modern disciples of Hahnemann have gone from the ludicrous extreme of infinitesimal dilution to the dangerous extreme of the greatest possible concentration of active and poisonous drugs.

ON THE RELATIONS BETWEEN DIPHTHERIA AND SCARLET FEVER.*

By ARTHUR RANSOME, M.A., M.D. Cantab.,
Lecturer on Public Health in Owens College, Manchester.

THE following case shows the close connection between these two diseases.

On February 23rd last year, a boy at school had scarlet fever. He was isolated until March 25th, when his brother saw him and took the complaint. Other cases occurred at the school on the 29th, 30th, and 31st. On April 2nd, one of the masters had a rigor and sore-throat; he left at once, and was seen by me next day. On the third day, diphtheritic patches covered the tonsils; but there was no rash. Slight albuminuria accompanied the pyrexia; both disappeared by the seventh day. On April 9th, six days after he arrived at home, his mother was attacked, membranous exudation appearing in both tonsils; but there was no albuminuria, and no rash.

The origin of these cases must surely be traced to the epidemic of scarlet fever at the school; for there were no cases of diphtheria either in the neighbourhood of the school or near their home. The close relationship of these diseases has often been pointed out. In the *Report of the Medical Officer to the Privy Council* for 1859, Dr. Burdon Sanderson gives cases of diphtheria and scarlet fever coexisting; there also I find three cases of diphtheria following three of scarlet fever, and the former spreading to a neighbouring cottage. These diseases have much in common. They are alike in their zymotic or epidemic character; both are characterised by high pyrexia; their chief point of attack is the throat and the glands of the neck; and a rash is found in both diseases. Their sequelæ also are similar: suppuration in the lymphatic glands, ulceration of the ears, arthritic affections with or without cardiac complication, kidney-disease and dropsy, general or localised paralysis of the nervous system. I have seen each and all of these consequences after both diseases.

Some physicians have been led by these points of resemblance to regard them as one disease, the same poison producing different manifestations; but, in spite of their similarity, I venture to think the differences are too important and too numerous to permit so simple a solution of the problem. In the first place, diphtheria is seen to arise directly from scarlet fever. I have never known the converse action. There are other etiological differences. Though both are contagious, diphtheria is less so, and more frequently occurs in single cases. Where several cases occur together, some general cause is probably concerned. Scarlet fever is more frequently conveyed by clothes or other fomites, by attendants on the sick, and it may be conveyed by means of milk. I am not aware that diphtheria has ever spread by any of these means. The two diseases differ in their geographical distribution. Diphtheria is common in India, where scarlet fever is unknown. Colonies and all new settlements seem peculiarly liable to diphtheria; it has appeared in the Australian colonies and in South America, attacking the otherwise healthy town of Buenos Ayres. In Lima, it was noticed that the black races withstand its contagion. It selects high and well drained districts, when low-lying lands in the vicinity escape. Dr. Blake, in the *Transactions of the Medical Society of California*, mentions a most fatal epidemic at an elevation of 4,000 feet. Many places in England subject to diphtheria are either high or well drained. Scarlet fever also haunts certain places; but these scarlet-fever fields are not at high levels, nor are they those in which diphtheria is especially prevalent. Diphtheria is often connected with defects in the house-drainage, not always due to imperfection in the house-connection with a general sewage-system, but to leakage from private cesspools; so that a question of spontaneous origin may arise in this disease, which is hardly admissible with respect to scarlet fever. Season has a different influence on the two diseases; scarlet fever being most prevalent in the autumn, low barometric pressure and greater humidity in the air favouring its diffusion; while diphtheria may arise at any season, and, in my own experience, its most virulent epidemics have occurred in the summer months.

* Abstract of a paper read before the Epidemiological Society of London, January 13th, 1875.