ANOTHER CASE OF POISONING BY HOMŒOPATHIC CAMPHOR.

By GEORGE JOHNSON, M.D., F.R.S.,

Professor of Medicine in King's College; Physician to King's College Hospital.

I AM indebted for the particulars of the following case of poisoning by homœopathic solution of camphor to my friend Mr. Gooch of Eton. He had read my last paper on the subject in the JOURNAL (Feb. 6th, 1875), and, thinking that the case of an Eton boy whom he had attended some time since would interest me, he has been so good as to send me a brief report, with permission to publish it in the BRITISH MEDICAL JOURNAL. The following is Mr. Gooch's narrative.

"About three years since, I was called to see an Eton boy, aged 14, who was supposed to have fainted. I found him lying on his bed, almost pulseless, and quite insensible, with his extremities cold, and his face and lips pallid. He had shortly before had his dinner, and I thought that, perhaps, he might be suffering from an attack of indigestion. Whilst I was administering stimulants, and applying hot bottles, etc., to his feet, a servant accidentally found a small bottle on his table, which was labelled, "Rubini's Homœopathic Camphor"; and, thinking that he might have taken some of its contents, I gave him, with difficulty, an emetic. This had the immediate effect of causing him to vomit his dinner, which was mixed with an immense quantity of camphor, which scented the whole room most powerfully. In about half an hour, he became sensible, and told me that he had a cold, for which he had purchased the camphor; but, instead of dropping two or three drops upon a lump of sugar, as directed, he poured it out on the sugar, and after taking it, immediately became insensible. In the evening, all the effects of the drug had gone off; he passed a good night, and had no unpleasant symptoms afterwards.

"He was a strong, healthy-looking boy. The bottle found in his room was very small, holding about half a drachm, and it was about half full when I saw it; so that he may have taken about fifteen drops of the solution, if an ordinary lump of sugar would hold so much."

This is the seventh case of poisoning by the homœopathic concentrated solution of camphor that has come to my knowledge during the last two years. I suspect that, if all the cases of this kind that have come under the notice of the profession within the United Kingdom during the same period were published, the list would be a very long one, and it might serve as a warning of the danger which attends the incautious use of this highly concentrated poison.

Since the above was written, I have read with much interest the two cases published in the last number of the JOURNAL (p. 243), by Dr. Legat and Mr. Ellerton.

ON SOME POINTS IN THE DIAGNOSIS OF BRIGHT'S DISEASE.

By CHARLES MACLEAN, M.B., Applecross.

IT may appear a work of supererogation to speak of the powers of the microscope in the investigation of renal disease, but we really cannot be too much impressed with the value of that instrument in diagnosis; and the chief object of this paper is to show how it can be made the principal medium in the performance of a delicate test towards the detection of true albuminuria when the urine contains blood. Bright's disease may easily be complicated with hæmaturia—indeed, it is often so; and it frequently becomes a question, whether the albumen in the urine of a patient is or is not to be explained by the presence of effused blood. We have, in fact, to find out how much blood-albumen the urine contains, and how much albuminuric albumen. The former can be estimated by counting the number of red blood-corpuscles to be seen on a given extent of the microscopic slide; it being known how much albumen, say in the ounce of urine, shall correspond to a given number of discs. The amount of albumen in the urine is easily found by coagulating, filtering, and drying that contained in a like quantity of urine (an ounce), and weighing it.

I find that, when forty blood-discs are to be seen in the whole field of an "Oberhauser" microscope, using object-glass No. 7 and eyeglass No. 4, if there be effused blood in the urine, the albumen contained in an ounce, on being dried, weighs one grain; therefore, whatever there is of dried albumen from the ounce of urine, over and above the proportion of one grain to the forty blood-discs, seen as above, must be the product of true albuminuria. I have a case of granular disease of the kidney in which the albumen obtained from one ounce of the urine, independently of blood, weighed eight grains when I saw it first.

It would be as fallacious to look upon coagulation by heat and nitric acid as conclusive of a case being one of albuminuria, without taking other circumstances, such as the presence of blood, etc., into account, as it would be to exclude the possibility of that disease being present merely because of the urine containing blood. It may be mentioned, that the urine has a distinct tinge of blood when as few as ten discs are to be seen on the field of the microscope. I had occasion to examine the urine of a lady, a visitor in this neighbourhood, who, before she left home, received a note from her medical attendant saying that she had had a severe attack, in which the noticeable symptoms were dark smoky-looking urine, which contained blood-discs, and which coagulated distinctly on the application of heat—essentially a tautological statement. The coagulation was twice referred to as an important point; whereas I think it quite likely that, if a test of the kind described in the foregoing remarks had been applied at the time, it would have been found that hæmorrhage sufficiently accounted for all the coagulation. I found the urine quite natural in every way, except that it contained great numbers of crystals of oxalate of lime; and the conclusion to which I came was, that the bleeding had been caused by a concretion of oxalate of lime forming in the kidney.

The importance of examining the urine in the diagnosis of disease cannot easily be exaggerated. The facility with which we obtain evidence in this way, and the certainty of the information when obtained, make the omission on our part of this means of diagnosis the more inexcusable. I was recently called to a case, in which the only observable symptoms were epigastric pain, nausea with feeling of great weakness, and a subacute bronchial affection. As the latter complaint did not appear to me to be capable of accounting for the constitutional symptoms, I examined the urine and found it loaded with albumen; and the microscope revealed numerous granular casts. All doubt as to the nature of the disease was thus at once dispelled. There was here not the least lumbar pain complained of, and no unusual feeling on micturition.

I had an instance of the great value of the microscope some time ago in a case of hæmaturia. It was an alarming case; and life seemed at the time in danger from the large quantity of blood lost, for pint upon pint of bloody urine (or, more properly speaking, of diluted blood) was passed. I gave the patient gallic acid to begin with, until I should examine the urine microscopically. This treatment made little or no impression, however; and, when I brought the microscope into use, I found crystals of the ammonio-magnesian phosphate in abundance. Thereupon I combined sulphuric acid with the gallic, and the almost immediate result was an abatement of the bleeding. The mineral acid probably dissolved any irritating particles of the triple phosphate that may have formed. The patient had a milder attack afterwards, but it yielded readily on the application of the same remedy.

THERAPEUTIC USE OF THE CONTINUOUS GALVANIC CURRENT.

By THOMAS E. CLARK, M.D.,

Late Physician to the General Hospital, Bristol; and formerly Surgeon to the Royal Infirmary, Bristol.

I SEND three cases treated by myself by the continuous galvanic current, believing that it is a subject of interest to the profession.

CASE 1. *Endometritis cured by the Continuous Galvanic Current.*—S. H., aged 39, a married woman, has had four children. After her fourth confinement, she suffered severely from leucorrhœa, bearing-down pains, tenderness in the left iliac region, and dysmenorrhœa. This continued for six months, when she was able to go about the house; this was in 1870. I saw her in June 1871, and, upon examination, found the fundus uteri lying obliquely and to the left, the os uteri patulous, and superficially ulcerated; and, during the introduction of the speculum, a porter-coloured fluid of a very offensive character flowed from the os. Under the local use of nitrate of silver, and the internal use of bromide of potassium, iron, and sumpbul, with rest, she became very much better, and indeed reported herself well. However, in the November following, she caught a severe cold, and all the old symptoms recurred. She was now confined to her room and bed, and no treatment seemed to have any beneficial effect upon her case. I saw her in June 1872, and ordered her away for a time to recruit her shattered nervous system, and at intervals of a week, applied the nitrate of silver; she also continued the bromide of potassium and iron mixture. At the end of August she returned home, and I recommended the continuous current, with the idea of altering the secreting property of the lining membrane of the uterus. Accordingly, I introduced a partly insulated wire into the cavity of the uterus, and attached