THE CROONIAN LECTURES ON ADDISON'S DISEASE.

Delivered at the Royal College of Physicians, London.

By E. HEADLAM GREENHOW, M.D., F.R.C.P., F.R.S.,
Physician to and Lecturer on Medicine at the Middlesex Hospital.

LECTURE II.

[Concluded from p. 371 of last number.]

IT will be remembered that, in dealing with the total number of three hundred and thirty-two cases bearing upon the subject of Addison's disease, I set aside, in the first instance, for subsequent discussion, all those which I found reported as bronzing or discoloration of the skin without disease of the suprarenal capsules. These cases are twenty-one in number; and a brief analysis will show that, in almost all of them, the abnormal pigmentation was due to one or other of the pathological conditions which I have mentioned.

Two of the twenty-one cases published by Vernois and Averbek precisely resemble the case of the woman exhibited by me at the Pathological Society; the age, circumstances, uncleanly habits and state of health, as well as the characters of the discoloration, having been nearly identical in all the three patients. The patients in these two cases died, and the capsules were found to be normal. It is significant of the confusion which prevails on this subject, that Averbek, who did not consider his own case as a true example of Addison's disease, nevertheless tabulated the parallel case of my patient, who recovered, among the so-called cured cases of that disease—which, as he justly observes, must be received with suspicion. In three other cases published by Puech and Gordon, there was constitutional syphilis. In two cases recorded by Sloane and Luton, the darkening of the skin was associated with phthisis. The first patient had suffered from liver-complaint with permanent discoloration of skin; and, although the whole surface was bronzed, the mammæ round the nipples were much paler than the surrounding parts, and there were many white spots on the breast and back. In two other cases related by Gibb and May, there was disease both of the lungs and the heart. Dr. Gibb's patient had aortic disease, which sufficiently accounted for the pale bronze hue of the face and neck; whilst in Mr. May's case the patient had also had gout, and the small brown spots scattered over the normal coloured skin point to the possible remains of some gouty eruption. In four cases published by Hutchinson, Harley, and Fricke, the discoloration dated from attacks of fever, jaundice, and hepatic derangement. Mr. Hutchinson's patient died of pneumonia, after an illness of fourteen days. His skin had been dark from the time he had had yellow fever in the Crimea. The entire surface was more or less bronzed, and the pigment was chiefly situated in the rete mucosum, but the face and hands were the lightest parts; whereas I need not say that they are generally the darkest parts in Addison's disease. In one of Dr. George Harley's cases, the darkening of the skin followed an attack of jaundice, and the patient remained five years in good health. The face, neck, and arms were dark; but the discoloration of the body was diversified with irregular white patches. The patient died of diseased liver and ascites. In Dr. Harley's other case, the patient had suffered during four years from occasional hepatic symptoms and gastric disorder. The face, arms, and trunk were all of a dark bronze colour, except the scar of a burn on the breast; but apparently the discoloration was quite uniform, and, unfortunately, the condition of the liver was not noted. In Dr. Fricke's case, the bronzing was limited to the forehead, face, and neck, and merged into jaundiced skin. The patient had suffered from ague, and died of cirrhosis of the liver. A case allied to these, but much more complicated, is reported by Dr. Merkel. The man's skin had been always dark; he had suffered from ague, and died of bronchitis and pleurisy. He was found, on *post mortem* examination, to have a large liver and spleen; and there was fatty degeneration of the heart, liver, and kidneys. The whole surface was bronzed, equally on covered and uncovered parts; and there were large grey-black patches on the buccal mucous membrane. Two cases of discoloured skin without disease of the suprarenal capsules, recorded by Dr. Bucknill and M. Regnard, occurred in women who were maniacal at the climacteric period of life—circumstances in which extreme duskiness of the skin is by no means uncommon. Two cases of internal cancer, not involving the suprarenal capsules, are reported by Mitchell and Fremy as presenting a certain degree of discoloration of skin;

though of the first, it is only said that the skin became of a leathern hue, much darker in eczematous scars and in the site of a blister than in other parts; and of the second case, that, some days before death, the skin assumed in places an evident bronze hue. Dr. Hodges found on the bodies of two old men, one of whom died insane, and the other of age and debility, general darkening of the skin, with a bronze hue on the genital organs; but the face and other parts were much less discoloured.

There still remains one case of bronzed skin without disease of the suprarenal capsules, published by Rossbach in Virchow's *Archiv* for 1870, which requires more special notice. The case is most fully reported, and, on the supposition that it is a true case of Addison's disease, is made the basis of a new theory on the pathology of that disease, to which I must hereafter revert. The patient, a lady aged 62, had always been prone to excitement and sleeplessness; and these natural tendencies had been aggravated by long continued distress of mind. She suffered from anæmia, debility, loss of appetite, occasional vomiting, and also from pains in the epigastrium and in the right scapula, which she had hurt by a fall. These pains gradually extended round her body with all the violence of the neuralgia of shingles, and were always worst at night. Scleroderma began to develop on the right hand and forearm, and spread more or less to other parts of the body. The skin and subcutaneous tissue became as hard as stone, split, and formed fissures; and, together with this rare and peculiar cutaneous affection, appeared a dark and extensive pigmentation of the skin, of which large patches sometimes developed in a single night. Meantime, the patient's restlessness of mind and body increased; she could bear no position in or out of bed for fifteen minutes together, and at length she became totally sleepless and actually deranged at night. Towards the close of life, her pulse was quick, hard, and occasionally intermitting. She had albuminuria and anasarca, and sank under a slight attack of pneumonia, after nine months' illness. The cerebellum was very soft; the brain and lungs were œdematous; and there was some fatty degeneration of the heart, liver, and kidneys: but the suprarenal capsules and neighbouring nerves were perfectly normal. The pigmentation of the skin was coextensive with the scleroderma. In the paler parts, it was produced by innumerable dark points, and on the darkest parts there were dendritic patches of colourless skin. On examination under the microscope, the pigment was seen to be deposited chiefly in the papillæ and upper layers of the cutis. I need not point out in detail the totally opposite characters of this discoloration of skin from those of the discoloration in Addison's disease. The chief constitutional symptoms—namely, the morbid excitement of mind and body—were also conditions totally uncharacteristic of Addison's disease; whilst the anæmia, loss of appetite, and other symptoms, which did resemble some of those of Addison's disease, were all fully accounted for by the obvious mental and physical ailments of the patient.

From this analysis of the twenty-one cases I have found published as cases of bronzing of skin without disease of the suprarenal capsules, I think it is evident that in all of them the discoloration of the skin was due to other pathological causes, and that it presented in none of them the really characteristic features of the bronzed skin of Addison's disease. I should add, what is even more important, that in none of these cases are any constitutional symptoms recorded but such as are easily referable to the morbid conditions I have mentioned as present in each. They are of course only representative cases, as others of similar characters are continually occurring, though I have not found them brought forward with reference to the question of bronzed skin and suprarenal disease.

The second allegation against the truth of Addison's discovery, viz., that disease of the suprarenal capsules not unfrequently exists without discoloration of the skin, is also, as I have said, perfectly true. Apart, however, from the small number of cases which have terminated before the development of the bronzing, it may be positively enunciated, that disease of the suprarenal capsules which is not attended with discoloration of skin, presents totally different characters from the particular lesion of those organs which I have described. Out of the thirty-seven cases which I have found published as examples of disease of the suprarenal capsules without bronzing of skin, twenty-four are cases of cancer of one or both capsules, invariably secondary to cancer of other neighbouring organs. In three of the other thirteen cases, the capsules had undergone fatty or amyloid degeneration. In one case, there was cirrhosis of the capsules, associated with kidney disease. In three cases, there were cysts; in three, apoplexy; in three, sarcomatous, fatty and plastic tumours, respectively; and, in seven cases out of the nine, only one of the two capsules was diseased. It is obvious that none of these lesions bear any resemblance to the one particular lesion of the suprarenal capsules which is attended by discoloration of skin; nor were any constitutional symptoms present in any of these

cases except such as were due to disease existing in other organs. I have myself, moreover, seen in the Middlesex Hospital several cases of cancer, in which the suprarenal capsules were involved, some of which I had had the opportunity of observing during life. The structure of the capsules was entirely destroyed in all those cases, yet in none of them had there been any constitutional symptoms or discoloration of skin resembling those of Addison's disease.

Hence, therefore, although it is quite true that bronzed skin does occur without the existence of any disease in the suprarenal capsules; and, again, that disease of the suprarenal capsules does exist, to the extent of entirely destroying their structure, without producing any bronzing of skin; it is also, on the other hand, equally clear that, in the one case, the bronzing of skin is not the peculiar discoloration indicative of the existence of Addison's disease; and that, in the other case (with the few exceptions already explained), the lesion in the suprarenal capsules is not the particular lesion of those organs which constitutes Addison's disease.

Were these the only misconceptions with regard to the relation of suprarenal disease to discoloration of skin, I might here quit this part of my subject; but, unfortunately, a small number of misunderstood cases have given a certain degree of prevalence to another opposite misapprehension, viz., that "any disease of the suprarenal capsules involving destruction of their tissues will produce the same effects as the particular lesion characteristic of Addison's disease".

Had not attention been too entirely directed to the bronzing of skin, to the neglect of the constitutional symptoms, and of Addison's original reservation that other pathological causes must first be excluded, such a misconception could never have arisen. It began, however, with Addison himself, who published four cases of cancer of the suprarenal capsules, as having been attended by some discoloration of the skin. I have also found published since his time, five cases of so-called cancer of the suprarenal capsules, and five cases of other diseases of the capsules, associated with bronzing of the skin. A glance at these fourteen cases will show that they afford no true cause for this misconception.

In the first of Addison's four cases, the patient was a woman aged 60—a time of life when the skin, even in healthy subjects, is apt to assume a dry brownish appearance; and she was suffering from extensive cancer of the left breast, lung, and surface of the liver. The only discoloration noted was a light brown swarthy hue on the face, arms, and chest. The second case was that of a woman, aged 53, who died of cancer of the stomach. She had previously had an eruption on the body; her skin was harsh, dry, and of a darkish hue; the folds of the axillæ were dark, and dark-coloured patches of the size of the palm of the hand were observed, raised in wrinkles and resembling slight ichthyosis. The third case was that of a woman, aged 28, who died of cancer of the uterus, and the only discoloration noticed was a peculiar dingy appearance of the skin, not very strongly marked. The fourth case was that of a man who died of cancer of the thorax and the lungs. When cancer was unexpectedly found in one of the suprarenal capsules, it was ascertained from the clinical notes, that the patient's face had presented a dingy hue. Addison had seen none of these cases during life, and knew nothing of their history or symptoms. Two more recent cases of cancer of the capsules, with discoloration of skin, are recorded by Drs. Gibb and Cayley. Dr. Gibb's patient had cancer of the liver and right capsule, associated with a proliferous ovarian cyst, and the only darkening of the skin observed was a brownish discoloration of the abdomen, especially below the navel; and of the chest, especially over the sternum. Dr. Cayley's case was that of a woman, aged 45, who died of cancer of the uterus, with secondary deposits in most of the abdominal organs. There was only a small nodule of cancer projecting from the surface of the suprarenal capsule; but during the last month of life the surface was very dusky and sallow, especially on the face and neck. I need scarcely point out that, in none of these six cases, was the discoloration observed in any degree characteristic of Addison's disease, and in none of them were any constitutional symptoms present, but such as were due to the disease in other organs.

The three remaining cases of so-called cancer of the capsules, with discoloration of skin, are of a very different character, having all presented, in a typical form, the constitutional symptoms and bronzing of skin described in my first lecture; and having been, in fact, as I believe, three typical cases of Addison's disease. In the first of these cases, Dr. Mettenheimer merely says, that the capsules were replaced by cancerous deposits; but he was so little clear as to the nature of such deposits, that he further reports the presence of some tuberculous or cancerous indurations of the left lung; and I think, as no cancerous disease was found in any other organ, there is no doubt that the deposits in the capsules were also of the so-called tuberculous character. Dr. Duclos, in the second case, says that the disease in the capsule was perfect scirrhus, but he had apparently made no microscopical examina-

tion, and his account of the condition of the organs is, on the contrary, a perfect description of the lesion peculiar to Addison's disease. The capsules formed tumours of the size and shape of eggs, were very hard, and much nodulated; and, on section, they grated against the scalpel and presented the appearance of lardaceous tissue. The third case, which is published by Dr. Gage of Worcester, United States, had a history which evidently led to the inference that cancer existed in the capsules, though it is not expressly asserted. The patient, a lady, aged 51, was said to have had scirrhus of the right breast, which, though of several years' growth, had not involved any of the neighbouring lymphatic glands, and had been removed two years before her death without recurrence. Her last illness was of six months' duration, and both the constitutional symptoms and the bronzing of the skin were characteristic of Addison's disease. Both the capsules were much enlarged, and joined together by a thickened mass of enlarged and diseased lymphatic glands. The right capsule was adherent to the under surface of the liver; and it is stated that from the point of attachment several broad lines of reddish-white deposit, resembling soft cancer, radiated into the substance of the liver for one inch and a half. In this case, also, there seems to have been no microscopical examination of the diseased organs; and, although the description is less clear than in the previous cases, the conviction that the disease was cancer having evidently been present to the mind of the writer, it bears on the whole a greater resemblance to Addison's disease than it does to cancer. Besides, it must be borne in mind that, if these three cases were indeed cases of cancer of the suprarenal capsules, they are the only cases on record in which cancerous disease of those organs has ever given rise to the constitutional symptoms and bronzing of the skin characteristic of Addison's disease, or has ever been attended by fatal results without the implication of other important organs—circumstances so incredible that they seem to justify the hypothesis which I have ventured to form concerning their real character.

Of the five cases of other diseases of the suprarenal capsules accompanied by discoloration of the skin, two are cases of amyloid disease. In one of these, published by Auerbeck, the patient, a man aged 53, presented a dirty grey brown colour of the skin, deepest on the shoulders and back, whilst the face was of a lighter hue. In the other case, recorded by Wolff, the patient had suffered for several years from a large hydatid cyst in the left kidney. A year before death, the face and forehead had assumed a dark sunburned hue, but the mucous membranes were anæmic, and the only signs of discoloration on the body were a few black spots on the neck, abdomen, and extremities. The debility, loss of appetite, vomiting, and epigastric pains, present in this case, were all, as well as the emaciation, obvious results of the general abdominal disease, and increasing discharge of foetid pus. In the remaining three cases, there was fatty degeneration of the capsules. One, related by Sir H. Thompson, was the case of an old woman, aged 81, who died of bronchitis. The skin was bronzed except on the face and hands, and the whole skin was permeated by colouring matter. In another of these cases, recorded by Charcot, the patient, a man aged 57, showed brown patches on most parts of the body, with a slight lichenoid condition of skin; the face, hands and feet remained of normal colour. The third case, published by Schet, is worthy of fuller notice, because the discoloration of the skin was, according to the description, identical with that seen in Addison's disease. It dated, however, from nine years' work under the glowing sun of a tropical climate, where the man had suffered from repeated attacks of intermittent fever, which continued at intervals during the four years he lived after returning to Europe, when he died of an epileptiform attack following copious spirit-drinking. The spleen was found to be much enlarged, and the blood contained pigment-granules and an excess of white corpuscles. The discoloration in this case, therefore, was doubtless due, in part to the effects of climate, but mainly to melanæmia, arising from malarious poisoning and enlarged spleen. It seems tolerably clear that these cases afford no real ground for the misconception that any disease of the suprarenal capsules, involving destruction to their tissues, may produce the same effects as Addison's disease of those organs.

Another misconception, nearly allied to the last, appears to exist in some quarters, to the effect that any abnormal condition of the suprarenal capsules is sufficient to raise the question of the existence of Addison's disease. Under the influence, apparently, of this misconception, a considerable number of cases have been published, in which the descriptions of the condition of the suprarenal capsules are too vague to justify any conclusions as to the nature of the lesion; whilst, in others, it seems doubtful whether there was any pathological lesion at all, and not rather some *post mortem* change. A few cases, also, of doubtful and anomalous affections of the suprarenal capsules have been published, as, for instance, one by Dr. Tuckwell, to show the necessity of caution in forming a diagnosis of such cases.

It is impossible to discuss in detail the thirty-two cases I have placed in this group; but it is certain that the characteristic features of the suprarenal lesion in Addison's disease are not described in any one of them. I must, however, advert to the most recent of these cases, published by Dr. Wickham Legg, in the last volume of the *St. Bartholomew's Hospital Reports*. The patient, a woman aged 37, unquestionably exhibited some of the constitutional symptoms of Addison's disease, for she suffered for eighteen months from debility, shortness of breath, occasional vomiting of food, and constipation, and she died after five weeks' residence in the hospital, of vomiting and exhaustion. But, on the other hand, there was a distinct bilious tendency to explain the vomiting; the appetite was fair, and there was great thirst, with a red fissured tongue, swollen spongy gums, and a temperature above the normal. Moreover, the patient had lost four stones in weight during her illness: an absolutely unprecedented circumstance in Addison's disease, unless when resulting from some serious coexisting wasting disease. The discoloured skin, also, was harsh, dry and warm, instead of being soft and cool, and was not more deeply pigmented in the axillæ or areolæ of the nipples than at other parts. At the *post mortem* examination, the right suprarenal capsule could not be found, its place being filled by a mass of yellow fat; and the left was only represented by a black body as thin as paper. This was a most peculiar and interesting condition of the suprarenal capsules, but it obviously bore no relation whatever to Addison's disease of those organs; nor were the constitutional symptoms and discoloration of skin, taken as a whole, characteristic of that disease. The most difficult point in the case is the absence of any apparent cause of illness and death.

One case remains of the whole number collected, which represents another misconception so remarkable, that it cannot be entirely passed over. The case is related by Dr. Gilliam of the United States, and is that of a man aged 45, who suffered from many of the constitutional symptoms of Addison's disease. Gradually increasing debility, dyspnoea and palpitation on exertion, loss of appetite, and tendency to vomiting, which became more and more obstinate until the patient died of exhaustion—were symptoms which, in the absence of any other discoverable cause might fairly lead to a diagnosis of Addison's disease, although no discoloration of skin was present. But when, after death, not only were the suprarenal capsules found to be healthy, but advanced degeneration and atrophy of the gastric tubules were discovered, which, as the author himself observed, fully accounted for all the symptoms, the diagnosis had clearly been a mistake. From this case, however, Dr. Gilliam concludes, first, that the bronzing of the skin is quite an unimportant feature of Addison's disease; and, secondly, that Addison's disease itself consists, not in a lesion of the suprarenal capsules at all, but in disease of the gastric tubules. The misconception lying at the root of this singular theory seems to be, that the presence of constitutional symptoms resembling those of Addison's disease was, in itself, sufficient evidence of the existence of that disease, and that the lesion which produced them might safely be regarded as the essential lesion constituting Addison's disease. That, in the case referred to, the lesion found in the stomach was the cause of the gastric and other symptoms, is obvious enough, but I trust I have made it equally clear that no symptoms, obviously referable to other morbid conditions can justify any assumption as to the existence of Addison's disease.

The question naturally arises, how misconceptions, so baseless in fact and disproved by such conclusive evidence, can have been so widely propagated and have survived so long. I believe this has been mainly due to the great rarity of the disease, which has prevented all but a small minority of medical men from acquiring even the degree of practical acquaintance with its true characters which the observation of a single typical case would afford.

If, however, the relation between the clinical phenomena and the specific suprarenal lesion in Addison's disease may be considered as proved, the obscure pathological processes involved in that relation, and the equally obscure causes of the suprarenal lesion itself, will still afford ample field for investigation. To these parts of the subject, I must endeavour to address myself in my next lecture, although perfectly conscious of the insurmountable difficulties they present, in the present vague and imperfect state of our knowledge with respect to many essential facts.

ACUTE GLOSSITIS IN SCARLATINA.—An instance of this extremely rare complication has lately been published by Dr. Wm. Moore, of Dublin. It occurred in a young gentleman aged 18, and was the only example Dr. Moore had ever seen in this disease. He strongly advocates the treatment by iron and stimulants, when this complication takes place. A somewhat similar case has been described as occurring in the practice of Dr. Banks, but the complication is unusual.

THE GOULSTONIAN LECTURES ON PUERPERAL FEVER.

Delivered at the Royal College of Physicians, London.

By ROBERT J. LEE, M.D., F.R.C.P.,

Assistant Physician at the Hospital for Sick Children, Great Ormond Street.

LECTURE III.—(Continued.)

WHEN Professor Lister remarks, "We now see it is not essential to assume the existence of a special virus at all, but that organisms common to all the sores in the ward may, for aught we know, assume specific properties in the discharges long putrefying under the dressings" (*Microscopical Journal*, vol. xiii), it is clear that he is indisposed to allow that a poison apart from bacteria can be brought into existence. It is not really because these points, so difficult at present to decide, are of practical importance that I refer to them. But they are interesting, as showing pretty clearly where the various parties of investigation have started from in search of fresh knowledge. We have claimed the privilege of testing all reports thus returned by the rules of general experience, supposing there is any great difference of opinion as to their correctness. There has been little occasion, however, to do this as yet; for the most valuable contributions we have received are the results of the labours of men of practical experience in the clinical phenomena of the disease produced by septic infection. It seems to me that, for this reason, we may accept the conclusions of Virchow, Lister, Billroth, Heiberg, and others, who are quite able for themselves, and are in the habit, of applying to their researches the test of clinical experience.

There is one case, for instance, in which we might find it difficult to explain, by the theory of bacteria, certain morbid phenomena, if we confined ourselves to the limited effect which the particles produce on the tissues: those changes, namely, which we term inflammatory. It is well known that there are cases in which we find no collections of pus, or secondary abscesses, and no well marked pathological alterations, cases usually of the acute form of puerperal fever, and where the constitutional symptoms have been the chief evidence of septic infection. In such instances, morbid anatomy reveals conditions not unlike those which we meet with in certain other contagious diseases: enlargement and softening of the spleen, congestion of the mucous and serous membranes, fluid condition of the blood, and slight general congestion of the pulmonary tissue. It is in these cases that Heiberg felt a difficulty in retaining the use of the term pyæmia; for there is neither a primary focus of purulent matter, nor are there any evidences of the presence of pus in any part of the body.

This is the reason why Professor Billroth uses the term *septhæmia* to signify the resorption of septic stuff, and pyæmia the resorption of pus. We have also septicæmia contrasted with icorrhæmia in some recent German works. These terms are intended to separate cases of acute septic infection from those of longer duration; for, as it is well known, in cases of abortion, retained placenta, and others, where the origin of the decomposing substance is not very doubtful, we observe a series of symptoms which lead to the belief that a slow process of poisoning is going on in the system of the patient. Such a process would be termed *ichorrhæmic*.

If we follow Professor Heiberg, as certainly we are inclined to do, on the grounds of clinical experience, we should regard these puerperal processes, in whatever form they present themselves, and whatever characters the symptoms and pathological changes may assume, as depending on one and the same cause, the action of which on the nervous system, the organs of circulation and respiration, on the temperature, and on the fluids and tissues of the body, may be modified by circumstances similar to those which we know to be capable of modifying the action of poisons generally.

For some reasons, therefore, we might feel disposed to rest satisfied with the old term putridity, which involves no such theory as septicæmia, pyæmia, and others; or we might fall back on the simple classical term sepsis as a convenient one for general use. In the clinical observation of cases in which septic agents are more or less actively at work in the system, whether in simple puerperal fever, in cases of abortion or retained placenta, in carcinoma uteri, or under any other circumstances, we are led to divide our attention between local