nor faintness on exertion, has ever been absent in the later stages of the disease. But there is not perhaps one of the other peculiar symptoms which is invariably observed. Every physician is sufficiently aware that this is equally true with respect to every other organic disease. It is only in the comparatively small number of cases in which an organic disease runs a long uncomplicated chronic course, that we can look for the full development of all its peculiar symptoms. The absence of some of these, therefore, in complicated cases, or in cases of short duration, cannot be regarded in Addison's disease, more than in any other organic disease, as militating against the connection between such symptoms and the pathological conditions with which they are associated.

THE GOULSTONIAN LECTURES ON PUERPERAL FEVER.

Delivered at the Royal College of Physicians, London.

By ROBERT J. LEE, M.D., F.R.C.P.,

Assistant Physician at the Hospital for Sick Children, Great Ormond Street.

LECTURE II.—(Concluded.)

THE interval between Gordon and Semmelweiss of half a century was occupied chiefly by those important investigations into morbid anatomy and pathology, both general and microscopical, which have contributed to the improvement of every department of our profession, and particularly to the practical knowledge of diagnosis and etiology. It is to this source that we must look for modifications in general opinion concerning some of the phenomena of puerperal fever; and it will be, I think, most convenient to defer our inquiry into the value of Semmelweiss's observations till we have reviewed the progress of pathological research. The real reason why there was such an universal movement in that direction about the end of last century may be easily traced to a few powerful minds, like the two Hunters and those nearly associated with them, who clearly perceived that the pathological study of the effects of disease promised more than the study of symptoms.

The result of this method, as far as it influenced the subject of puerperal fever, was the introduction of several new terms into our nosology which were intended to describe the pathological conditions of different organs, structures, and tissues. Thus originated puerperal peritonitis, phlebitis, metritis, and some others. These terms, as we know, did not include any explanation of the remote cause which induced the changes observed by the pathologist. They all more or less involved the assumption, that what is termed inflammation was their exciting cause, and the practical object of these terms was to connect local inflammatory processes with local and constitutional symptoms.

The subject of the pathology of puerperal fever is an extensive one, and I shall not attempt to treat it in the same manner as that of the etiology of the disease. There may be some excuse for doing this, as I feel more equal to deal with this part from personal experience and observation than with the other, which we have regarded as a matter of history. There is one point which would not be difficult to prove, if it were not self-apparent, and that is the fact, that neither morbid anatomy nor pathology assisted the theory of Gordon: I mean to say, that the idea of a specific poison would never have been evolved from such pathological work.

It was of very great value, however, in another way. Such theories as that which accounted for the swelling of the leg, or phlegmasia dolens; by supposed metastasis of lacteal secretion, very soon retired before the scalpel of the pathologist. The result of all this is, that we have some chance now of succeeding, after the separation of many pieces which would not fit together, in completing a puzzle of really no great complexity.

If it were asked why Gordon's theory did not find acceptance with pathological investigators (for we may observe that there were some of these who were distinguished for an extensive knowledge of the disease in all its forms, and had studied carefully all its symptoms), we must look for an answer in the supposition, that the pathological mind is of such a character as never to feel satisfied of the existence of anything without direct evidence, and then only when the method of demonstration employed is one with which it is familiar. The pathological mind is not a credulous one. It has a good deal of confidence in its own powers, and it likes to use its own visual organs. It would meet such a theory as Gordon's with an avowal of the greatest willingness to believe in the existence of a specific poison, provided only it could be shown the poison. To imagine that it could be contented with less is to do

injustice to its common sense and acuteness. For such a mind to conceive of an atom is almost impossible; it appears to it practically useless to do so, and to range beyond the sharply defined cordon of the demonstrable is scientifically reprehensible.

Let us examine to what pathology led. The principal effect was a very excellent classification of morbid changes, which has been followed closely by continental authors, and is still preserved by systematic writers in all countries. When it was found that descriptions of particular forms of inflammation were true for the particular cases in which they occurred, but not for all forms of the fever, we can understand that general views would soon embrace the apparently unconnected series of various morbid appearances which have been recorded. It soon appeared quite possible for various structures in the abdomen to be severally or simultaneously affected, or even that there might be well marked, though very different, pathological alterations, and yet that the cases were correctly described as forms of puerperal fever. Thus a simple classification was adopted under the general term uterine or puerperal inflammation, which included: 1. Inflammation of the peritoneal covering of the uterus and the peritoneal sac; 2. Inflammation of the uterine appendages; viz., the ovaria, Fallopian tubes, and broad ligaments; 3. Inflammation and softening of the proper or muscular tissue of the uterus; 4. Inflammation and suppuration of the absorbents and veins of the uterus. Each separate structure was thus carefully examined, and its alterations recorded.

There is a great deal to admire in the very exact and truthful descriptions, which make the works of the pathologists, to whom we are indebted for these valuable aids to our knowledge of puerperal fever, so well deserving of our regard. Probably the very fact, that a rigid exclusion of etiological influences in connection with these pathological conditions was observed by the school of pathologists to which I have referred, may be the cause of their work being so valuable. That publication of Dr. Baillie's, which contained nothing but careful and elaborate engravings, with the simplest descriptions possible, was the type of scholarly composition and masterly anatomical and pathological skill. There can be no doubt that, since the productions of that period from which we have traced the commencement of our history, nothing has ever appeared to compare with those of our own country. There were no uncertainty as to facts, no hasty generalisation, and no attempts at artificial classification, amongst those that followed close upon Baillie's model. So carefully did they attach themselves to the firm rock of certain truth, and so sure were they of the ground on which they stood, that they would have wondered at many since their day who have been looking so long into the hazy atmosphere that surrounds those difficult regions of the origin of life and disease, as to have forgotten to move on whilst dreamingly attempting to define the misty outlines of forms of things beyond their power of perception or comprehension. Such is the feeling they would have experienced, combined with one of pity, if they had heard some of us say that the very foundations of pathology are uncertain; that we know not quite where we are, or what next we may see. This brings me back to an urgent claim for the careful preservation amongst us of our peculiarly national method of observation, which, while accurate in detail and cautious in expression, still considers the best proof of its value to be the benefit it confers by its practical application.

To return to our subject, we have to notice the important extension of the theory of inflammation to the explanation of the connection between that condition which was known as phlegmasia dolens, and certain morbid changes in the structures of the venous system of the uterus; that is to say, when the symptoms which had previously been referred to metastatic lacteal secretion or pressure upon the lymphatics, were found to depend upon obstruction to the venous circulation.

The term phlebitis assumes the possibility of the veins, as separate structures, being affected by inflammation. Various changes of their lining membrane, and the existence of pus within the vessels, were considered satisfactory evidence of inflammatory action; and then followed the theory of pyæmia or puerperal purulent infection, arising from suppurative phlebitis. We will not stop to discuss just now whether these views were altogether correct. There is no doubt that, as far as the facts were concerned, they led to very important results.

It was shown that phlebitis might occur, not only in the puerperal condition, but also as a consequence of suppressed menstruation; in cases, too, of malignant ulceration of the os and cervix of the uterus; in certain forms of tuberculous disease; also as a consequence of abortion, and after operations for fibrous and other tumours of the uterus. The theory of puerperal purulent infection was supported by the popular school of medicine in France, and was ably argued by the late Professor Trousseau. It never was very generally received amongst ourselves, for the reason that we did not find it in accordance with clinical experience.

I can recommend the analysis of Trousseau's lectures on this subject to those who are interested in the comparison between French and the English character of mind, as influencing the scientific study of the phenomena of disease. Trousseau entertained the idea that puerperal fever depended on purulent infection, and that such infection was always to be traced to a wound; or that suppurative phlebitis might follow a wound, and pus might thus be introduced into the system. He assumed, or rather, fell back upon, an old theory, which has often been employed to explain many circumstances connected with the puerperal condition. As far as it bears upon our subject, which, I may remind you, is the specifically poisonous origin of the fever, we can fairly allow it to be passed over, at least at present. The theory to which I allude is this: that a certain diathesis characterises the puerperal state.

We can easily understand how the theory of a particular diathesis was also employed to explain that condition of the veins which we term thrombosis, or rather, that state of the blood which was supposed to be the primary cause of its coagulation; and a further application of somewhat the same theory to tuberculous and cancerous cases, which assumed in the latter instance the more familiar title of cachexia, was conveniently used to explain the occurrence of the various pathological conditions, and the symptoms arising from them which were found to be common to these diseases as well as to the puerperal state.

The last edition of Trousseau's lectures was published some years after Semmelweis's work, in which, as we shall see, it was clearly shown that cases of carcinoma of the uterus in the same ward with puerperal cases might produce that fever. This was undoubtedly, though a simple observation, one of considerable importance in bringing clinical evidence into consistent accordance with pathological facts. There is one remark of Trousseau's, however, which clearly shews that he was quite aware of the weakness of his argument in favour of diathetic influence. "The same conditions", he says, "exist in nearly every recently delivered woman, or in all cachectic persons", yet "phlegmasia dolens is not a necessary complication in either class of patients"; from which he concludes very candidly that "consequently there exists a special cause which is unknown". It happened about this time that the doctrines of the German pathologists were exciting a good deal of attention in their bearing on the theory of inflammation. Those important changes which are observed in the various forms of puerperal inflammation were subjected to the theory which Virchow expounded, of the origin of inflammation in certain alterations in connective tissue. It is not essential to our argument to consider whether that theory be correct or not; for it is, as far as we can judge at present, not of very great practical importance whether the process of inflammation commence with active changes in the cellular elements of connective tissue, or in the exudation of the corpuscular elements of circulating fluids. They both agree in allowing the existence of some primary cause for the tissue-changes; they only differ in respect to the earliest effects which that cause produces. It is now generally admitted that the connective tissue is that which is chiefly affected in puerperal inflammation in all its various forms; that is to say, in metritis, phlebitis, and others; and the term cellulitis is more commonly used than any other, at least by those who receive the recent doctrines on inflammation.

Clinical experience and pathological examination of this question lead me to concur entirely in these views. Instead of regarding a vein as a separate structure, and thus admitting the term phlebitis to be pathologically correct, it appears that we must refer the coagulation of the blood, which is the consequence of so-called phlebitis, to changes in the connective tissue which surrounds the vessels. On this point, therefore, we are obliged to differ from the pathologists whom Trousseau followed, as well as from those who consider that the tissue-changes are induced by coagulation of the blood or thrombosis, whatever that may depend upon.

There is not much difficulty in comprehending the process by which a vein becomes filled with coagulum in phlegmasia dolens, whether it occur in the puerperal state, or under any other circumstances. Perhaps the best instance from which we can inform ourselves of the relation which the process of coagulation bears to that of cellulitis, is one of those rapidly fatal cases of the disease in which the large vessels which unite to form the vena cava ascendens are filled with coagula. By removing the main trunks contained in the pelvis, together with the adjacent arteries, as high as the diaphragm, we can determine by dissection the exact points where the coagula terminate. If we then examine the condition of the cellular tissue surrounding the vessel under inspection, it will be found that the external changes are in advance of the coagulum, assuming that both processes are taking place in the direction of the venous circulation. As a matter of demonstration, we may insert a needle through the vein at the point where the inflammatory changes are seen to terminate; and, on comparing its position with that of the termination of the coagulum, we may measure the distance be-

tween them. This may amount to an inch or more; but, to a great extent, it appears to depend on the number of the lateral veins that enter the vessel between those two points. The orifices of the lateral veins are generally distinguished by a small projecting coagulum, which is much less firm than that of the principal coagulum. The space between the two points is not empty, but the blood which it contains is in the same condition as that which fills the vena cava itself, and generally distends the right cardiac chambers. It is satisfactory to find that this view of the pathology of phlebitis, using the word under correction, is admitted by the leading pathologists in Germany, and that they are inclined to adopt it in preference to the theory of thrombosis or primary coagulation.

We may dismiss this part of our subject with one or two brief remarks. It may be asked, How are we to explain the formation of pus in veins on the supposition that the connective tissue externally to them is primarily affected? We should reply to this: that there is evidence to prove that the different tissues of which the venous canals are composed participate very soon in the changes of the connective tissue; that the termination of the inflammatory process in any locality is the formation of pus, and that an abscess may form in such a position and in such a manner as to allow of a communication between the collection of pus and the venous canal. If, again, the question be put to us, Can the lining membrane of a vein secrete pus in the same way that the pleural membranes, the pericardium, or peritoneum are known to do? we can only answer that the epithelial surface of a vein may exhibit the ordinary changes which succeed one another in the stages of inflammation, and that in the final stage pus-corpuscles may be secreted, or rather developed, but only when the epithelial surface has been destroyed. The practical value of these pathological researches has been to afford us very valuable assistance, not in the elucidation of the etiology of puerperal fever, but in the explanation of several symptoms which were previously unaccountable, such as those resulting from embolism, coagulation of blood in the right side of the heart, the formation of secondary abscesses in various organs and tissues, and the constitutional disturbance which is produced by them.

I shall now request your attention to those contributions which were made in Germany to the etiology of puerperal fever by Semmelweis. Quite independently of his relative personal merits as a scientific observer, we may simply confine ourselves at first to the evidence which he offers on the subject of the contagious nature of the fever. After perusing the first few pages of his work, we find that he presents himself to us as an independent witness. There is also the manner which favourably impresses us with the idea, that he would not assert anything which he could not prove; on the contrary, we feel that he rather tires us with details. He does not mix up conjecture or theory with fact, though he may indulge too much in arguments which are intended to prove certain conclusions almost self-evident. The method of investigation pursued by Semmelweis was much the same as that followed by Gordon, and the proposition he supported was, like his, that puerperal fever is a disease (essentially) produced by the introduction of a septic poison into the system. Gordon, as we have seen, declined to express any opinion on what the poison might be; but Semmelweis did not hesitate to state that it was generated in the foul organised substance which results from the decomposition of animal matter. The work does not contain any original contributions to the pathology of the disease, and in that respect, as an observer of pathological conditions, he appears to me to have been decidedly inferior to Gordon.

Semmelweis argues his point somewhat in the following manner. "We see that the great lying-in institutions are singularly subject to occasional outbreaks of this disease, and the danger to puerperal women is very much greater in those institutions than in separate domiciles. In those hospitals where students are instructed, and are in the habit of coming to the wards, and to the examination of cases from the dissecting-room, the disease is more frequent and the mortality greater than when only midwives are admitted for instruction. To prove this, we have such and such a tabulated register of cases, which shows a mortality under one system of 10 per cent., as compared with $3\frac{1}{2}$ in the other. In Vienna, before 1841, the mortality in the lying-in hospital was small, when compared with that of a few years later, when the hospital had been converted from its first intention to serve for the clinical instruction of students."

"To prove that the poison was generated in the process of decomposition, and might be conveyed by the hands of attendants, a case of cancer of the uterus in a pregnant woman was the cause of a dozen fatal cases in the same ward, and the probable means of conveyance of the poison were the sponges and other things employed by the nurses, or in the course of examination by the medical attendants. We had a case, too, of necrosis of a knee-joint, which was also the cause of some fatal cases. These were clearly established facts, and they point to the

same conclusion." This is the nature of the argument, which is fairly well worked out. To show that the dissecting-room might be the source of danger, he makes the declaration:—"I myself was appointed for the second time to the clinical assistantship of the first *clinique*, and was in the habit of going into the wards after making explorations in the dead-room. The mortality in April and May was 18 per cent. and 12 per cent. respectively. During the four preceding months, I was only occasional assistant, and was not engaged in *post mortem* inquiries, nor was I obliged to examine the patients *seriatim* before the visits of the Professor. Look at the difference between the mortality in the two cases."

Then he goes on to show us from his tables that there is no connection between seasons of the year and the occurrence of the fever; so that it could not be said in any way to depend on epidemic conditions of atmosphere or temperature. There is a great deal more in the way of collateral evidence adduced by Semmelweis, but the main line of argument he pursued is thus broadly indicated. It was essentially based on statistics fairly arranged, without taking advantage of such data, and without overstraining them.

He did not neglect an important test of his views by ascertaining the effect on living animals of the artificial introduction of healthy pus into the circulation, thus proving that the symptoms could not be referred to such a cause as simple purulent infection.

I have said that Semmelweis's work consists chiefly of a collection of observations made by himself, without direct reference to the opinions of others. He does not omit, however, to mention the views of some few whom he was led to regard as the representatives of scientific thought, both in his own country and abroad. In this part of his work, however, it must be acknowledged that he gives proofs of possessing very limited knowledge indeed of the literature of his subject, and we cannot feel surprised that he finds but little difficulty in disposing of his opponents to his own satisfaction.

He brings his arguments together into a condensed conclusion at the end of his treatise in brief and forcible language. "I have shown", he says, "above that the excess of mortality in the first lying-in *clinique* in comparison with that in the second was occasioned by putrid matter, which was introduced by the hand of the examiner. I have shown that, in October 1847, the foul material of a foul cancer of the uterus produced puerperal fever. I have shown that, in November 1847, a decomposing material from a carious knee-joint produced puerperal fever; how the fever was connected with the visits of the surgeons in one hospital in particular, and how the mortality during six years was reduced to less than one per cent. when all connection with the surgical wards was broken off. It was the finger of the surgeon that carried the poison in the latter case; it was the atmosphere in the case of the knee-joint, and the air and the foul linen together in the *clinique* at Pesth, where the infection spread in spite of the washing with chlorine-water."

It may be mentioned that he had adopted the use of chlorine-water with signally good results, and derived from this fact increased support to his views of the septic origin of the disease. We will not venture to make any further comparison between the value of Semmelweis's treatise and that of Gordon. As far as the merit of priority is concerned, though the observations and opinions of the one may have been quite independent of the other, we can certainly claim for Gordon precedence in date by a considerable period.

In their style of composition, in their arrangement of data, and their logical powers, it is more than probable that we should be inclined to claim for Gordon decided superiority over Semmelweis. At the commencement of these lectures, I briefly alluded to the condition of mind which it is necessary for us to obtain in order to weigh fairly scientific evidence. It is easy to decide on the value of such evidence when we have tested it on the anvil of time. Observations and theories which have been conclusively analysed by that process naturally lose their interest. But, when we are studying the pages of history, it is the lives and characters of those who thought, and worked, and benefited us, that we feel an interest in contemplating, as well also as the lives and characters of those who did not think and did not work, and stood in the way to oppose the progress of scientific and practical knowledge. The history of this fever, if it contained nothing but facts, with the dates of their discovery, would not be a long one. The history of the erroneous ideas and statements which were advanced on the subject of its origin and nature would far surpass it in dimensions. It has been said by one who has displayed singular scientific acumen in philological research that, "for the discovery of truth, there is nothing so useful as the study of errors". It may be useful certainly to the individual who pretends to originality; but, for our purpose, I venture to think we have done well in confining our attention to the more limited object with which we started, of tracing the progress of truth through its various stages of development.

CLINICAL LECTURE ON CLUB-FOOT AND ITS TREATMENT.

By GEORGE BUCHANAN, A.M., M.D.,

Professor of Clinical Surgery in the University of Glasgow, Surgeon to the Western Infirmary, etc.

THERE are few operations in surgery which the surgeon approaches with more satisfaction than those for the cure of natural deformities. Amputation of limbs, resection of joints, and excision of tumours, are necessary and important as a means of saving or prolonging life after other treatment has failed to remove or arrest disease; but these are evidences of the imperfection of our knowledge of the nature and origin of many affections, and, at all events of the inefficiency of our therapeutic applications. Let us hope that the advance of pathological and therapeutical knowledge will one day diminish the number of radical operations, which at present are necessary in the treatment of surgical diseases.

But we cannot expect that any extension of our researches will diminish the number of children born with congenital deformities. Into the popularly supposed causes of these defects, it is not our place to enter; it is sufficient that they are common enough to demand a careful consideration on our part. It is very probable that many of you who are destined to become general practitioners, may never be called on to perform operations for the cure of deformities, delegating those duties to those who are to take the position of operating surgeons; but some of you will, I have no doubt, have as your sphere some place where you must rely on your own resources for every medical service demanded by your patients; and all of you will have it in your power to advise and direct patients as to the proper course to be followed. This is a most important position, for it is in early years that most of these defects can be most readily remedied. Never was this more impressed on my mind than by the case now before you. Here is a young gentleman, eighteen years of age, who has been the victim of talipes varus since infancy. The deformity seemed so hopeless to the medical attendant, that he did not venture to suggest any treatment; and the various surgeons, who were from time to time consulted, dissuaded the parents from any operative interference. Six months ago, he came to me, and, though the deformity was extreme and all the structures implicated were rendered rigid by the length of time they had been in their unnatural position, I did not despair of being able to improve his state, at least in a material degree. The result has exceeded our most sanguine expectations, and the patient is here to walk across the theatre on the sole of his foot, a thing he never did in his life till a month ago.

The further illustration of this subject will be found in connection with the case I shall now introduce, and also another in the wards, which will be operated on a short time hence. Here is a boy, aged seven years, with a well-marked example of talipes varus, or inverted club-foot. Observe the particulars of the distortion. First, the heel is raised and twisted towards the opposite foot. Second, the inside of the foot is twisted round so that the sole of the foot looks upwards, and the boy walks on the dorsum of the tarsus turned down. The sole of the foot is also curled inwards, so that the ball of the great toe is much closer to the heel than it ought to be. The whole foot is so distorted, that the only kind of boot which the boy can wear for the protection of the skin is a sort of leather bag with a strong thick sole. Let us now consider the structures concerned in this deformity. Let me remind you, then, club-foot exists in four varieties—talipes equinus, talipes calcaneus, talipes valgus, and talipes varus. Each of these varieties may depend on atony of one set of muscles, or an undue contraction of the antagonistic set. Hence we have tonic and atonic talipes. Now it is important to recognise this difference, because, though in both you may, by mechanical or operative means, restore the normal shape of the foot, yet it is in the tonic variety alone that you can promise a really favourable result as regards the future power of walking. I am far from taking a hopeless view of cases depending on muscular atony, or paralysis as it is often called, for I have had under my care a great many children in whom by persevering mechanical, gymnastic, and operative treatment I had had every reason to be satisfied with the result. The whole subject of the proper treatment of muscular atony, whether the result of deformity or prolonged retention in one position, is well deserving of notice, and I shall call your attention to it in another lecture.

The case before us is one of the tonic variety, in which the distortion is produced, or at least kept up, by the contraction or rigidity of the muscles, tendons, ligaments, and fasciæ. With regard to the