I have seen cases in which, on examination of the tumour after removal, I discovered the epithelium of the cysts in a thickened and villous condition. If the cysts of such a tumour have remained whole, and there has been no escape of epithelial elements from it into the peritoneum, then there need be little fear of malignant peritonitis arising in that patient in the manner we have described.

We make it a rule now to examine very carefully the sediment obtained from ascitic fluids, never mind what the nature of the tumour may be. Malignant ovarian tumours are in most cases surrounded by ascitic fluid; and if, in ascitic fluids, such little masses of proliferating epithelium as I have described are found, then in these cases we may, I hold, with certainty diagnose malignant peritonitis.

Dr. Keith long ago arrived at the conclusion that a burst ovarian cystic tumour should be removed as quickly as possible; and, as the result of pathological inquiry, it would seem that such procedure is to be strongly recommended.

INTRATHORACIC TUMOURS.*

By J. W. F. SMITH-SHAND, M.D.,

Physician to the Aberdeen Royal Infirmary, etc.

I.—PRIMARY CANCER OF THE LUNGS.

I GIVE the following cases as a contribution to the history and diagnosis of intrathoracic tumours. Notwithstanding all that has been written about them, they still too frequently baffle the diagnostic skill of the physician, and often the exact nature of the malady is only first revealed at the *post mortem* examination. But, as observations of these cases are multiplied, the difficulties connected with them will disappear. As Dr. Graves, in his Clinical Lectures, observes about a case of cancer of the lungs, which during life was a mystery and puzzle to himself and Dr. Stokes, "rare cases should not be looked upon as mere matters of curiosity, but should be attentively studied, with the view of enabling us to recognise the true nature of similar cases when they again occur. Were the history of diseases at present reputed to be extremely uncommon published by all those who meet with them, facts now apparently single and insulated would serve as nuclei round which future experience and observation might cluster together similar facts in groups sufficiently numerous to illustrate and explain each other." It is with this object that I record the following cases; and I take this opportunity of expressing the great pleasure and instruction I derived from the perusal of Dr. Risdon Bennett's Lumleian lectures on the Natural History and Diagnosis of Intrathoracic Cancer, as published in the *BRITISH MEDICAL JOURNAL* for 1870.

CASE I.—The first case that I shall lay before you is that of Anne Mackie, domestic servant, aged 36, who was admitted into the Aberdeen Royal Infirmary on May 18th, 1869, complaining of cough and loss of power in the right leg.

Previous History.—The patient stated that, having previously enjoyed good health, she had for the last six months been troubled with a cough, with slight expectoration, which had on two or three occasions contained small streaks of blood. Any pain that she occasionally felt in her chest was easily removed by the application of a mustard poultice. About five weeks previously to admission, on coming downstairs one morning, she felt her right foot slip out behind her, but she did not fall; and since then the loss of power in the leg had increased. She was treated with cough mixtures, and a liniment for the supposed sprain, but without benefit.

Condition on Admission.—The patient was of spare habit of body, with dark hair and muddy complexion, and an anxious expression of countenance. When walking, she dragged her right foot behind her, with the toes touching the ground, and, on examining the leg, it was found that she had no power of moving in any way the toes or ankle. Sensation was not much affected, except to a slight degree over the dorsum of the foot. Her voice was weak and husky, and had altered in character since her illness commenced; but she said that she had not lately had any sore-throat. On examining the chest, the superficial veins of the left side were seen to be enlarged. The left thorax looked fuller and more rounded than the right, with entire absence of movements of expansion or elevation on that side. Percussion, dull all over the left side, was normal on the right. On auscultation, there was entire absence of respiratory murmur over the left lung, except behind, on a level with the spine of the scapula, where tubular breathing was heard over a very limited space. Vocal resonance was diminished, and vocal fremitus was absent on the left side. The breath-sounds over the right lung were puerile. The patient had a rather frequent dry

hollow cough, with scanty frothy expectoration. She complained of no pain in the chest, and could lie on either side. There seemed a general impulse or heave over the cardiac region, but no apex-beat could be felt. A soft systolic murmur was heard in the region of the aorta, loudest behind and over the first bone of the sternum in front. Pulse 104, small. No palpitation was complained of. The tongue was furred; the appetite fair; but there was dysphagia. Bowels regular. The urine was acid, of specific gravity 1030, with a slight trace of albumen. Menstruation was regular, but the last period was missed; and she had been troubled with headaches since that time.

Treatment and Progress of Case.—May 19th. She was ordered five grains of iodide of potassium, in a mixture, three times a day.—May 25th. The patient reported herself much the same; but, on careful examination, it was found that the symptoms of hemiplegia had increased, as the tongue was protruded to the right side, the right orbicularis palpebrarum was affected, and there was loss of power in the right hand. The pupils were movable under the influence of light, but the left pupil was more contracted than the right.—June 3rd. Pain was to-day complained of in the right side, and the existence of pleuritis was ascertained by the usual physical signs. Under this fresh attack of illness, she rapidly sank, and died on June 6th.

Post Mortem Examination.—I am indebted in great part to Dr. J. M. Bruce, house-physician at that time, and now of the Charing Cross Hospital, London, for the following observations. A large soft cancerous tumour was found on the left side of the chest, consisting of the lung of that side. It was adherent to the pericardium, completely surrounded the great vessels at the root of the neck, and compressed the left vagus and recurrent laryngeal nerves. The lung, weighing 78 ounces, was perfectly full of soft cancer, and the bronchus was plugged with an encephaloid mass growing into it. In the right pleura, there were found recent pleuritic adhesions and 20 ounces of effusion. The right lung weighed 22 ounces. The heart was normal; there was no valvular disease. There was slight atheroma in the aorta. The liver weighed 53 ounces, and was fatty. Both kidneys were fatty; the right weighed 5½ ounces, the left 6 ounces. The spleen weighed 6 ounces, and was healthy. The brain weighed 43 ounces, and contained deposits of soft cancer in several places: 1. On the inner surface of the left process of a cerebello ad testes; 2. On the left pes hippocampi; 3. On the right hippocampus minor; 4. On the right Sylvian fissure. The optic thalami and corpora striata were healthy. A few more spots of soft cancer were scattered through the cerebrum, especially in the posterior lobes, one side not seeming to be more diseased than the other.

REMARKS.—This is a good example of that rare disease, primary cancer of the lung. As a secondary affection, it is more common, and, in that case, a diagnosis may often be formed with absolute certainty; but the primary form, as Dr. Biegel says in the article on Cancer of the Lung in Reynolds's *System of Medicine*, in the majority of cases, admits of no diagnosis.

On examining the patient, it was evident to me that I had not to deal with an ordinary case of chest-disease. The full and more rounded appearance of the left thorax, together with the enlargement of the superficial veins and the absolute dullness on percussion, led me to conclude that there was pleuritic effusion, and also intrathoracic tumour. At the same time, the presence of paralysis, the history of sore-throat, the husky voice, and the general appearance of the patient, made me ascribe a syphilitic origin to these symptoms, and accordingly I treated her with the iodide of potassium. Impressed with this idea, the probability of carcinoma, I must confess, did not enter my mind.

There are many points of clinical interest about this case. The heaving impulse over the cardiac region, and the aortic murmur (which was probably caused by pressure, as there was no valvular disease), and the contracted pupil, might have given rise to a suspicion of aneurismal complication. It is difficult to say why, with such a large mass occupying the left thorax, the heart did not appear displaced to the right side. It may have been on account of the intimate adhesions between the pericardium and the cancerous mass which surrounded it. But, if it had been, it would have been almost impossible to doubt the existence of a large pleuritic effusion. Dr. Risdon Bennett, in his Lumleian lectures, mentions a case which occurred in the practice of his colleague Dr. Sutton, where there was great dyspnoea and lividity of countenance, with the left side of the chest bulged, absolutely dull before and behind, respiration inaudible and heart displaced. The patient, aged 11, was repeatedly examined by Dr. Herbert Davies and others, and all decided that the chest was full of fluid. Thoracentesis was twice performed, but nothing issued, except a little dark blood. The patient died some time afterwards, and the left side was found to be entirely occupied by a mass of medullary cancer, the left lung being collapsed and pushed backwards. The dysphagia complained of by Mackie was doubtless the result of the compression of the vagus nerve; but it was

* Read before the Aberdeen, Banff, and Kincardine Branch.

singular that there was no dyspnoea or distress until the pleuritic attack took place on the right side, when we consider the pressure, if not destruction, to which the recurrent laryngeal nerve was exposed. Dr. Risdon Bennett expresses his belief that, if there be any considerable degree of pressure or irritation of the recurrent laryngeal nerves, there will invariably be a great amount of both dyspnoea and distress: at the same time, he observes that it is somewhat remarkable that symptomatic vomiting or other disturbance of the stomach is not more frequently met with when there is destruction of continuity of one pneumogastric nerve. The experience of this case, then, shows that, even when the recurrent laryngeal nerve is involved, as indicated by the changed and husky voice, dyspnoea and distress may not necessarily be produced.

The normal temperature observed corresponds with the experience of Wunderlich, who says that it is a peculiarity of cancer cases that elevated temperatures are comparatively rare. The absence of pain also corresponds with what has been observed in the majority of cases of cancer of the lung.

I regret I was not at the time acquainted with the valuable sign of tumour in the neck first pointed out by the late Dr. Kilgour of this city, with whose acute powers of observation and diagnostic skill many of us are well acquainted. I cannot, therefore, say whether the glands were affected or not; but I think, if they had been to any extent, it would have been noticed.

Dr. Cockle, in his able and learned work on *Intrathoracic Cancers*, lays great stress upon the importance of external tumour; but one of his cases, in which the diagnosis was difficult, illustrates also the necessity of ascertaining the *duration* of the tumour when present, as the patient in question had had three sebaceous tumours on his neck for many years, while other tumours, cancerous in nature, appeared on the abdomen only shortly before death. Primary cancer of the lungs is said by Walshe to give rise very seldom to secondary deposits elsewhere, although the lungs themselves are frequently affected as a sequel of disease in other organs. The case of Mackie is, therefore, interesting, on account of the secondary cerebral complication connected with it.

(To be continued.)

REMOVAL OF THIRD METATARSAL BONE : ANÆSTHESIA BY CHLORAL-HYDRATE.

By G. J. S. NAIRNE, Surgeon, and ALEX. NAIRNE, Physician,
Glasgow.

ON April 1st, 1875, Christina D., aged 3½, a delicate looking girl, was put under the influence of chloral, and had the third metatarsal bone of her left foot removed for disease. The operation had nothing special to note, but the narration of the administration of the chloral may be interesting.

An aqueous solution was prepared, of the strength of twelve grains to twenty minims. A solution of strychnine (two grains to the ounce), was also prepared. At 10.30 A.M., ten minims of the chloral solution were injected subcutaneously into the right lower leg. At 10.38, there was no perceptible effect; the child cried continuously. At 10.40, ten minims more were injected subcutaneously into the lower part of the left leg, and the child was given to its mother to carry about. At 10.45, the patient was drowsy, but easily roused; she started and cried whenever the mother would lay her down. At 10.55, ten minims more were injected; at 11, she was sleeping apparently soundly. There was no flushed face, but the pulse was rather quick. She awoke and cried. At 11.5, ten minims were administered by the mouth. She dropped off to sleep in a few seconds, and began to snore. Pinching made her cry, but she fell asleep immediately. At 11.18, ten minims more were administered by the mouth. At 11.20, she was in a deep sleep.

The operation was now performed. During its performance she began to cry, but made no struggling to speak of; the eyes remained nearly shut, and the crying seemed that of a dreaming child. Immediately she fell asleep again as sound as ever. Strong tea was prepared, of which she was made to drink a cupful. Her pulse was quick but firm, so we let her alone. She slept deeply for over two hours. More strong tea and milk were then given her, and, in another three-quarters of an hour, she sat up dazed, looking quite unterrified, and apparently oblivious to any pain. Next morning, she took her porridge and milk with quite a good appetite.

Altogether, the administration of chloral seemed better than that of chloroform—in this instance, at least. There was no convulsion, nor apoplectic appearance, and no unnatural disturbance of the breathing. There was no subsequent sickness nor any apparent ill effects. We

conceive that chloral would be much safer and better than chloroforming in ophthalmic operations.

We were glad we had no cause to use the strychnine, but would not have hesitated to inject it subcutaneously if there had appeared any necessity. We have administered strychnine frequently after chloroforming, and generally found an immediate good effect in brightening up the patient and preventing sickness. Subcutaneous injection would, therefore, very likely be successful in restoring animation in cases of chloroform poisoning.

Next time we use chloral we intend to have a much more dilute solution, and a larger syringe, and to inject it into a vein. By these means, we hope to avoid the pain caused by the injection, provide as far as possible against phlebitis, prevent the formation of painful swellings or ecchymoses, and secure the more immediate introduction of the fluid into the circulation. Total unconsciousness does not seem at all necessary to produce anæsthesia, as, although the girl woke up and cried during the operation, she made no struggling, and was asleep before her cry was finished.

CLINICAL MEMORANDA.

RAPID PULSE.

I HAVE read an article in the JOURNAL of this date (June 12th) by Dr. Robert Farquharson of St. Mary's Hospital, on Rapid Cardiac Pulse, and I feel particular interest in the cases which he has recorded, having at the present time a patient under my care whose pulse at the wrist it was difficult to count. I happened to be called to this gentleman, a foreigner, aged 65 to 70, fond of alcoholic drinks, and of bilious temperament. I found him at two o'clock in the morning scarcely able to breathe, and sitting on the side of his bed, surrounded by his relatives, who were in the greatest alarm. The pulse at the wrist was irregular, and 140. There was the same rapid irregular "to-and-fro" action of the heart, apparently caused by aortic insufficiency; but, as in Dr. Farquharson's case, no *bruit* could be detected. As it turned out afterwards, it was the beginning of an attack of bronchitis, and, when free expectoration took place, there was a marked improvement in the circulation, the pulse coming down to 130, then to 120, where it has remained. The urine was scanty, high coloured, acid, and of specific gravity 1010. No albumen could be detected by the heat and nitric acid test. The patient is still under treatment, and relieved as regards the bronchial attack, though still suffering from what would be considered a very rapid pulse (120 to 130), yet not coming up to the abnormal rapid pulse recorded by Dr. Farquharson.

J. D. CRONIN, M.D., F.R.C.S.E., Fleet-Surgeon R.N. Retired,
Queenstown, Ireland.

NOTES ON A CASE OF INSANITY INDIRECTLY CAUSED BY PHIMOSIS.

A. J. K., aged 26, had been formerly a schoolmaster, but now devoted himself exclusively to the study of music, practising on the piano or organ eight hours daily. On inquiry, it was found that he was obliged to give up the school, as he found he could not exercise sufficient authority or control over the boys. There was a slight loss of memory, and he had some difficulty in remembering his age. He stated that the symptoms first noticed were pains in the back, but was unable to say when they commenced. He had lately suffered from a sensation in the occipital region as if the skull were being depressed. He was a seven months' child, and did not walk until he was three years old. When an infant, he suffered from constant diarrhoea and prolapsus ani. He was always very excitable, but his excessive mental and bodily weakness prevented his following any regular occupation. He is passionately fond of music, and has overtaken his strength by his studies. He had for the last six months been under the delusion that certain people had been following him about and making indecent gestures at him. He could not speak of these facts without bursting into tears and becoming hysterical. He believed that these people had done their best to prevent his getting a living. On inquiry, it was found that he was addicted to masturbation. This habit he attributed entirely to a state of irritation and itching which were constantly present in the penis. On examination, it was found that there was a condition of congenital phimosis, the secretion from the glandula Tysonii never having been cleared away. The case was treated temporarily with bromide of potassium and aperients, and partial circumcision was recommended. The operation was successfully performed; and, from a letter subsequently received, it appears that the habit has