The head and shoulders were not drawn over the table, but the pillow was removed, and artificial respiration, drawing out of the tongue, and cold water dashed on the face, were tried without effect.]

ON CERTAIN PHYSICAL PHENOMENA CONNECTED WITH THE CIRCULATION, RESPIRATION, SECRETION, AND NUTRITION.*

By GEORGE JOHNSON, M.D., F.R.S.,

Professor of Medicine in King's College; Physician to King's College Hospital.

I WISH now to direct attention to some of the physical results of an impeded return of blood through the systemic veins. In order that the blood may maintain its normal composition and its physical properties, it must move freely through the vessels. Dr. Polli (*op. cit.*, p. 306), by a series of experiments, clearly proved that the blood is rendered more dense by inducing an artificial stagnation or engorgement by means of the fillet applied for some time previously to opening the vein; and his conclusion is, that "venous blood, when it is prevented by any obstacle from circulating freely in its veins, becomes more dense, or loses a certain portion of its watery principle". And he asks: "Does not this fact constitute a principal pivot of the doctrine which explains œdemas, cellular dropsies, and other infiltrations, by considering them as exudations induced by obstructed circulation?" Without doubt, the œdematous swelling of a limb, whose principal veins are obstructed, is the counterpart of the blood-thickening which results from the transudation of the watery constituents. The force which drives the water through the walls of the vessels is derived from the left ventricle of the heart. The blood is driven through the capillaries into the veins, and its passage through the venous channels being impeded, a portion of its water is forced through the capillary walls into the tissues outside. The result is an œdematous dropsical swelling of the limb, and an increased thickness and consistency of the blood within the veins. This simple experiment explains some forms of blood-thickening which have perplexed pathologists.

It explains the blood-thickening of apnoea. It is a well known fact that a great impediment to the aëration of the blood is always associated as one of its consequences with an impeded circulation through the lungs, and a consequent engorgement of the right side of the heart and of the whole systemic venous system. One result of this impeded venous circulation is a puffy œdematous condition of the subcutaneous tissue, especially of the face and neck; and another result is, that the blood becomes dark, thick, and treacly, and coagula often form in the right cavities of the heart and in the pulmonary artery.

Dr. George Buchanan, in one of his papers on tracheotomy in croup and diphtheria, read before the Medico-Chirurgical Society of Glasgow, in giving the history of a case in which he opened the wind-pipe in an advanced stage of laryngeal diphtheria, states that, "as soon as the wound was made, a quantity of dark tarry-looking blood flowed from the edges, and was with difficulty cleared away by firm and repeated applications of a moist sponge". Dr. Buchanan, again referring to the state of the blood, says "it was quite evident that that fluid was in a dark and viscid state. The darkness might be accounted for (he says) by the impeded respiration; but the tarry consistency struck me as a condition which might be one of the pathological elements of the disease (diphtheria) in an advanced stage." The tarry consistency of the blood is, in fact, one of the usual results of extreme and prolonged apnoea. Not long since, during the operation for tracheotomy on account of laryngeal obstruction of long standing, I saw a decolorised fibrinous coagulum coughed out of one of the cut veins in the neck. When I say it was coughed out, I mean that it was thrust out of the vein during the expiratory effort of coughing, which, as we continually see, always forcibly drives the blood into the superficial cervical veins.

There is also chemical evidence of the blood-thickening which results from apnoeal conditions. Thus the late Dr. Dundas Thomson, who published an elaborate paper on the Chemistry of the Blood in Cholera (*Medico-Chirurgical Transactions*, vol. xxxiii), states that one specimen of blood "from a patient who laboured under an affection of the mucous membrane of the air-tubes contained a greater excess of solids in proportion to water than he had found even in cholera blood".

I now come to one of the main objects of this communication, which is to suggest a more complete explanation than has hitherto been given of the increased density and viscosity of the blood during the collapse

stage of cholera. In order to explain the phenomena, we must have the facts clearly set forth.

The main facts, then, are these. The blood-thickening is a phenomenon which is present only during the stage of collapse; it does not occur during the diarrhoeal stage, and it quickly passes away when reaction occurs. We have already seen that mere diarrhoea, however profuse, does not thicken the blood; the reason being that, so long as the circulation of the blood is unimpeded, water passes from the tissues into the blood as rapidly as it escapes from the blood into the bowels. Dr. Garrod, many years ago, made careful analyses of the blood in cases of cholera, and he states that "the amount of intestinal evacuations in any case is by no means an indication of the extent to which the blood has become altered". (*London Journal of Medicine*, vol. i, p. 409.)

The blood-thickening, as a rule, bears a direct relation to the degree and the duration of the collapse. Now, there is one condition which is common to cases of apnoea and of choleraic collapse; and that is an impeded flow of blood through the lungs. This impeded circulation, with the consequent engorgement of the systemic venous system, is the main cause of the blood-thickening, which is also common to the two pathological states. The blood is thickened partly in consequence of its slow movement and partial stasis within the veins; but mainly, perhaps, by the passage of its watery portion through the walls of the capillaries. In the choleraic cases, it is probable that the passage of water from the distended venous system into the tissues is facilitated and much increased by the previous partial dehydration of the tissues during the diarrhoeal stage. The less water the tissues contain, the more readily would water pass into them through the walls of one of the distended capillaries.

There is another condition, common to cases of apnoea and choleraic collapse, which must tend to lessen the stream of water which naturally sets from the tissues into the blood. The impeded circulation through the lungs during the stage of collapse involves, as a necessary consequence, a diminished supply of oxygen, and, therefore, a diminished oxidation of the tissues. Many of the products of oxidation are crystalloids; for example, urea, uric acid, creatin, and the bile acids. These crystalloids, when formed in the tissues, pass into the blood, and with them they carry, or are carried by, a current of water. The crystalloid products of oxidation may be said normally to have an interstitial diuretic influence similar to that which urea and other urinary crystalloids exert upon the kidneys. During the collapse stage of cholera, the main products of oxidation—carbonic acid, urinary and biliary constituents—are simultaneously and greatly diminished. We see, then, that, at the same time that the distension of the systemic veins tends to force water from the blood into the tissues, the formation of crystalloids is much lessened; and, in a corresponding degree, the flux of liquid which, under their influence, normally passes from the tissues into the blood, is impeded. It is obvious that, during the stage of collapse, with the physical conditions already described favouring the passage of water from blood to tissues, and retarding the flow from tissues to blood, a continued escape of water through the alimentary canal would have much more influence on the composition of the blood than an equal drain of liquid at a time when the circulation of the blood, and the chemical and physiological changes resulting therefrom, are unimpeded.

I shall, perhaps, render my doctrine more intelligible if I recapitulate now the main phenomena of cholera upon which the blood-changes depend. In the diarrhoea stage, the discharge of water from the bowels draws an equal amount of water from the tissues, but does not thicken the blood. In the stage of collapse, the impeded circulation through the lungs involves, as a necessary consequence, venous distension, a limited supply of oxygen to the blood and the tissues, and hence a lessened formation of crystalloids in the tissues. It consequently happens that two influences tend simultaneously to deprive the blood of water: (1) the arrest or diminution of the current of water-borne crystalloids from the tissues; and (2) the passage of water from the distended veins into the more or less dehydrated tissues. If, at the same time, there be a continuance of the intestinal discharges, the blood-thickening is yet further increased.

Before I conclude, I wish to direct attention to the general conditions under which the circulation of the blood is arrested or much retarded by the influence of the minute blood-vessels. The phenomena to which I refer may be observed especially in cases of apnoea, cholera-collapse, and chronic renal disease.

The first effect of paralysing the respiratory muscles by curara is to increase the blood-pressure in the systemic arteries. Then, after a time, the pressure falls; and, after death, a large accumulation of blood is found in the right cavities of the heart, while the left are comparatively empty. The explanation of the phenomena appears to be

* Concluded from page 8 of last number.

this. The passage of unacrated blood into the systemic vessels excites, through the vaso-motor nerves and centre, contraction of the muscular arterioles. This explains the increased arterial tension. The blood which returns from the systemic capillaries through the veins must be of abnormal quality. The oxidation of the blood and tissues is suspended; and, therefore, the products of oxidation usually present in the venous blood are absent. This abnormal blood reaching the lungs, then excites contraction of the pulmonary arterioles, and the circulation through the lungs is thus impeded. The diminished pressure in the systemic arteries in the advanced stage of apnoea is explained partly by the diminished flow of blood through the lungs, and the consequent depletion of the systemic arteries; partly, perhaps, by weakening of the muscular walls of the heart, consequent on the defective blood-supply to the coronary arteries, the blood being at the same time unoxegenised.

After death in the collapse stage of cholera, the state of the lungs and of the heart's cavities is essentially the same as in cases of death from acute apnoea. The main phenomena of choleraic collapse are explained by the impeded pulmonary circulation.

The condition of a patient dying with a plug in the pulmonary artery is, in many respects, identical with that observed in cases of choleraic collapse.

M. L. Doyère, in his *Mémoire sur la Respiration et la Chaleur Humaine dans le Choléra*, Paris, 1863, has shown, by a series of most exact analyses, that one of the most constant chemical phenomena of cholera is a more or less considerable diminution of the amount of carbonic acid exhaled in proportion to the volume of oxygen absorbed by the blood. In the normal state, the volume of oxygen absorbed and of carbonic acid exhaled, are almost exactly equal. In cholera, with rare exceptions, the amount of carbonic acid given out is much less than that of the oxygen taken in. These exact observations indicate that abnormal chemical changes take place in the blood and tissues of a cholera-patient. It is probable that, during the blood-changes which result from the action of the cholera-poison, a portion of the oxygen enters into combination with the abnormal products, forming compounds which, as it were, take the place of the carbonic acid. Thus the normal respiratory changes in the lung would be interfered with; and, the morbid blood exciting the contraction of the minute pulmonary arteries, the phenomena of collapse occur.

From the study of cases of renal disease, we obtain evidence of an analogous class of phenomena. In the advanced stages of all forms of chronic Bright's disease, in proportion to the destruction of the gland-cells, we find hypertrophy of the muscular walls of the minute renal arteries and of the minute arteries in various tissues, but more especially in the skin and mucous membrane of the intestines; and, together with this arterial hypertrophy, there exists a corresponding hypertrophy of the left ventricle of the heart. The destruction of the renal gland-cells has for its necessary consequence an accumulation of urinary excreta in the blood. This urinous blood excites contraction of the systemic arteries in general, just as blood, unacrated and charged with carbonic acid, excites systemic arterial contraction. An excess of urinary crystalloids in the blood must interfere with the normal interchange of materials between the blood and the tissues; and so the blood conveyed to the kidneys will have the same abnormal relation to the gland as exists between the venous blood and the pulmonary vessels in cases of apnoea. Thus the minute renal arteries are excited to contract upon their morbid contents. This excessive contraction of the systemic arterioles in general, and of the renal arterioles in particular, results in hypertrophy of their muscular walls, and calls forth a corresponding excessive contraction and resulting hypertrophy of the left ventricle of the heart.

A careful comparison of the phenomena of apnoea or asphyxia, as it is commonly called, with those of renal disease and of cholera, appears to warrant the conclusion that one general physiological feature is common to all these cases; namely, that the interchange of materials between the blood and the tissues concerned in respiration, secretion, or nutrition, being gravely impeded, there results from this impediment a retarded circulation consequent on excessive contraction of the muscular arterioles. This excessive contraction, of course, takes place under the influence of the vaso-motor nerves; and, when prolonged, as in cases of chronic renal disease, it results in hypertrophy of the muscular walls of the minute arteries.

In the present communication, I have made no reference to those subtle and special affinities which result in the separation of peculiar products by particular secreting tissues. How it happens that urine is secreted by the kidneys, bile by the liver, and sweat by the skin, has never yet been explained. A complete explanation of the physics and the physiology of secretion is beyond the range of our present knowledge of vital phenomena.

THERAPEUTIC MEMORANDA.

TREATMENT OF SEA-SICKNESS.

HAVING been for some time engaged as surgeon on board some of the transatlantic steamships, I beg to offer a few remarks on the subject of sea-sickness, especially in respect of its alleviation by the subcutaneous method of administering morphia. Sea-sickness, as it occurs in crossing the Atlantic Ocean, sometimes assumes a very formidable character, and it would be a desideratum that the subject of its relief should be investigated more fully than it has been done heretofore. The measures that I adopted for its palliation were as follows.

I generally administered a few grains of blue pill with colocynth, or a grain or two of resina podophylli (which latter drug is especially applicable, on account of the minuteness of the dose), as I found that patients who were labouring under sea-sickness were also obstinately constipated. I then prescribed slop-diet, such as arrowroot and beef-tea, with a little brandy mixed with the former, if there were any symptoms of collapse. Sucking lumps of ice I also found to be extremely grateful, especially in the tropics. With these simple measures, together with free ventilation and absolute rest in the horizontal posture, most patients are able to get on tolerably well. In some cases, however, the indisposition assumes a more serious aspect, resulting from almost total want of food and rest, and also from mental depression, especially in females, who are apt to be hysterical. It is in these distressing cases that the hypodermic injection of acetate of morphia acts like a charm. It exerts its beneficial influence generally in from a quarter to half an hour. It soothes the patient wonderfully, allaying all that restlessness and extreme sense of illness and dependancy so often to be observed in sea-sickness. The patient is able to retain food after its administration, and thereby gain strength; and I have never seen its careful use followed by any pernicious effects.

L. TONSON COLTHURST, M.D., U.S. Mail Service.

SUBCUTANEOUS INJECTIONS OF WARM WATER.

THE substitution of warm water for morphia in subcutaneous injections for the relief of pain having lately been before the readers of the JOURNAL, I venture to add my experience, which is limited to four cases, the result being the same in all. In the first, there was pain in the course of the sciatic nerve, which lasted four or five days. At intervals of two or three weeks, when there was entire freedom from pain. The patient had not had any anodyne nor hypodermic injection. On one occasion, when the pain was severe, a subcutaneous injection of six drops of warm water was given at the seat of pain, but without any relief. The injection was repeated an hour subsequently, but with the same result. The other three cases were of carcinoma uteri. In one, the disease was at its commencement; the pain was peculiarly periodic, occurring daily at 4 P.M., and lasting several hours; sometimes through the night. One-fifth of a grain of acetate of morphia in four minims of water, given subcutaneously in the arm, subdued the pain; and, if used before the usual time of its occurrence, prevented it. Thinking the pain might be in part under the influence of fear, I advised that, without the knowledge of the patient, for two days the strength of the solution should be reduced one-half. This change was carried out; but each night the patient complained that the injections were losing their effect. Four drops of warm water were then substituted for the morphia-solution. That night the patient said the injection was useless. It is now eighteen months since the first symptoms in this case showed themselves. She now has an injection under the skin of the arm of six minims of a solution of bimeconate of morphia (one grain in six minims) three times in twenty-four hours. The solution was prepared by Messrs. Southall of Birmingham. The bottle containing it requires to be placed in a tumbler of hot water each time before use, to perfect the solution of the morphia. Emaciation precludes the administration of a larger quantity of fluid. The pain is thus kept under, though the uterus and vagina are one morbid mass; and no bad symptoms are caused by such large doses of morphia. There is a tendency to sleeplessness at night; but this is not due to pain, and is removed by ten grains of chloral-hydrate. After carefully watching the three cases of carcinoma uteri similarly treated, I arrived at the conclusion that the three women were delicate morphometers, and that a drop of the solution lost in the administration was detected by them in the result; that the amount of morphia had to be gradually increased that its efficacy might be retained; and, on this account, and on account of the insolubility of morphia salts in neutral solutions, I think a substitute for morphia as an anodyne is a desideratum. EDWIN RICKARDS, M.D., Birmingham.