led them to conclusions different from those taught by their lecturer should be also heard, instead of being ignored, or perhaps contemptuously spoken of, and practically debarred from telling the very different story the same "nature" has taught them.

Dr. Tyndall concludes his paper with the information that, if it were thought desirable, he would willingly make arrangements to enable "medical men" to see his "infusions protected and unprotected", though the apparatus is now in disarray. The "medical men" who have studied Dr. Tyndall's lecture probably do not want "to contrast the hyaline purity of the protected infusions" with the "muddy turbidity of the unprotected"; but not a few having a knowledge of the results of previous investigations would like to be informed concerning the exact method or methods pursued by Dr. Tyndall, which enabled him to so confidently conclude that not one of the hundred and thirty-nine flasks contained bacteria, and that "there was neither mistake nor ambiguity about the result". There are, unfortunately, some scientific persons, and probably not a few "medical men", who are not quite convinced that a few hundred or thousand bacteria suspended in a *clear* fluid or atmosphere would be certainly made manifest by the "searching beam".

A COMMENTARY ON DR. PATRICK BLACK'S THEORY OF ASPHYXIA.

By GEORGE JOHNSON, M.D., F.R.S.,

Professor of Medicine in King's College; Physician to King's College Hospital.

DR. BLACK has done me the favour to send me a copy of his recently published lecture on the Physiology of Respiration;* and, having carefully read the pamphlet, I venture to take up my pen in defence of the commonly received theory of asphyxia, which is assailed by Dr. Black. He tells us in his preface that "the doctrine of asphyxia, as laid down by the highest authorities, cannot, he believes, be maintained; and, as this subject is very closely connected with the heart's physiology—so closely, indeed, that his reasoning on each of these subjects must stand or fall together—it is necessary to give such an account of the latter as shall be in perfect harmony with all the phenomena of the former". He goes on to say that "the heart's physiology has been in some important points entirely misapprehended", and he again brings before the public the views to which he gave utterance more than twenty years ago in the Croonian Lectures delivered by him at the Royal College of Physicians. He now "challenges the verdict of philosophic minds on these, and respectfully asks that they may be considered".

Dr. Black's doctrine is, that "the primary end" of respiration is not the replacement of the carbonic acid of the blood by oxygen, but that "the movement of respiration is absolutely necessary to the maintenance of life, for, without this movement, the blood is obstructed in its progress through the lungs. Arrest the movement, and you have immediate asphyxia; allow the movement, even if the air admitted contains no free oxygen, and is, therefore, wholly unfitted to vivify or arterialise the blood, and the circulation will go on for a considerable time without let or hindrance". Professor Huxley, amongst other physiologists, he says, "has fallen into the error of referring asphyxia to the double cause of privation of oxygen and excessive accumulation of carbonic acid in the blood". In a subsequent part of his lecture, Dr. Black repeats the statement that, while "the movements of the chest are absolutely necessary to enable the blood to pass through the lungs, this passage of the blood is entirely independent of the chemical changes which are in a manner incidental to the movement". (P. 32.) He goes on to explain the final cause of the relative weakness of the right side of the heart. "Nature (*i.e.*, the Author of Nature), in placing this barrier to the pulmonary stream, has by this very act ensured the blood's aëration. For, had she endowed the right heart with sufficient power to sweep the hindrance before it, instead of requiring that it should be lifted away by the act of inspiration, the necessity for this act would not have been felt; medulla's function would not be invoked; the blood would pass through the lungs without her sanction; it would not put on its red garment, for it would pass unchanged. But Nature has so weakened the heart, that the barrier is effectual until another power is called in; and, as every inspiration lifts off this barrier, so the circulation through the lungs is continued, and the blood's changes are duly accomplished." (P. 33.)

I have given Dr. Black's theory as much as possible in his own words, and he will doubtless admit that I have fairly stated his doctrine.

* *Respiration; or, why do we Breathe?* A Lecture delivered before a popular audience. By Patrick Black, M.D., Physician to St. Bartholomew's and Christ's Hospitals. London: 1876. 8vo. Pp. 36.

In reply to Dr. Black, I commend to his notice Dr. John Reid's well known essay *On the Order of Succession in which the Vital Functions are arrested in Asphyxia*. He will there find that the old Hallerian doctrine, which attributed the arrest of the circulation through the lungs to the cessation of the mechanical movements of the chest, is completely disproved by very simple and conclusive experiments. I give the following in Dr. Reid's own words.

"A tube with a stop-cock was fixed into an opening in the trachea (of a dog), and one of Poiseuille's hæmadynamometers was introduced into the femoral artery, for the purpose of obtaining definite information upon the force with which the blood was transmitted along the arterial system. The stop-cock of the tube in the trachea was then shut; and, when the respiratory process had been suspended sufficiently long to cause a decided fall in the column of mercury supported by the blood sent along the femoral artery, a large bladder full of pure nitrogen, with a brass nozzle provided with a stop-cock, was fixed in the tube in the trachea, which it fitted accurately; and both stop-cocks were then opened. After the effects of the respiration of nitrogen gas had been ascertained, a bladder of the same size as the other, similarly provided with a nozzle, and full of atmospheric air, was then substituted for the bladder containing the nitrogen, and the results compared. The difference between the effects of the respiration of the nitrogen gas and the atmospheric air was most marked, and of such a nature as could not be mistaken; for, while the mercury continued to fall in the instrument during the respiration of the nitrogen gas, it rose very rapidly immediately after the atmospheric air had entered the lungs and acted upon the blood. In this experiment, the same mechanical movements of the chest, which failed to renew the free circulation of the blood through the lungs when nitrogen gas was inspired, rapidly effected that object when atmospheric air was permitted to enter the lungs, even when tried on the same animal subsequently to the failure of the nitrogen, and consequently at a more advanced stage of the process of asphyxia. This experiment was repeated several times, and, when the requisite care was taken to procure and employ pure nitrogen (unmixed with atmospheric air), invariably with the same result."

Surely Dr. Black must see that the results of this experiment are utterly irreconcilable with his statement that the circulation through the lungs is entirely dependent on the respiratory movements, and in no degree dependent on the chemical changes effected between the blood and the respired air. If the arrested pulmonary circulation were, as Dr. Black believes, a mechanical result of the suspension of the respiratory movements, the arrest of the pulmonary circulation should follow immediately upon the suspension of those movements. But this is not so. Dr. Reid, in the essay before referred to, clearly demonstrates that the first impediment to the circulation consequent on the exclusion of air from the lungs occurs, *not in the pulmonary, but in the systemic vessels*. For a period of about two minutes after shutting off the air, black blood continues to pass freely through the lungs into the left side of the heart; and a dynamometer in the femoral artery shows a continuous increase of pressure, in consequence, as Dr. Reid says, of "an impediment to the passage of the venous blood through the capillaries of the systemic circulation". After about two minutes, the pressure in the artery rapidly falls, in consequence of the impeded circulation through the lungs, and the resulting comparative emptiness of the left side of the heart.

Dr. Reid's observations on the blood-pressure were somewhat interfered with by the struggling of the animal; but, when the experiment is done upon a dog under the paralyzing influence of curare, as I have seen it done by the present distinguished Professor of the Institutes of Medicine in the University of Edinburgh, the results are in strict accordance with Dr. Reid's statement that the first effect of suspended respiration is to cause an impeded systemic circulation, while the impediment and the final arrest of the pulmonary circulation occur later. The most probable explanation of the impeded circulation through both systems of vessels is that which attributes it to the contraction of the muscular arterioles, excited by the influence of the abnormal blood upon the vaso-motor nerves and centre. (See my paper on Certain Physical Phenomena connected with the Circulation, etc., *BRIT. MED. JOURNAL*, Jan. 8th, 1876, p. 44.)

Dr. Black makes the surprising statement that the two ventricles of the heart discharge "their contents through orifices of exactly the same diameter". I can imagine Dr. Herbert Davies' dismay at seeing such an assertion, after his ingenious demonstration of the mathematical relations of the areas of the four orifices of the heart. The relatively greater thickness of the walls of the left ventricle is required, not merely, as Dr. Black states, to overcome the greater impediments which exist in the terminal systemic vessels, but, the orifice of the aorta being narrower than that of the pulmonary artery, a more vigorous left

ventricle is necessary to drive an equal volume of blood in the same time through the smaller orifice of the aorta.

Dr. Black speaks repeatedly of the weakness of the right heart. It is, of course, absolutely weaker than the left; but, relatively to the work which each heart has to perform, the two hearts are of equal strength. If they were not, they could not work so harmoniously together, as they often do, for fourscore years and upwards. I think that I have now adduced sufficient reason for my belief that Dr. Black's "explanation of the heart's thin pulmonary chamber" is untenable, and his theory of asphyxia inconsistent with unquestionable facts.

ON A CASE OF LARGE OSSEOUS TUMOUR, REMOVED SUCCESSFULLY, WITH ANTISEPTIC PRECAUTIONS, FROM UNDER THE LIGAMENTUM PATELLÆ.

By THOMAS ANNANDALE, F.R.S.E.,

Surgeon to the Edinburgh Royal Infirmary, and Lecturer on Clinical Surgery.

THE following case seems to me well worthy of a special report, for two reasons: first, from its great rarity; and, second, as an illustration of the value of antiseptic treatment in operations on or near important joints.

D. D., aged 62, was admitted into my wards on the 19th September last. Three years before his admission, he observed a small swelling at the lower part of the anterior aspect of his right knee-joint, and since then this swelling steadily increased in size, and, from its bulk, lately became troublesome, and interfered with the proper movements of the joint.

An examination determined the existence of a tumour on the anterior aspect of the joint: it was of the size of two closed fists, was hard to the touch, and was circumscribed and free on all sides except at its base, which appeared to have a firm attachment to the head of the tibia. The tumour, when grasped, felt quite fixed to the bone when the

knee was flexed; but as, in the extended position of the joint, it admitted some lateral movement, it was evident that no real connection to the bone existed. The patella was displaced upwards and inwards, and could be felt immediately above the tumour. The movements of the joint were limited, especially that of flexion. Fig. 1 is taken from a cast of the knee and tumour, and well illustrates the appearance of the parts before the operation.

On September 22nd, I exposed the anterior aspect of the tumour by a free incision carried along its outer aspect, and a second transverse and smaller one in a direction inwards across the front of the knee.

Fig. 1.

The first incision was made down to the tumour, in a direction parallel and external to the ligamentum patellæ and patella. The second one was superficial, and only divided the skin and cellular tissue. The object of these incisions was to thoroughly expose the tumour, and, at the same time, avoid as much as possible injury to the synovial membrane of the joint and to the ligament of the patella. When the tumour was brought into view, it was found that the ligament of the patella was stretched quite it, and ran along a groove on its superficial aspect. The tumour had apparently originated in the bursa under the ligament, and the connection of this ligament to the tumour explained its fixed condition, which simulated an osseous attachment. By careful dissection, I succeeded in removing the tumour without injury to the ligament; and, although a large surface of syno-

vial membrane was exposed, the cavity of the joint was not opened into.

The operation was performed under the carbolic spray, and the wound was dressed with the usual precautions. Not the slightest local or constitutional irritation followed the operation, and the patient was dismissed on October 29th, the wound being soundly healed, and the joint and limb perfectly movable and natural in every respect.

Fig. 2 illustrates a sectional view of the tumour, which was composed

Fig. 2.

of cancellated osseous structure, with here and there portions of cartilaginous tissue. The tumour was surrounded by a strong fibrous capsule.

REMARKS.—From the history and situation of this tumour, I am inclined to think that it originated in connection with the bursa under the ligamentum patellæ. Fibrous enlargements of this bursa have been recorded, but the occurrence of an osseous growth in this situation is not referred to in recent works on surgery. The tumour had no attachment to, or connection with, any bone; and it may, therefore, be classed under those isolated osseous growths sometimes met with in the areolar and other textures, and of which Paget (*Lectures on Surgical Pathology*, 3rd edition) remarks, that they "are interesting for little more than their rarity".

The successful result obtained in this case, and the perfect recovery of the patient with a completely movable joint, must be attributed, in my opinion, to the employment of antiseptic precautions. By no other means do I think that so free an exposure of the synovial membrane could have been made without the slightest irritation resulting.

ALEPPO BOIL OR "BISKRA BOUTON".

By GEORGE THIN, M.D.

THE Aleppo boil—*mal* of Aleppo—was discovered by Richard Pococke, an English traveller, in 1745, and formally introduced into medical science as an independent malady by Dr. Alexander Russell in 1756. The credit of the discovery belongs, therefore, to England. But Aleppo has no monopoly of the disease, which, we are told, is also endemic in Cairo, Alexandria, Bagdad, Candia, Cyprus, and as far westwards as Biskra, an Algerian station in the Sahara. Such is the terror excited by this perennial plague, that Turkish officials and soldiers who are sent to Aleppo receive double pay as compensation for the risk they incur by being exposed to its attack.

The two observers who have most recently made inquiries regarding its nature have arrived at contradictory conclusions. Whilst Dr. Van-dyke Carter supports the theory which regards it as a specific malady, Dr. Geber regards the whole thing as resting on mistaken diagnosis, maintaining that there is not, and never has been, any such disease, and that the host of observers, from Dr. Russell downwards, have failed to recognise the cases of syphilis, lupus, scrofula, chronic eczema, and badly treated dermatitis, which passed as "bouton". Dr. Geber's