ventricle is necessary to drive an equal volume of blood in the same time through the smaller orifice of the aorta.

Dr. Black speaks repeatedly of the weakness of the right heart. It is, of course, absolutely weaker than the left; but, relatively to the work which each heart has to perform, the two hearts are of equal strength. If they were not, they could not work so harmoniously together, as they often do, for fourscore years and upwards. I think that I have now adduced sufficient reason for my belief that Dr. Black's "explanation of the heart's thin pulmonary chamber" is untenable, and his theory of asphyxia inconsistent with unquestionable facts.

ON A CASE OF LARGE OSSEOUS TUMOUR, REMOVED SUCCESSFULLY, WITH ANTISEPTIC PRECAUTIONS, FROM UNDER THE LIGAMENTUM PATELLÆ.

By THOMAS ANNANDALE, F.R.S.E.,

Surgeon to the Edinburgh Royal Infirmary, and Lecturer on Clinical Surgery.

THE following case seems to me well worthy of a special report, for two reasons: first, from its great rarity; and, second, as an illustration of the value of antiseptic treatment in operations on or near important joints.

D. D., aged 62, was admitted into my wards on the 19th September last. Three years before his admission, he observed a small swelling at the lower part of the anterior aspect of his right knee-joint, and since then this swelling steadily increased in size, and, from its bulk, lately became troublesome, and interfered with the proper movements of the joint.

An examination determined the existence of a tumour on the anterior aspect of the joint: it was of the size of two closed fists, was hard to the touch, and was circumscribed and free on all sides except at its base, which appeared to have a firm attachment to the head of the tibia. The tumour, when grasped, felt quite fixed to the bone when the

knee was flexed; but as, in the extended position of the joint, it admitted some lateral movement, it was evident that no real connection to the bone existed. The patella was displaced upwards and inwards, and could be felt immediately above the tumour. The movements of the joint were limited, especially that of flexion. Fig. 1 is taken from a cast of the knee and tumour, and well illustrates the appearance of the parts before the operation.

On September 22nd, I exposed the anterior aspect of the tumour by a free incision carried along its outer aspect, and a second transverse and smaller one in a direction inwards across the front of the knee.

Fig. 1.

The first incision was made down to the tumour, in a direction parallel and external to the ligamentum patellæ and patella. The second one was superficial, and only divided the skin and cellular tissue. The object of these incisions was to thoroughly expose the tumour, and, at the same time, avoid as much as possible injury to the synovial membrane of the joint and to the ligament of the patella. When the tumour was brought into view, it was found that the ligament of the patella was stretched quite it, and ran along a groove on its superficial aspect. The tumour had apparently originated in the bursa under the ligament, and the connection of this ligament to the tumour explained its fixed condition, which simulated an osseous attachment. By careful dissection, I succeeded in removing the tumour without injury to the ligament; and, although a large surface of syno-

vial membrane was exposed, the cavity of the joint was not opened into.

The operation was performed under the carbolic spray, and the wound was dressed with the usual precautions. Not the slightest local or constitutional irritation followed the operation, and the patient was dismissed on October 29th, the wound being soundly healed, and the joint and limb perfectly movable and natural in every respect.

Fig. 2 illustrates a sectional view of the tumour, which was composed

Fig. 2.

of cancellated osseous structure, with here and there portions of cartilaginous tissue. The tumour was surrounded by a strong fibrous capsule.

REMARKS.—From the history and situation of this tumour, I am inclined to think that it originated in connection with the bursa under the ligamentum patellæ. Fibrous enlargements of this bursa have been recorded, but the occurrence of an osseous growth in this situation is not referred to in recent works on surgery. The tumour had no attachment to, or connection with, any bone; and it may, therefore, be classed under those isolated osseous growths sometimes met with in the areolar and other textures, and of which Paget (*Lectures on Surgical Pathology*, 3rd edition) remarks, that they "are interesting for little more than their rarity".

The successful result obtained in this case, and the perfect recovery of the patient with a completely movable joint, must be attributed, in my opinion, to the employment of antiseptic precautions. By no other means do I think that so free an exposure of the synovial membrane could have been made without the slightest irritation resulting.

ALEPPO BOIL OR "BISKRA BOUTON".

By GEORGE THIN, M.D.

THE Aleppo boil—*mal* of Aleppo—was discovered by Richard Pococke, an English traveller, in 1745, and formally introduced into medical science as an independent malady by Dr. Alexander Russell in 1756. The credit of the discovery belongs, therefore, to England. But Aleppo has no monopoly of the disease, which, we are told, is also endemic in Cairo, Alexandria, Bagdad, Candia, Cyprus, and as far westwards as Biskra, an Algerian station in the Sahara. Such is the terror excited by this perennial plague, that Turkish officials and soldiers who are sent to Aleppo receive double pay as compensation for the risk they incur by being exposed to its attack.

The two observers who have most recently made inquiries regarding its nature have arrived at contradictory conclusions. Whilst Dr. Van-dyke Carter supports the theory which regards it as a specific malady, Dr. Geber regards the whole thing as resting on mistaken diagnosis, maintaining that there is not, and never has been, any such disease, and that the host of observers, from Dr. Russell downwards, have failed to recognise the cases of syphilis, lupus, scrofula, chronic eczema, and badly treated dermatitis, which passed as "bouton". Dr. Geber's