duction of material which, in their deficiency, perhaps cannot be turned to vital purpose.

The measures I have commonly—and numerous experience justifies me in saying beneficially—used, have been liberal diet, with beef-tea, Liebig's extract, and vegetables which abound in potash-salts, together with liquor potassæ, phosphate of potash, iron, quinine, and sometimes cod-liver oil. The alkaline remedies have been urged with the most obvious advantage during the persistence of suppuration.

Nothing now remains for me to do but to thank this accomplished and critical audience for the forbearance with which they have listened to many tedious and possibly familiar details. I trust I have shown how the varieties of renal disease are not only complicated but often inextricable; but how, according to the bias which prevails in each instance, different classes of secondary results follow; that among these the cardio-vascular change, however else it may arise, is a result of the vascular tension of uræmia, which, though more common with some forms of disease than others, may ensue upon any. Finally, in touching upon treatment, I have sought to show in what instances it is to be guided by a consideration of the special organic state, and where more frequently by the results which it has produced upon the patient.

THE GOULSTONIAN LECTURES

ON

THE NATURAL HISTORY OF PULMONARY CONSUMPTION.

Delivered at the Royal College of Physicians, London.

By A. B. SHEPHERD, M.A., M.B., F.R.C.P.,

Physician to the Victoria Park Hospital for Diseases of the Chest; Assistant Physician to St. Mary's Hospital; etc.

LECTURE III.—*Concluded.*

THE title I chose for these lectures debars me from speaking with the faintest air of authority, or to any extent, of the various drugs used to oppose the ordinary course, and therefore considered to have some effect on the natural history of consumption. The key to any rational treatment of the disease is probably to be found in a careful consideration, in any individual case, of the present condition of the phthisical process—whether it be still active, or for the time being stationary; and of the constitutional habit of the patient—whether of the weakly excitable type characterised by excessive reaction, or of the vigorous, more heavy one, in which reaction is less marked (*see Braun*). But of one medicine, or rather food, I must speak briefly, because, in its effects on the patient, it offers a distinct bedside result. To use the words of a former President of the College, “you may learn much (and so, no doubt, you ought) by prying into the arcana of the night-chair” (*Watson, Lectures, 1857, vol. ii, p. 715*); and yet, in these days of numberless instrumental aids to diagnosis, the learning gained from that inspection, like the splendid knowledge of the pulse and appreciation by hand of increased temperature which marked a generation happily not passed away, seems to be in danger of leaving us, who more unworthily trust to mechanical assistance. But an inspection of the night-chair of a consumptive patient taking large quantities of cod-liver oil will reveal, as perhaps physiological theories ought to have predicted, a very great increase of fatty or oily matters evacuated. This, again, is no new observation, though it is one, I think, which may well teach us its lesson. Over twenty years ago, Sir William (then Dr.) Gull had recorded (*Guy's Hosp. Rep., ser. iii, vol. i, p. 369*) his own observations on “fatty stools from disease of the mesenteric glands”, and written down in no half-hearted way his opinion as to the careless and utterly unscientific use of cod-liver oil. It cannot be denied that, by some means or other, this drug had, I might even say still has, come to be looked upon as a “specific” for “consumption”; and I use the word as out-patients use it, as I have used it in these lectures—as consumption from disease of the lungs. Patients themselves tell us they must have phthisis, because the doctor who prescribed for them gave them cod-liver oil. At any rate, in hospital practice later, I repeatedly examined the faces of phthisical patients, perhaps because the teaching of the wards of Guy's was unconsciously with me; and I found, as every one who performs the same experiment will find, that in these phthisical patients, beyond a certain amount of oil administered, varying of course in individuals, but, on the average, four drachms daily, large quantities of it pass through the intestines almost unchanged. And to this simple waste, not too much to be decried, is superadded very often increased disturbance of diges-

tion and nutrition. Still, to quote Sir W. Gull, “if it be ever admissible to limit the attention to one organ in treating disease, it is so” here, where “the condition of the gastro-intestinal membrane is of paramount importance”.

And what I have said with respect to cod-liver oil, and its possibly harmful waste, applies quite as much, if not more, to a highly vaunted drug, which, to judge from the weight of an ordinary pig's pancreas, would demand for each patient, if I be not very wrong, hecatombs of victims; unless, indeed, there be a peculiar breed of swine with a larger pancreas,

“ὄψις θρεψέ σὺβάτης ὄρχαμος ἀνδρῶν.”—*Odyssey, 14, 22.*

Only a fortnight ago, a gentleman who had taken the emulsion had noticed for himself the fatty nature of his stools: a fact which my remark as to one of the effects of cod-liver oil led him to mention.

In approaching rapidly the close of my subject, I cannot but congratulate myself, representing for the moment the youngest Fellows of the College, that the work of the very latest time leads to a conclusion which in its pathological results, though not at all in its reasoning from those results, has the support of the perhaps oldest living authority on pulmonary consumption in this country. Dr. C. J. B. Williams, in the discussion to which I have already referred (*Path. Soc. Trans., vol. xxiv, p. 340*), has given us his “explanation of the twofold seat and origin of tubercle, or rather of consumptive disease: 1. Lymphatic, which is miliary, infective, and scattered; 2. Inflammatory, which is diffused in form and local in extent.”

I would call the attention of those gentlemen who wish to examine them after the lecture to the microscopical preparations placed on the table of the various forms of pulmonary consumption described. [Explanations of the preparations, etc., followed.]

To sum up briefly the views, by no means original, which I have attempted to support respecting pulmonary consumption in these lectures, I am disposed to conclude:

That, in the affection to which the name “acute general miliary tuberculosis” is given, the hard nodules occurring in the lungs are due mainly to growth of the perivascular and peribronchial cords, together with alveolar catarrh of smaller or larger extent.

That this perivascular growth is secondary to a cheesy focus not necessarily present in the lungs.

That the commonest form of pulmonary consumption in this country is due, in the first instance, to an intra-alveolar and intrabronchial catarrh, accompanied by proliferation of the epithelium more or less rapid; followed generally by impermeability and disappearance of the alveolar capillaries, and by changes in the septa of the alveoli; and resulting in the destruction of lung-tissue: the whole process giving rise to the clinical symptoms of phthisis.

That the same clinical symptoms of phthisis are caused by the pathological changes present in chronic bronchitis and emphysema, and in other cases by fibroid growth of the alveolar septa themselves, leading to the disappearance of their cavities.

That catarrhal pneumonia is the most common accompaniment of any affection of, or even injury to, the lungs; and that its occurrence often gives rise to the appearance of bodies possessing, to the naked eye, all the characteristics of the “tubercle” of authors.

That there is nothing in common between the pathological results of acute tuberculosis and those of chronic cheesy degeneration of the lungs, except it be in the occasional presence in the latter of the secondary lymphatic (?) products almost always, if not always, to be found in the former.

That the cheesy form of pulmonary phthisis ought not to be looked upon as one absolutely hopeless and fatal.

That, in the majority of cases of this form of phthisis, the whole bent of our treatment should be against repeated and chronic inflammations of the lung, in the sense of catarrh of the air-vesicles and finer tubes, and as far as possible in aid of increased assimilation and nutrition, especially in the encouragement of those methods of hygiene which consist not only in change of air and climate, but in the more ordinary matters of baths and suitable clothing.

And, sir—to pause for one moment—the mention of these hygienic measures reminds me that, since our meeting here on Wednesday last, the College mourns the death of one of her greatest ornaments and most illustrious Fellows, Professor Parkes of Netley. Further of him to-day it is not for me to speak.

Portal, publishing his work on the *Nature and Treatment of Pulmonary Phthisis* in “the first year of the French Republic”, as his title-page tells us (1792), speaks more boldly than did Bayle, who, with some sort of solemn religious feeling, confesses that to a large extent he has not dared to “explain the part that Nature plays in the production of phthisis”. (*Op. cit., Préf., p. xiii.*) But Portal, conceding that medicine is but a branch of natural history, thought he could not do better

than conform strictly to the method followed in the different portions of the latter; that is, to give a history of facts observed, arrange them in order, and draw from their comparison the true character of the object treated of. "We ought never," he says, "to hope to guess at or to domineer over Nature; we can but listen in silence to, and meditate attentively over, the phenomena we observe; we can but discuss them carefully, and gather from them such truths as shall be useful." My aim has been to work after this fashion.

In conclusion, sir, if I have not in very word complied with the conditions of the bequest of Dr. Theodore Goulston, "that a dead body was, if possible, to be procured, and two or more diseases treated of", I have at least brought the pictured results of the *post mortem* room within the College, and treated, however imperfectly, of an affection which is Protean in its anatomy, if not in its features. Following out, at any rate, the feeling which, as Censor of this College between the years 1615 and 1626, he had probably attempted to inculcate, that not without the scalpel would the bounds of knowledge be widened, I may say, in the words which his perhaps greater successor in this Fellowship makes use of, "Non solum manus pro virili medicas languidis corporibus adhibendi, verum etiam defunctorum cadavera introspectiendi quamsepissimè ansam arripui" (Morton, *op. cit.*, Ad Lect.). Yet it may be that whatever I have said on the subject I have had the honour to bring before you only illustrates once more the twice-told sentence of Bacon (*Nocturn Organon*, liber i, xlv; and *Advancement of Learning*, book ii): "The spirit of man feigns in Nature a simplicity and uniformity greater than really is". It may be that the conclusions to which I have been almost irresistibly led as to what seems to me to be the simple nature of pulmonary consumption may fail to be real, when scanned by some more searching eye than my own; and, if it be so, for work at least honestly attempted some consolation may be found in the words in which Heberden speaks of the uncertainty of modes of treatment: "No aphorism of Hippocrates holds truer to this day than that in which he laments the length of time necessary to establish medical truths, and the danger, unless the utmost caution be used, of our being misled, even by experience." (*Comm.*, 1802, p. 71.)

ON THE OPEN AIR TREATMENT OF CONSUMPTION.

By JAMES BLAKE, M.D., F.R.C.S., San Francisco, California.

AN article in the JOURNAL for October 24th, 1874, recalls an intention I have had for some time of sending a communication on the open air treatment of consumption, a plan I have advocated for many years, both on the grounds to which Dr. Marcet alludes as to its being a septicæmia, and also on account of the advantages which such a treatment offers for improving the digestive organs. In a paper published in the *San Francisco Medical and Surgical Journal*, in 1860, I pointed out the advantages to be derived to the digestive functions by living in the open air; but looking at the septicæmic element of the disease, no other treatment, it appears to me, can so effectually combat it as living in the open air, the only condition in which a patient with diseased lungs can avoid rebreathing the poisoned air he has expired, laden with the germs for intensifying the putrefactive processes going on to his lungs. There is undoubtedly a germ of truth in the theory of Dr. Mac Cormac of Belfast, that the chief cause of consumption is rebreathed air, but not, I think from its being overcharged with carbon, as he supposes, but because it is loaded with a much more subtle poison in the putrefactive germs which it contains. Some twelve years ago, I published some cases in the *American Journal of Medical Sciences*, showing how many cases of consumption had been arrested, and some cured, by what I called the open air treatment of the disease; and a longer experience has convinced me that this method offers the best chance for our consumptive patients. But, before the profession can be induced to employ it, I am aware that an accumulated mass of prejudice has to be removed, not only amongst physicians, but more particularly amongst the public, as regards the evil of exposure and living in the open air. The idea of advising a patient in the third stage of consumption, suffering from cough and night-sweats, to sleep in the open air, is a proposition which in England, I am aware, would be considered not only as dangerous, but almost as a sign of lunacy. Even here, where it is no very uncommon thing for persons to sleep out of doors, and where the dewless nights of our mountain ranges during our rainless summers render any covering but blankets and a tree quite superfluous, I often met with objections to following such a course. And yet I am convinced that it is the best method that we possess for arresting and curing consumption. In England possibly, and in fact in most parts of Europe, the occasional summer

rains and the absence of dry mountain ranges, with their pine-covered ridges, offer obstacles to such a treatment being fully carried out; but the principle once recognised, a great deal may be done even there towards employing it. In order, if possible, to remove to a certain extent this prejudice, I would state that I have never in a single instance seen any ill effects result from it. I have sent out patients to sleep in the open air who were so far reduced that they could not even ride on horseback, but had to be conveyed in a carriage. The difficulties and annoyances of living in the open air are, I believe, entirely in the imagination. The most agreeable holidays I have ever spent have been whilst camping out with a sensible party of ladies as well as gentlemen, and never have we broken up camp to return to houses without regret by all the members of the party. When ladies are in the party, it is better to have a tent, the front of which must always be fully open at night. During the winter months, the southern part of the State about San Diego or Santa Barbara is the most desirable place for patients, as there are seldom more than a dozen rainy days during the year; but in the summer the coast range of mountains north of San Francisco offers by far the most congenial climate, far better than that of the Sierras, owing to the greater equability of the temperature. From the middle of May to the end of October, as a general thing, living in the open air can be enjoyed without any fear of rain. As the summer heat increases, the higher mountains, up to 4,000 or 5,000 feet, ensure an agreeable climate, where the thermometer never rises above 85 deg. (this, in the dry air of the mountains, is about the same, as far as our sensations are concerned, as 70 deg. in England), and never descends below 55 deg., ranging generally from 60 deg. to 75 deg. during the twenty-four hours. As a general thing, at height above 1,500 feet, the camp can be made in pine-woods, and I believe that there is something antiseptic in the exhalations of these trees; certain it is that they impart a most agreeable odour to the air, particularly in warm weather. As for the cost, the expense is slight, as there are no hotel bills to pay; the journey across the continent is now rendered so easy that an invalid can generally support it without inconvenience, and in fact improve during the trip. In the cold weather, the trip can be made by the steamer to Colon, and thence to San Diego by the Pacific Mail Company's ships.

THERAPEUTIC MEMORANDA.

CASE OF RHEUMATIC FEVER TREATED BY SALICINE.

MISS O. R., a strong healthy girl, aged 15, came under my care on May 3rd, suffering from rheumatic fever. She had been ill for four days, but had not been seen by any medical man. On May 3rd, she had high fever; pulse 120; temperature 105 deg. The tongue was moist and furred. The knees and ankles were painful and swollen; the right foot and ankle cold and livid. I prescribed bicarbonate of potash in ten-grain doses every third hour, and a linseed-meal poultice to the right knee and foot; the nourishment to consist of beef-tea and milk. On May 5th, the bowels had been moved four times during the night. She did not sleep, and was very thirsty. The pain had moved to the wrist and knuckles. Morning temperature 105 deg.; pulse 126; evening temperature 105.6 deg. I discontinued the bicarbonate of potash, and ordered chalk mixture and catechu, and the joints to be enveloped in cotton-wool. On May 6th, she was very delirious during the night. The bowels had been moved six times. The urine was scanty; I did not see any of it. Owing to her helplessness and pain, she would not use the bed-pan. Temperature, morning 105 deg.; evening 105.3 deg.; pulse 120, very weak. This day I had the advantage of a consultation with my friend Dr. McClelland of Newtown-Mount-Kennedy. The pain was principally in the back and hips. The heart-sounds were normal.—May 7th. She had a very restless night. The bowels were moved several times. She had perspired freely since the commencement of her illness. The tongue was furred and moist; the knees were swollen and very painful. She could not bear the weight of the bedclothes. Pulse 120; temperature, morning 105.2 deg.; evening 105.2 deg.

From the favourable mention made of salicylic acid and salicine in the treatment of rheumatic fever in the BRITISH MEDICAL JOURNAL of May 6th, I determined to give salicine a trial (not being able to procure the acid at the time), and ordered twelve grains every third hour, discontinuing all other medicine. On May 8th, she had a better night. The pain had moved to the shoulders and back of the neck. The bowels were moved only once. Pulse 110, and strong; temperature, morning, 103 deg. (a fall of 2.2 deg.). She said she felt better, and thought the powders had done her good. Temperature, evening,