kindle a fire which it is beyond our power to extinguish. Let us consider well our responsibility, that we may not be the means of driving our fellow creatures into temptation. Let us take care that no unhappy wreck can point to us as the cause of her ruin, and if, by the reading of this paper, but one British maid or matron be kept out of temptation, or arrested on the road to ruin, I will not consider that it has been read in vain.

OBSERVATIONS ON THE EFFECTS OF CUCA LEAVES.

By WALTER BERNARD, L.K.Q.C.P.I., Londonderry.

HAVING read the interesting observations of Sir R. Christison on the effects of cuca on himself, as reported in the JOURNAL of April 29th, I determined to try a similar experiment on myself. I therefore procured, from Messrs. Duncan and Flockhart of Edinburgh, a package of the leaves, which I found to be a pretty fair sample of the article, if I may judge by Sir Robert's description, only that the leaves were green rather than "deep green", as in his account, and did not occasion such warmth in the mouth. I selected for the experiment a day with the maximum of fatigue in it. This was Monday, May 22nd, 1876. I was constantly on foot from 8.45 A.M. to 10 P.M., with the exception of an hour for dinner (from five to six o'clock); and my dinner consisted of roast lamb, some vegetables, no stimulants, and was followed by tea at seven o'clock, and mechanical rest, but no sleep up to the hour (12.15 A.M.) when I commenced to prepare for my trial trip. Thus, without food or drink since seven o'clock, I set out at 1 A.M., accompanied by Professor Leebody, analyst for this city, for a twenty-four miles' walk, and the ascent at the end of it of a hill 1,381 feet high. Commencing my journey, I certainly did not feel inclined for much exertion; but after some miles brisk walking, I felt more equal to the work. The renal secretion, noted every two hours by my friend, shows the subjoined results. We did 13½ miles in three hours, giving an average of 4½ per hour. We could have done 4¾ easily, as a smooth level footway extended to this point, but rain fell heavily for the first two hours and a-half, rendering the footpath slippery, and thus interfering with a quicker pace. Up to this time, we did not feel fatigued. Here we commenced, at the end of three hours, to chew forty grains of the cuca. We traversed a very bad hilly road, strewn with loose stones, which considerably impeded our progress. Walking under these difficulties, and now fatigued, we found no material benefit from chewing and swallowing the active principle of the leaves; a doubt, however, remained in my mind as to this conclusion, as I fancied my step, on ascending the little hills, was more elastic, and the feeling of fatigue slightly mitigated. This idea soon ripened into a belief. At six o'clock, and not far from our destination, each of us had a second dose of forty grains. I now resolved to finish well up by giving it a fair trial. Testing my strength to the utmost, and chewing quickly, I advanced towards the hill with a rapid pace, leaving Mr. Leebody in the rear. Arriving at Mamore gap, I made a very rapid ascent of Mamore hill, 1,381 feet high. Hurried breathing and fatigue began to be felt when I was about half-way up. Here I commenced to chew another forty grains; and, before I arrived at the top, though I still continued my rapid pace, power was restored. My breathing was considerably relieved. I felt in excellent spirits, and neither hungry nor thirsty. It was seven o'clock when I arrived at the top, somewhat later than we expected; but the loss of time was mainly due to the zig-zag course I followed in endeavouring to avoid the loose stones. Yet, were it not that along the rough road we were cheered by a beautiful clear morning (for all clouds and threatening of rain had vanished) and an invigorating balmy breeze from off the sea, our arrival would have been somewhat later. I remained ten minutes on the top, admiring the coast line, the entrance to Lough Swilly, the cultivated lands lying low and quite flat at the east of the hill between me and the sea, as well as the grandeur of the Donegal mountains free from haze. I felt now quite refreshed and buoyant, with no sensation of hunger, thirst, or fatigue. On descending, I bounded like a doe, jumping from knoll to knoll. Some short distance down, I met my companion, making for the top. I asked him had the cuca any effect on him? He said not. I offered him another forty grains, but he would not accept it, being somewhat disheartened. He was, however, still chewing what he had remaining of the second forty grains. I left him, and made my way to a house where my servant had put up my horse and vehicle, which followed us during our course. On arriving here, I was cool and comfortable; for soon after I had put on the rapid pace I had become bathed in perspiration from the lumbar region upwards. Gently

retracing my steps from the foot of the hill afforded me time to be come cool.

On leaving home, my pulse was 60; temperature 98 deg. Now the pulse was 94; temperature 96 deg. I waited a quarter of an hour for Mr. Leebody, and was agreeably surprised to find him beginning to change his views as to the efficacy of the leaf. He had now lost almost all feeling of lassitude. But I refer to his own notes, which are appended to this. We had determined to walk home, to give the cuca a fuller and fairer trial; but both being now agreed in thinking that there was good service rendered to us by its use, and that a further prosecution of the experiment would not add much more accurate knowledge to that which we had now gained, we decided to drive—first walking briskly a long high hill—and we arrived at home at 10.30. The greater part of the way home I had a strong taste of the shrub in my mouth, resembling somewhat that of tea. On my arrival, I did ample justice to a good breakfast. I worked unceasingly and willingly all day, and did not feel at all fatigued, and did my work on foot. I had no desire to sleep. Those who knew that I had been walking for so many hours during the night and morning, mentioned that I neither looked jaded nor worn, and I certainly did not feel so. Moreover, I state positively (and this can be substantiated by my wife) that when I have had occasion to sit up watching, I have been fagged and tired the next day, having browache not unfrequently from exhaustion. I felt thirsty at two o'clock, and had a small bottle of Apollinaris water. I dined at five on mutton, pancakes, and four glasses of good Burgundy. At six o'clock, I retired to my room to write out my notes, but felt disinclined for work for the first time during the day. This disinclination was excited by the Burgundy, as alcohol, in any shape or form, has always had this effect upon me. At seven o'clock, I took eight ounces of tea, which so refreshed me that I was able to complete my notes, and finished them at nine o'clock. I took them across the street to compare them with those of Mr. Leebody. I retired to bed at a few minutes after ten o'clock; fell at once into a sound sleep, and awoke refreshed at a quarter to six o'clock. I then had a bath and was dressed at seven, and I feel all right to-day (Wednesday, May 24th).

In May 1873, I walked twenty-seven Irish miles with Mr. Stewart, manager of the Bank of Ireland in Carlow, and Mr. Craig, manager in this city, taking twelve hours to accomplish the journey, and dining and resting; drinking one bottle of porter and a glass of whisky, with a view to assist my progress; and, on another occasion, I did eighteen miles, and took the same amount of stimulants. Long before reaching home, I felt extremely fatigued, and on both occasions did not sleep for the first two or three hours, owing to palpitation and restlessness. Since then, I have found, by experiments on myself, that although a stimulant restores power for the first few miles, it has a strong tendency to induce depression of spirits, lassitude, and disinclination for mental or bodily work. Further, it helps to make me restless in the beginning of the night, waking with a dry tongue in the morning. I had not my usual accompaniments of fatigue on lying down, *i. e.*, palpitation, and a dull aching pain in the left eyebrow. This will lead me to try the effects of cuca as a cardiac tonic.

In conclusion, I may mention that I am of opinion the specimen I received was not so good as that described by Sir Robert Christison, as it was slow in its action, and required a larger quantity to produce good effects. Having many times during my life undergone bodily fatigue from watching, and from long walks, and having suffered effects such as I have before mentioned, I think it is not a hasty generalisation to come to that, on this occasion, the cuca prevented the browache, palpitation, and restlessness, as well as afforded relief, for which it is, in my opinion, becoming justly celebrated.

THE ACTION OF CUCA.

By J. R. LEEBODY, M.A.,

Professor of Experimental Science in Magee College, Londonderry; and Public Analyst for the City and County of Londonderry.

I HAVE made the following notes of a walk taken by me on May 23rd, in company with my friend Dr. Bernard, for the purpose of testing the properties of cuca.

To insure thorough fatigue on my part, I took, in addition to the exercise involved in my ordinary work, a long row on the river on the afternoon of May 22nd. I had a steady pull of over three hours, for the greater part of the time against wind and tide. The consequence was, that when I went to bed at 10.45 P.M., I was more than usually tired. I slept till 12.15 A.M., and then rose, feeling sleepy and fatigued. At 1 A.M., we started for our walk. Before starting, I ate one ounce of bread, to prevent headache, from which I usually suffer when I take

exercise in the morning without previously partaking of food. At 4 A.M.—having walked thirteen miles and a half along a level road—we each chewed forty grains of the leaf. We then walked seven miles along a very rough mountain-road, continually ascending and descending steep hills. At 6 A.M., we had reached the base of a mountain which we proposed to ascend. We each took forty grains more *cuca*. I walked steadily up the hill at a moderate pace. Dr. Bernard, on the other hand, walked very rapidly up, and met me on his return journey before I had quite reached the summit. I rested for seven minutes on the top of the hill, and then descended and rejoined Dr. Bernard, who had completed the ascent and descent in ten or fifteen minutes less than I occupied. We then walked half a mile up a steep hill, got on a car, which was waiting for us, at 7.45 A.M., and, after a rapid drive, reached Derry at 10.30 A.M. I then had breakfast; and after breakfast, feeling drowsy, I lay down without undressing, and slept for an hour and a half. I rose at 1 P.M., spent the rest of the day attending to my usual duties, had also an hour's row on the river, went to bed at 10.30 P.M., and slept soundly till eight the next morning.

I noted the sensations experienced by me during the day as follows. On starting at 1 A.M., I was sleepy and tired; walking in the open air refreshed me, however, and at 4 A.M. I was not excessively fatigued, but was somewhat hungry. On chewing forty grains of *cuca*, my hunger ceased; but I did not notice my sense of fatigue diminished. At 6 A.M. I was very tired, and I did not notice any special effect produced by chewing the second forty grains of *cuca*. From this time, however, my fatigue did not increase. On descending the hill I felt hungry, and I then ate two ounces of bread. On arriving in Derry, my feeling of fatigue was quite gone, and I had an excellent appetite for my breakfast. Throughout the afternoon and evening I felt no more fatigue than I usually do after a busy forenoon's work in my laboratory.

On neither occasion after chewing the leaf was I conscious of its producing any special effect on me, either in diminishing fatigue or causing increased elasticity of spirits; and I was therefore at the time disappointed in my expectations, and somewhat sceptical as to the restorative properties of *cuca*. It was only when I had completed my walk and compared my sensations with those I certainly would have experienced had I attempted the same exertions without *cuca*, that I began to recognise its power; and when I found myself during the afternoon going about my usual avocations without being conscious of any feeling to remind me that I had spent the morning in an unusual manner, I fully appreciated its value.

The total distance walked was twenty-four miles, ten of which were along a most fatiguing mountain-road. The hill ascended was 1,381 feet high. The total nutriment taken by me before breakfasting at 11 A.M., was three ounces of bread and eighty grains of *cuca*.

The following are the results of three examinations of the urine.

	Quantity.	Sp. gr.	Total solids.
At end of first 2 hours	2,844 grains.	1.028	185.50 gr.
„ second „	1,750 „	1.028	114.20 „
„ third 2½ „	1,750 „	1.0275	110.25 „

The specific gravity and total solids were very high.

SUPPURATION OF THE MIDDLE EAR IN RELATION TO LIFE INSURANCE.

BY LLEWELYN THOMAS, M.D.,

Physician to the Royal Academy of Music; Surgeon to the Central Throat and Ear Hospital, London.

AMONG the various, and in some instances vexatiously minute, forms of questions propounded by different Life-offices, for the guidance of their medical examiners, I believe, in very many cases, the question, "Have you ever had a discharge from the ear?" is omitted. Now, it may be considered that this subject would be included in the general question, "Have you had any complaint not already mentioned?" but, from my experience of aural patients, I am confident that very few would volunteer any information on the subject, as many people appear to consider a slight running from the ears as a normal condition; and, even if the fact were admitted, many practitioners treat otorrhœa with such contempt that they would probably pass the matter by with indifference.

I propose, therefore, to demonstrate that the subject is one of great importance in the prognosis of the probabilities of a life being "good," and suggest that, in a doubtful case, the verdict should be held in abeyance until the suspected ear has been carefully examined and tested. I can well imagine the referee objecting to make a tedious examination with the aural speculum and other tests, in addition to his multifarious duties, for the ordinary fee of a guinea, or, in some cases, for as little

as half-a-guinea; yet the subject is of the greatest importance both to the office and to the proposer.

The causes of chronic suppuration of the middle ear are acute and subacute catarrhal inflammation, and acute suppurative inflammation of the tympanum, which may result from small-pox, scarlet fever, measles, diphtheria, pneumonia, ordinary catarrh, mechanical injuries, and gouty, scrofulous, or syphilitic dyscrasia: the results are polypi and exostoses, disease of the mastoid cells, caries and necrosis, cerebral abscess, pyæmia, phlebitis, and paralysis, or even possibly insanity.

The late Mr. Toynbee, and Dr. Roosa, the talented New York aurist, have published tables, from which it may be seen that death may occur from chronic aural disease at any age and at any period from the commencement of the discharge. Mr. Toynbee cites a case, terminating fatally after the existence of otorrhœa for thirty-five years, and Dr. Roosa one after forty-one years. It is thus evident that chronic suppuration may lie dormant for years, and may then be kindled into a most fatal disease, either by some accidental circumstance, as exposure to cold, an injury to the head, a deteriorated state of health, or by the mechanical results produced by the affection itself.

A careful consideration of the anatomy of the tympanum will at once show what a dangerous locality it is in which to have a chronic suppuration as a constant inhabitant; in fact, I am acquainted with no other region of the body which is in relation with such terribly vital parts. The roof of the tympanum is in contact with the meninges of the brain; the bone is here often thin and porous, or even entirely absent, so that the tympanum actually forms a portion of the cavity of the cranium. The floor of the tympanum is sometimes bony, sometimes membranous, and, lying lower than the floor of the external auditory meatus and the orifices of the mastoid cells and Eustachian tubes, is frequently the seat of purulent accumulations in suppuration of the middle ear, which may result in phlebitis of the jugular vein. The inner wall is the outer boundary of the labyrinth, and is in front of the promontory in contact with the carotid artery; caries, in this situation, may produce hæmorrhage from the carotid, or suppuration extending through the labyrinth into the cavity of the skull. The posterior wall opens into the mastoid cells, which are bounded internally by the lateral sinus—a fact of considerable importance in suppuration of the mastoid, or in attempts to remove polypi springing from this locality. Polypi and exostoses of the tympanum, produced by the irritation of the chronic suppuration, may both prove fatal, either by pressure and caries, or by blocking up the discharge and producing meningitis, the pus in these cases rarely finding its way through the Eustachian tubes, as they are generally rendered impervious by the long-continued inflammatory processes. The mastoid cells are probably always more or less affected in a suppuration of the tympanum; but at any time a more dangerous affection may be lighted up, either acute periostitis or caries, from accumulation of the discharges, both of which frequently prove fatal from extension to the meninges, especially in cases where the gravity of the symptoms has not been early recognised and actively treated. Caries of any portion of the temporal bone may take place, and is always dangerous, as it may result in meningitis, abscess, hæmorrhage, or phlebitis. It is not at all easy to determine the existence of caries in many cases, as the probe cannot be used with anything like the same degree of boldness as in other parts. If, however minute it may be, a suppuration persistently resist treatment, caries of some portion of the bone probably exists: caries even of the external meatus, from the proximity of the dura mater to the upper wall and of the mastoid cells, may produce meningitis. Pyæmia also occurs, probably from the entrance of pus into the circulation through the mastoid veins or lateral sinus.

One of the most interesting facts in connection with the subject of chronic aural disease is its connection with cerebral abscess, for, from the published cases of German and English authors, it appears that from one-third to one-half of the cases of abscess of the cerebrum arise from aural disease. This is readily understood, if we consider the intimate connection of the dura mater with the roof of the tympanum. There is usually meningitis in connection with the site of the abscess; but sometimes the membranes are healthy, and even a portion of healthy brain-substance may intervene between the temporal bone and the abscess. Collections of pus are also not at all uncommon in the cerebellum after chronic disease of the mastoid cells, and here occurs another possible source of error in prognosis; the discharge, when investigated, may appear to come from the meatus only, as the membrane is seen to be intact, yet there may be in such a case caries of some portion of the temporal bone and disease in the mastoid cells, without perforation of the drum head. Such cases are rare, though they are on record. Apparently, no rule can be laid down regarding the danger of necrosis of different portions of the temporal bone; in many cases, a fortunate kind of hyperplasia takes place around the tympanum, and