clerk, which occupation involves long standing. He is at this time free from all appearance of swelling, which originally extended from the upper third of the thigh to the foot, and is able to walk any reasonable distance without pain, and with complete freedom of action of the foot.

NEW FORMS OF VACCINATORS.

By A. B. STEELE, L.K.Q.C.P., Member of the National Vaccine Establishment, Liverpool.

THERE is much reason to doubt whether the practical efficiency of vaccination is promoted or impeded, by the numerous attempts to complicate by ingenious mechanical contrivances a really simple though important little operation. Those who desire to witness the highest degree of uniformity and typical excellence which has probably been attained in this country in the production of Jennerian vesicles, may gratify their wishes by visiting the station of Mr. Marson, in Lambeth, whose experience and success in vaccination are probably unequalled, certainly not exceeded, anywhere. They will find that these desirable results are obtained by the use of the old-fashioned bleeding lancet, that being the instrument ordinarily used by Mr. Marson for the insertion of lymph. In attaching undue importance to the form of the instrument used, there is some fear lest the far more essential points, namely the selection of lymph, and attention to other minute details, well understood by every experienced vaccinator, should be overlooked.

The latest novelty, the new form of vaccinator by Dr. P. M. Braidwood, figured and described in the JOURNAL of the 18th May, is really the Danish knife, a most efficient little instrument, with the addition of a joint in one blade.

The only real difference between this so-called new vaccinator and the Danish knife, which has been for many years used by Mr. Ceely (by whose recommendation I was led to adopt this mode of vaccinating), consists in the jointed blade which can be drawn back for the purpose of cleaning it. This object has been attained by a simpler and, as I think, preferable contrivance, which I adopted a few years since, viz., by the construction of the Danish knife on the principle of the cross-action forceps, so that by simple pressure on the handles both blades are separated and can be readily cleaned. Although it did not appear to me sufficiently important to publish this as a new invention, I had the pleasure some two years since of showing it to Dr. Braidwood; and if I am not mistaken it was at the Liverpool station of the National Vaccine Establishment that he first witnessed the mode of vaccinating on the Danish plan, which I quite agree with him in considering very efficient and satisfactory in its results. The instrument which I use can be made by any surgical cutler; it is kept in stock by Wood, surgical instrument maker, Liverpool, and will probably be less costly, a manifest recommendation when efficiency is not sacrificed to economy.

FORCIBLE FEEDING OF THE INSANE.

By H. SUTHERLAND, M.B., M.R.C.P., Blacklands House, Chelsea.

IN the JOURNAL of March 30th, there appears a paper on "Forcible Feeding of the Insane", in which the writer states that he believes Paley's apparatus will be found efficient for feeding even in the severest cases of refusal of food. In the most severe cases that have come under my notice, Paley's feeder was not of the slightest use, although, doubtless, a valuable instrument in the subacute forms of insanity where food is refused. At the West Riding Lunatic Asylum, where there were sometimes as many as four patients to be fed by artificial means three times a day, endeavours were always made first to induce them to take their food naturally, feeding themselves. If that failed, an attendant would try to feed them, as one would feed a child or sick person, with a spoon. Sometimes an attendant's wife would persuade a male patient to take his food; and *vice versa*, the physician would often succeed with a female patient when the nurse had failed. If the patient still resisted, Dr. Williams's method of feeding with the spoon was tried, and, if that again proved a failure, the œsophageal tube was usually resorted to.

In the milder cases of refusal of food, the spoon, or Paley's feeder, will often answer the purpose very well; but in the severest cases, what are we to do? The severest cases that I have met with are those of acute mania, occurring in the course of general paralysis. It may require five men to get such a patient properly into position; but if Dr. Williams's method be adopted, three ought to be sufficient to hold him while he is fed. His method of holding the patient is as follows. One at-

tendant kneels at the head of the bedstead, with a knee on each side of the patient's head, which should be slightly raised; care being taken to put a soft towel between the knees and the patient's ears, which otherwise might be injured. This attendant also holds the patient's head with his hand. A sheet is thrown across the patient's body, and drawn down tightly upon him, especially at the knees. Two attendants then kneel on the sheet, one on each side of the patient's knees, so that his legs are held tightly in their place without the least risk of injury. Each of the attendants then grasps one of the patient's arms, which should be left outside the sheet. The patient is thus held firmly, and without danger, by the three attendants. If there be more attendants, they are only in one another's way, and interfere with the movements of the medical man. The physician then opens the patient's mouth with the ordinary screw-gag, which the attendant holding the head should look after and keep in position. The tube is then passed. It is first dipped into the basin of liquid food, which is of course ready at hand and properly warmed. The tube may require to be slightly bent, about two inches from the end in some cases. The physician then inserts the tube, and along with it his forefinger, with which he feels for the epiglottis, slides it over it, and pushes about eighteen inches of the tube down the œsophagus. If this length of tubing go down without any grating or stoppage of any kind, he may be sure he has not passed his tube into the larynx. The food is then injected by the stomach-pump. I should mention here that I have never seen the liquid food pass down the sides of the tube. It is true the patient may vomit the food; and, if this occur, he should be immediately made to sit up, and the tube should be at once removed. When the food has all been pumped in, the tube should be withdrawn, and the basin placed under the chin of the patient, to prevent the liquid which remains in the tube from running over the bedclothes.

A far better apparatus than any mentioned above is a large tin-funnel, to the narrow end of which is attached an India-rubber-tube about 18 inches long, the other end of which fits into the stomach-tube, which should be about two feet in length and five-eighths of an inch in diameter. When the tube is fairly passed, the basin of liquid food is simply emptied into the funnel and finds its way into the stomach by the force of gravity.

Now, the reason for adopting the above method, instead of using Paley's feeder, in severe cases of acute mania, is this: that these patients will often holla and shout all the time they are being fed, so that it is impossible to prevent food, introduced by Paley's feeder, from entering the larynx, which of course may be followed by serious consequences.

With regard to the indications for using the spoon, Paley's feeder, or the nasal tube, instead of the stomach-pump, I think such methods are to be adopted in the milder cases requiring forcible feeding, or when it is wished to make the process as disagreeable as possible, as in the case of lazy patients who refuse food merely by force of habit. But in the severest cases, there is no method equal to the stomach-tube, both for making a certainty of the patient getting his proper amount of food, and for providing for his safety more effectually than by any other system of artificial feeding.

OBSTETRIC MEMORANDA.

PREMATURE RUPTURE OF THE MEMBRANES.

WITHIN the last month or two the consequences of this accident have been discussed in the JOURNAL, and opinions upon the subject have been so various, and even conflicting, that I had some weeks ago nearly resolved to give the facts of a case which presented itself many years since in my practice. The case left a strong impression upon my mind; and, as through a period of forty years I have not had a similar one, I now report it as completely supporting the experience and views of Dr. Thorburn, of Manchester, who reproduces the subject in the JOURNAL of May 18th, and whose third case exactly resembles mine in respect of the duration of gestation after the escape of the liquor amnii, while it shows at the same time that a term of six weeks' gestation after the accident is not incompatible with the safety and health of the foetus.

In the middle of January, 1836, I was called to a woman, aged 27, in her second labour, she having, as she stated, passed her seventh month of gestation. A regular course of pains soon completed the first stage, when the membranes suddenly ruptured, disclosing that the vertex was presented naturally—a fact which I could easily make out by digital examination. The point of my finger was distinctly in contact with the sutures, thus leading one to the conclusion that true liquor amnii had been discharged. I waited through the day in the expectation of speedy delivery, but, like Dr. Thorburn, I got tired of