paralysis or excitement. For the three following days she appeared better during the day, but at night-time was submaniacal, noisy, and uncontrollable; and on the fourth night I was summoned to the infirmary, at 11 P.M., when I found the patient, who, after returning from a walk, had seated herself in a chair and commenced screaming. She had refused her food, and had refused to move or to speak, and was in a submaniacal condition. After a short time, I succeeded in persuading her to rise, and then, with assistance, to go upstairs. She was put to bed, and I then examined her, and found that she was unable to speak, though she endeavoured to answer "yes". The excitement appeared to have moderated considerably, and I ordered her an increased dose of belladonna. She slept well, but the next day complained of headache, and her face was covered with petechiæ. I then ordered that her belladonna should be taken every six hours; and the treatment was persevered in with an intermission of two days only, when she suffered from bilious vomiting, for three weeks, when she was discharged, not having had any return of the symptoms, and having been under treatment over three months. I recommended that the treatment should be continued for twelve months, and that she should use a tepid bath every morning. Upon the application of the friends, I consented to readmit the patient should her malady return; but as I have not had a renewal of the application, the conclusion is favourable for the patient's state.

The case is one of both value and significance in illustrating the frequent consequential mental excitement of epilepsy, particularly in that class of cases to which I have several times drawn attention, where the epilepsy is slight, sometimes so slight as hardly to be observable, and wherein the mental disturbance seems to bear a sort of inverse ratio to the epilepsy; the epilepsy often being expressed as a very transient loss of consciousness, whilst the mental disturbance consequent upon it is profound.

The medico-legal aspect of these cases is in the greatest degree of importance, since the responsibility of a person committing a criminal act under the influence of the fleeting mania following epilepsy, is or ought to be annulled. The petechiæ I have frequently commented upon, and it is perhaps unnecessary to enter upon the subject here. I may add that the treatment with belladonna was one which I frequently adopt, and with great success, when there is no indication for the use of absorbents, and especially when the uterine functions appear in any way disturbed.

MASKED SMALL-POX: PETECHIÆ: RAPIDLY FATAL.

By FREDERICK PAGE, M.D., Senior House-Surgeon to the Infirmary, Newcastle-on-Tyne.

A. S., a comedian, aged 23, applied at the Newcastle-on-Tyne Infirmary during the afternoon of Monday, April 22nd, complaining of a very acute pain over his left lumbar muscles. He had fallen down two or three stairs upon his back. He was of intemperate habits, and acknowledged that he had been drinking. The pain was spasmodic. Upon a careful examination, no indication of a bruise could be found. There had been no sickness, and the urine was stated to be as usual. Two or three ounces of blood were taken from the loin by cupping, with marked relief; a calomel and jalap purge was administered, and the patient was told to come back on the following day if the pain returned. The next morning he again applied, and had a hypodermic injection of a quarter of a grain of morphia, after which he walked home. But little relief and no sleep followed the injection. A man who slept in the same room said that he noticed no rash on his companion when they went to bed, and nothing had occurred to give rise to the idea of imminent danger. After a restless night, the patient, between three and four in the morning, suddenly died.

As I was unable to give a certificate as to the cause of death, having seen nothing in the man's condition when I examined him to indicate danger to life, an inquest was asked for. The necropsy was made on Thursday, April 25th, at 1 P.M., at the Infirmary, in the presence of the physicians and some of the surgeons. The body was in fair condition. Rigor mortis was absent. On the left arm were two vaccination-marks. Around the puncture made by the injection-needle was a large effusion of dark fluid blood. No blood was poured out where the cupping-glass had been applied. The trunk of the body was suffused with a small thickly-set petechial rash of mulberry colour. The thighs were copiously, the legs and arms sparsely, and the face but slightly, if at all, sprinkled with the same kind of eruption. Both conjunctivæ were completely ecchymosed. The face was livid, and the body generally, but more particularly upon the left side, deeply discoloured

by suffigation. Upon opening the chest, the muscles were found to be very soft and dark coloured. A few old strings of false membrane existed between the pleuræ on both sides, and in each pleural cavity was a quantity of thin bloody fluid. The lungs were much congested, and their surface was sprinkled extensively with stellate petechiæ. The heart was rather small, flaccid, and spotted with petechiæ; the valves were all normal. The liver was similarly affected with petechiæ over its surface, and the gall-bladder was full of bile. The intestinal canal was very vascular and dark-coloured generally, petechiæ and large patches of extravasated blood appearing between the layers of the mesentery and beneath the peritoneum in all directions. Externally both kidneys were blotched with numerous bloody effusions, and into the fat surrounding them blood was also poured out. The bladder contained a few ounces of bloody urine, and its mucous surface was deeply stained with blood. The brain was covered with minute arborescences of dark vessels, but no petechiæ or any effusions of blood were seen, either on the surface or in the interior of the organ. The brain-substance seemed sound, perhaps rather soft. The blood was throughout the body unusually dark and in a very fluid state.

It was the universal opinion of those present at the necropsy that the man had died from malignant small-pox. During the epidemic of small-pox, which has raged here for twelve months, and which has now subsided, occasional cases strongly resembling the above have occurred, proving fatal in a few days, where no eruption whatever of small-pox appeared, but where petechiæ, varying in size and number, spread over the whole surface of the body, with great impairment of the muscular power, but with perfect integrity of mind and unconsciousness of danger. Such cases, too, have not always occurred in the intemperate.

This case is certainly remarkable, inasmuch as the powers of the nervous centres and of the muscles were alike preserved, and no eruption whatever appeared up to at least four or five hours before death. It will have been noted, this man was quite able to go about, and did so to the last, and that the only symptoms he had was the pain in the loins, not over the spine, but in the left lumbar muscles. Intemperance may have rendered him an easier prey to infection, and the bad consequences of poisoned blood. In the JOURNAL (page 497) another extraordinary case of malignant small-pox masked by injury is reported by Mr. Gascoven, of St. Mary's and the Lock Hospitals; and reading it has induced me to send this.

OBSTETRIC MEMORANDA.

ALBUMINURIA IN THE SIXTH MONTH OF PREGNANCY.

M. W., a multipara, aged 32, a laundress, slipped and fell, without unconsciousness, in a steep garden, on the 1st February. On the 3rd, I found the uterus gravid and prolapsed, with the os just within the vulva; the patient stated it had been down since the fall. She supposed herself to be between the third and fourth months of pregnancy, her last menstruation having been in the middle of October. The uterus was easily replaced, and the patient was ordered to keep in bed; but, on the 6th February, she got up, reproducing the prolapse. On the 7th, I applied a flattened caoutchouc pessary, which kept the uterus *in situ* until a week after quickening (which was on the 15th), when the pessary slipped down and was withdrawn.

I was summoned on the 23rd March, and found that the patient had acute and general anasarca, constipation with severe headache, dry and harsh skin, and was passing a very small quantity of urine; hot-air baths, diaphoretics, and saline purgatives were ordered; these caused profuse diaphoresis and watery stools. On the 24th, the urine became solid when heated; no tube-casts or blood-corpuscles were visible in it; the fœtus moved until this day. The patient was much better on the 26th, passing more urine, which was less albuminous. On the 6th April, she commenced taking ten minims of the tincture of the perchloride of iron with twenty minims of nitrous ether four times daily. The patient did not have any convulsion, and in a week the albuminuria had ceased. Labour began on the 29th, and lasted five-and-a-half hours, resulting in the birth of a putrid female fœtus of between five and six months' development. On the 30th April and 1st May, the patient was convalescing favourably. I was unable to attend the labour, which was conducted by a nurse; no chloroform was given. The first pregnancy terminated in the birth of an eight months' fœtus, six weeks after its death; the second in the birth of a girl, and the third in that of a boy (both living now); the fourth in a miscarriage in the third month, three years ago; this last being the fifth pregnancy. This case supports that published at page 447 of the JOURNAL for April 27th.

STANLEY HAYNES, M.D., Malvern Link.